

# Coloring $k$ -partite sparse digraphs

Ararat Harutyunyan<sup>a</sup>, Louisa Harutyunyan<sup>b</sup>, Narek Hovhannisyan<sup>c</sup>

<sup>a</sup>*Université Paris-Dauphine, PSL Research University, CNRS, UMR 7243, LAMSADE*

<sup>b</sup>*University of Toronto Mississauga, Department of Computer Science*

<sup>c</sup>*Concordia University, Department of Computer Science and Software Engineering*

---

## Abstract

In this paper, we study the colorings of  $k$ -partite sparse digraphs. The *chromatic number* of a graph  $G$  is the smallest integer  $k$  such that the vertices of  $G$  can be colored with  $k$  colors such that each color class is an independent set. The *dichromatic number* of a digraph  $D$  is the minimum  $k$  such that the vertices of  $D$  can be colored with  $k$  colors with each color class inducing an acyclic subdigraph. This coloring invariant shares many similarities with the graph chromatic number and can be thought of as its analogous digraph generalisation.

Our main result in this short note shows that there exist sparse  $k$ -partite digraphs which have dichromatic number  $k$ . This, in particular, not only implies that there exist graphs with equal chromatic and dichromatic number, but that they can be taken to be somewhat sparse.

*Keywords:* Graph coloring, Dichromatic number,  $k$ -partite digraphs

---

---

*Email addresses:* [ararat.harutyunyan@dauphine.fr](mailto:ararat.harutyunyan@dauphine.fr) (Ararat Harutyunyan ),  
[louisa.harutyunyan@utoronto.ca](mailto:louisa.harutyunyan@utoronto.ca) (Louisa Harutyunyan), [narekh98@gmail.com](mailto:narekh98@gmail.com) (Narek Hovhannisyan)

The research of A. Harutyunyan is supported by ANR grant DAGDigDec (DAGs and Digraph Decompositions)21-CE48-0012

# 1 Introduction

The digraphs considered in this paper will be oriented graphs, i.e., not containing loops, parallel arcs or directed cycles of length 2. A subset  $S$  of vertices of a digraph  $D$  is called *acyclic* if the induced subdigraph on  $S$  contains no directed cycle. The *dichromatic number*  $\vec{\chi}(D)$  of  $D$  is the smallest integer  $k$  such that  $V(D)$  can be partitioned into  $k$  sets  $V_1, \dots, V_k$  where each  $V_i$  is acyclic. Note that, equivalently, the dichromatic number is the smallest integer  $k$ , such that the vertices of  $D$  can be colored with  $k$  colors so that there is no monochromatic directed cycle. It is easy to see that, for any undirected graph  $G$  and its bidirected digraph  $D$  obtained from  $G$  by replacing each edge by two oppositely oriented arcs, we have  $\chi(G) = \vec{\chi}(D)$ . The dichromatic number was first introduced by Neumann-Lara [7]. In recent years, there has been considerable attention devoted to this topic, and many results have demonstrated that this digraph invariant generalizes many results on the graph chromatic number (see, for example [1, 2, 5, 4]).

For an undirected graph  $G$ , the *dichromatic number* of  $G$ , denoted by  $\vec{\chi}(G)$ , is the maximum dichromatic number over all its orientations.

In 1970s, Erdős and Neumann-Lara conjectured the following:

**Conjecture 1.1.** ([3]) *For every integer  $k$ , there exists an integer  $f_k$  such that if  $G$  is any graph satisfying  $\chi(G) \geq f_k$ , then  $\vec{\chi}(G) \geq k$ .*

It is clear that  $f_1$  and  $f_2$  exist ( $f_2 = 3$ ). It is unknown if  $f_3$  exists.

A graph  $G$  is  $k$ -partite if  $V(G)$  can be partitioned into  $k$  independent sets, i.e.,  $\chi(G) \leq k$ . For two positive integer  $n$  and  $k$ , let  $K_{n*k}$  denote the complete  $k$ -partite graph with  $n$  vertices in each vertex class. The main result of this paper is to show that the conjecture of Erdős and Neumann-Lara holds for very sparse subgraphs of  $K_{n*k}$  in the following strong form.

**Theorem 1.2.** *Let  $k \geq 2$ . There is an integer  $n_0$  such that for any  $n \geq n_0$  the following holds. Let  $G_{n,k}$  be any  $k$ -partite graph with vertex classes  $V_1, \dots, V_k$  with  $|V_i| = n$  for all  $i \in [k]$  satisfying the property that for all distinct pairs  $i, j$  and all sets  $V'_i \subset V_i$  and  $V'_j \subset V_j$  with  $|V'_i| \geq n/k^2$  and  $|V'_j| \geq n/k^2$ , the number of edges between  $V'_i$  and  $V'_j$  is at least  $100n \log k/k^2$ . Then  $\vec{\chi}(G_{n*k}) = k$ .*

**Remark 1.3.** *We note that a random subgraph of  $K_{n*k}$  where the probability of an edge between any two vertices in different partite sets is at least  $f(n)/n$ , where  $f(n)$  is a function increasing arbitrarily slowly, satisfies the condition of Theorem 1.2.*

## 2 Proof of Theorem 1.2

In this section, we prove Theorem 1.2. First, let us recall the following result from Manber and Tompa [6].

Let  $S$  be the set of all graphs with  $n$  vertices and  $m$  edges and let  $A(m, n)$  denote the maximum number of acyclic orientations that a graph in  $S$  can have.

**Theorem 2.1** ([6]).  $A(m, n) \leq \prod_{v \in V} (1 + \deg(v)) \leq \left(\frac{2m}{n} + 1\right)^n$ .

**Lemma 2.2.** *Let  $i$  and  $j$  be distinct, and let  $G_{ij} = G_{n*k}[V_i \cup V_j]$  be the bipartite graph induced on vertices  $V_i$  and  $V_j$ . There is an orientation of the edges of  $G_{ij}$  such that the resulting digraph  $D$  satisfies the property that for every  $V'_i \subset V_i$  and  $V'_j \subset V_j$  verifying  $|V'_i|, |V'_j| \geq n/k^2$ , the digraph  $D' = D[V'_i \cup V'_j]$  has a directed cycle.*

*Proof.* We employ the probabilistic method. Let  $D$  be the digraph obtained from  $G_{ij}$  by taking a uniformly random orientation of its edges.

Let  $V'_i \subset V_i$  and  $V'_j \subset V_j$  be two fixed sets with  $|V'_i| = |V'_j| = \frac{n}{k^2}$ . Let  $D' = (V'_i \cup V'_j, E)$  be the digraph induced on  $V'_i$  and  $V'_j$ . Let  $m$  be the number of edges of  $D'$  and recall that  $m \geq 100n \log k/k^2$ .

Now, using Theorem 2.1, we obtain that

$$\begin{aligned} \mathbb{P}[\exists V'_i, V'_j : D[V'_i \cup V'_j] \text{ is acyclic}] &\leq \binom{n}{n/k^2}^2 \frac{\left(\frac{2m}{2n/k^2} + 1\right)^{2n/k^2}}{2^m} \\ &\leq (ek^2)^{2n/k^2} \frac{\left(\frac{100n \log k/k^2}{n/k^2} + 1\right)^{2n/k^2}}{2^{100n \log k/k^2}} \\ &\leq (ek^2)^{2n/k^2} \frac{(100 \log k + 1)^{2n/k^2}}{k^{100n/k^2}} \\ &\leq \left(\frac{ek^2(100 \log k + 1)}{k^{50}}\right)^{2n/k^2} < 1 \end{aligned}$$

This completes the proof. □

Now we finish the proof.

*Proof of Theorem 1.2.* Let  $D$  be the digraph obtained by taking an orientation of  $G_{n*k}$  such that for every pair  $i, j$  the conclusion of Lemma 2.2 holds for  $G_{ij}$ . We will show that  $[\vec{\chi}(D) = k]$ .

Assume  $D$  is colored with any number of colors. We can say that if there exist two partitions  $V_i$  and  $V_j$  and subsets of vertices  $V'_i \subset V_i$  and  $V'_j \subset V_j$  of the same color, with  $|V'_i|, |V'_j| \geq n/k^2$ , then there is a monochromatic directed cycle. Indeed, this follows from the aforementioned lemma.

Note that  $\vec{\chi}(D) \leq \chi(G_{n*k}) \leq k$ .

Now let us prove that  $\vec{\chi}(D) = k$  by contradiction. Assume it is possible to color  $D$  with  $k - 1$  colors. As  $D$  is validly colored with  $k - 1$  colors it means that there cannot be two sets of the same color in different partitions verifying  $|V'_i|, |V'_j| \geq \frac{n}{k^2}$ .

If there are  $k - 1$  colors, then there is at least one color (hereupon called *dominant color*) so that there are at least  $\frac{k \cdot n}{k-1}$  vertices colored with that color. Let  $(c_1, c_2, \dots, c_k)$  be the number of vertices that are assigned the dominant color in  $(V_1, V_2, \dots, V_k)$  respectively.

As mentioned above there cannot be  $c_i$  and  $c_j$  such that both are at least  $\frac{n}{k^2}$ . In the best case scenario only one of them can be more than  $\frac{n}{k^2}$ . Without loss of generality assume that  $c_k \geq \frac{n}{k^2}$ . Obviously, the following inequality should hold.

$$\sum_{i=1}^k c_i \geq \frac{k \cdot n}{k-1} \quad (1)$$

Moreover,

$$\sum_{i=1}^k c_i = \sum_{i=1}^{k-1} c_i + c_k \leq (k-1) \cdot \frac{n}{k^2} + c_k$$

If Equation 1 is valid, then the following should hold too.

$$(k-1) \cdot \frac{n}{k^2} + c_k \geq \frac{k \cdot n}{k-1}$$

$$(k-1) \cdot \frac{n}{k^2} \geq \frac{k \cdot n}{k-1} - c_k$$

In order to minimize the right-hand side, we select the maximum possible value for  $c_k$ , which is  $\max(c_k) = |V_k| = n$ .

$$(k-1) \cdot \frac{n}{k^2} \geq \frac{k \cdot n}{k-1} - n = \frac{n}{k-1}$$

As  $k$  is a constant and  $n$  is any number satisfying  $n \geq n_0$ , it is obvious that the aforementioned inequality is invalid for any  $n$ . Thus, there necessarily exist at least two partite sets with more than  $\frac{n}{k^2}$  vertices of the dominant color, which means with probability  $1 - o(1)$  there will exist a monochromatic cycle.

Thus,  $\vec{\chi}(D) = k$ . □

**Corollary 2.3.** *For every  $k$ , there is a simple graph  $G$  such that  $\chi(G) = \vec{\chi}(G) = k$ .*

## References

- [1] Julien Bensmail, Ararat Harutyunyan, and Ngoc Khang Le. List coloring digraphs. *Journal of Graph Theory*, 87(4):492–508, 2018.
- [2] Drago Bokal, Gasper Fijavz, Martin Juvan, P Mark Kayll, and Bojan Mohar. The circular chromatic number of a digraph. *Journal of Graph Theory*, 46(3):227–240, 2004.
- [3] Paul Erdős. Problems and results in number theory and graph theory. *Proceedings of Ninth Manitoba Conference on Numerical Math. and Computing*, pages 3–21, 1979. [https://www.renyi.hu/~p\\_erdos/1980-30.pdf](https://www.renyi.hu/~p_erdos/1980-30.pdf).
- [4] Ararat Harutyunyan and Bojan Mohar. Gallai’s theorem for list coloring of digraphs. *SIAM Journal on Discrete mathematics*, 25(1):170–180, 2011.
- [5] Ararat Harutyunyan and Bojan Mohar. Two results on the digraph chromatic number. *Discrete mathematics*, 312(10):1823–1826, 2012.
- [6] Udi Manber and Martin Tompa. The effect of number of hamiltonian paths on the complexity of a vertex-coloring problem. *SIAM Journal on Computing*, 13(1):109–115, 1984.
- [7] Victor Neumann-Lara. The dichromatic number of a digraph. *Journal of Combinatorial Theory, Series B*, 33(3):265–270, 1982.