

Cooperative Games

Lecture 4: The Bargaining Set

Stéphane Airiau

ILLC - University of Amsterdam



Today

- If agents desire the kind of stability offered by the core, they will be unable to reach an agreement.
- ☞ they have no choice but to **relax** their stability requirements.
- We would like a solution that allows agents to always reach an agreement, while guaranteeing some stability.
- The bargaining set is one such solution.

A second solution concept:

The bargaining set.

R.J. Aumann and M. Maschler. **The bargaining set for cooperative games**, in *Advances in game theory (Annals of mathematics study)*, 1964.

M. Davis and M. Maschler. **Existence of stable payoff configurations for cooperative games**, *Bulletin of the American mathematical society*, 1963.

Let (N, v, \mathcal{S}) be a game with coalition structure and x an imputation.

The bargaining set models stability in the following sense:

Any **argument** from an agent i against a payoff distribution x is of the following form:

I get too little in the imputation x , and agent j gets too much! I can form a coalition that excludes j in which some members benefit and all members are at least as well off as in x .

The argument is **ineffective** for the bargaining set if agent j can answer the following:

I can form a coalition that excludes agent i in which all agents are at least as well off as in x , and as well off as in the payoff proposed by i for those who were offered to join i in the argument.

Definition (Objection)

Let (N, v, \mathcal{S}) be a game with coalition structure, $x \in X_{(N, v, \mathcal{S})}$ (the set of all feasible payoff vectors for (N, v, \mathcal{S})), $\mathcal{C} \in \mathcal{S}$ be a coalition, and i and j two distinct members of \mathcal{C} ($(i, j) \in \mathcal{C}^2$, $i \neq j$).

An **objection of i against j** is a pair (P, y) where

- $P \subseteq N$ is a coalition such that $i \in P$ and $j \notin P$.
- $y \in \mathbb{R}^p$ where p is the size of P
- $y(P) \leq v(P)$ (y is a feasible payoff distribution for the agents in P)
- $\forall k \in P$, $y_k \geq x_k$ and $y_i > x_i$ (agent i strictly benefits from y , and the other members of P do not do worse in y than in x .)

An objection (P, y) of i against j is a **potential threat** by coalition P , which contains i but not j , to deviate from x . The goal is not to change \mathcal{S} , but to obtain a side payment from j to i , i.e., to modify x within $X_{(N, v, \mathcal{S})}$.

Definition (Counter-objection)

An **counter-objection to** (P, y) is a pair (Q, z) where

- $Q \subseteq N$ is a coalition such that $j \in Q$ and $i \notin Q$.
- $z \in \mathbb{R}^q$ where q is the size of Q
- $z(Q) \leq v(Q)$ (z is a feasible payoff distribution for the agents in Q)
- $\forall k \in Q, z_k \geq x_k$ (the members of Q get at least the value in x)
- $\forall k \in Q \cap P, z_k \geq y_k$ (the members of Q which are also members of P get at least the value promised in the objection)

In a counter-objection, agent j must show that it can protect its payoff x_j in spite of the existing objection of i .

Definition (Stability)

Let (N, v, \mathcal{S}) a game with coalition structure. A vector $x \in X_{(N, v, \mathcal{S})}$ is **stable** iff for each objection at x there is a counter-objection.

Definition (Pre-bargaining set)

The **pre-bargaining set** (*preBS*) is the set of all stable members of $X_{(N, v, \mathcal{S})}$.

Lemma

Let (N, v, \mathcal{S}) a game with coalition structure, we have
$$\text{Core}(N, v, \mathcal{S}) \subseteq \text{preBS}(N, v, \mathcal{S}).$$

This is true since, if $x \in \text{Core}(N, v, \mathcal{S})$, no agent i has any objection against any other agent j .

Example

Let (N, v) be a 7-player simple majority game, i.e.

$$v(\mathcal{C}) = \begin{cases} 1 & \text{if } |\mathcal{C}| \geq 4 \\ 0 & \text{otherwise} \end{cases}.$$

Let us consider $x = \langle -\frac{1}{5}, \frac{1}{5}, \dots, \frac{1}{5} \rangle$. It is clear that $x(N) = 1$.

Let us prove that **x is in the pre-bargaining set** of the game $(N, v, \{N\})$.

Objections within members of $\{2, 3, 4, 5, 6, 7\}$ will have a counter-objection by symmetry. ✓

Example

Let (N, v) be a 7-player simple majority game, i.e.

$$v(\mathcal{C}) = \begin{cases} 1 & \text{if } |\mathcal{C}| \geq 4 \\ 0 & \text{otherwise} \end{cases}.$$

Let us consider $x = \langle -\frac{1}{5}, \frac{1}{5}, \dots, \frac{1}{5} \rangle$. It is clear that $x(N) = 1$.

Let us prove that **x is in the pre-bargaining set** of the game $(N, v, \{N\})$.

Objections within members of $\{2, 3, 4, 5, 6, 7\}$ will have a counter-objection by symmetry. ✓

Let us consider the objections (P, y) of 1 against another member of $\{2, 3, 4, 5, 6, 7\}$. Since the players $\{2, \dots, 7\}$ play symmetric roles, we consider an objection (P, y) of 1 against 7 using successively as P $\{1, 2, 3, 4, 5, 6\}$, $\{1, 2, 3, 4, 5\}$, $\{1, 2, 3, 4\}$, $\{1, 2, 3\}$, $\{1, 2\}$ and $\{1\}$. We will look for a counter-objection of player 7 using (Q, z) .

- $P = \{1, 2, 3, 4, 5, 6\}$. We need to find the payoff vector $y \in \mathbb{R}^6$ so that (P, y) is an objection.

- $P = \{1, 2, 3, 4, 5, 6\}$. We need to find the payoff vector $y \in \mathbb{R}^6$ so that (P, y) is an objection. $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \dots, \frac{1}{5} + \alpha_6 \rangle$.

- $P = \{1, 2, 3, 4, 5, 6\}$. We need to find the payoff vector $y \in \mathbb{R}^6$ so that (P, y) is an objection. $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \dots, \frac{1}{5} + \alpha_6 \rangle$.

The conditions for (P, y) to be an objection are the following:

- each agent is as well off as in x : $\alpha > -\frac{1}{5}, \alpha_i \geq 0$
- y is feasible for coalition P : $\sum_{i=2}^6 \left(\alpha_i + \frac{1}{5} \right) + \alpha \leq 1$.

- $P = \{1, 2, 3, 4, 5, 6\}$. We need to find the payoff vector $y \in \mathbb{R}^6$ so that (P, y) is an objection. $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \dots, \frac{1}{5} + \alpha_6 \rangle$.

The conditions for (P, y) to be an objection are the following:

- each agent is as well off as in x : $\alpha > -\frac{1}{5}, \alpha_i \geq 0$
- y is feasible for coalition P : $\sum_{i=2}^6 \left(\alpha_i + \frac{1}{5} \right) + \alpha \leq 1$.

w.l.o.g $0 \leq \alpha_2 \leq \alpha_3 \leq \alpha_4 \leq \alpha_5 \leq \alpha_6$.

- $P = \{1, 2, 3, 4, 5, 6\}$. We need to find the payoff vector $y \in \mathbb{R}^6$ so that (P, y) is an objection. $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \dots, \frac{1}{5} + \alpha_6 \rangle$.

The conditions for (P, y) to be an objection are the following:

- each agent is as well off as in x : $\alpha > -\frac{1}{5}$, $\alpha_i \geq 0$
- y is feasible for coalition P : $\sum_{i=2}^6 \left(\alpha_i + \frac{1}{5} \right) + \alpha \leq 1$.

w.l.o.g $0 \leq \alpha_2 \leq \alpha_3 \leq \alpha_4 \leq \alpha_5 \leq \alpha_6$.

$$\text{Then } \sum_{i=2}^6 \left(\frac{1}{5} + \alpha_i \right) + \alpha = \frac{5}{5} + \sum_{i=2}^6 \alpha_i + \alpha = 1 + \sum_{i=2}^6 \alpha_i + \alpha \leq 1.$$

- $P = \{1, 2, 3, 4, 5, 6\}$. We need to find the payoff vector $y \in \mathbb{R}^6$ so that (P, y) is an objection. $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \dots, \frac{1}{5} + \alpha_6 \rangle$.

The conditions for (P, y) to be an objection are the following:

- each agent is as well off as in x : $\alpha > -\frac{1}{5}$, $\alpha_i \geq 0$
- y is feasible for coalition P : $\sum_{i=2}^6 \left(\alpha_i + \frac{1}{5} \right) + \alpha \leq 1$.

w.l.o.g $0 \leq \alpha_2 \leq \alpha_3 \leq \alpha_4 \leq \alpha_5 \leq \alpha_6$.

$$\text{Then } \sum_{i=2}^6 \left(\frac{1}{5} + \alpha_i \right) + \alpha = \frac{5}{5} + \sum_{i=2}^6 \alpha_i + \alpha = 1 + \sum_{i=2}^6 \alpha_i + \alpha \leq 1.$$

$$\text{Then } \sum_{i=2}^6 \alpha_i \leq -\alpha < \frac{1}{5}.$$

- $P = \{1, 2, 3, 4, 5, 6\}$. We need to find the payoff vector $y \in \mathbb{R}^6$ so that (P, y) is an objection. $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \dots, \frac{1}{5} + \alpha_6 \rangle$.

The conditions for (P, y) to be an objection are the following:

- each agent is as well off as in x : $\alpha > -\frac{1}{5}$, $\alpha_i \geq 0$
- y is feasible for coalition P : $\sum_{i=2}^6 \left(\alpha_i + \frac{1}{5} \right) + \alpha \leq 1$.

w.l.o.g $0 \leq \alpha_2 \leq \alpha_3 \leq \alpha_4 \leq \alpha_5 \leq \alpha_6$.

$$\text{Then } \sum_{i=2}^6 \left(\frac{1}{5} + \alpha_i \right) + \alpha = \frac{5}{5} + \sum_{i=2}^6 \alpha_i + \alpha = 1 + \sum_{i=2}^6 \alpha_i + \alpha \leq 1.$$

$$\text{Then } \sum_{i=2}^6 \alpha_i \leq -\alpha < \frac{1}{5}.$$

- We need to find a counter-objection for (P, y) .

- $P = \{1, 2, 3, 4, 5, 6\}$. We need to find the payoff vector $y \in \mathbb{R}^6$ so that (P, y) is an objection. $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \dots, \frac{1}{5} + \alpha_6 \rangle$.

The conditions for (P, y) to be an objection are the following:

- each agent is as well off as in x : $\alpha > -\frac{1}{5}$, $\alpha_i \geq 0$
- y is feasible for coalition P : $\sum_{i=2}^6 \left(\alpha_i + \frac{1}{5} \right) + \alpha \leq 1$.

w.l.o.g $0 \leq \alpha_2 \leq \alpha_3 \leq \alpha_4 \leq \alpha_5 \leq \alpha_6$.

$$\text{Then } \sum_{i=2}^6 \left(\frac{1}{5} + \alpha_i \right) + \alpha = \frac{5}{5} + \sum_{i=2}^6 \alpha_i + \alpha = 1 + \sum_{i=2}^6 \alpha_i + \alpha \leq 1.$$

$$\text{Then } \sum_{i=2}^6 \alpha_i \leq -\alpha < \frac{1}{5}.$$

- We need to find a counter-objection for (P, y) .

claim: we can choose $Q = \{2, 3, 4, 7\}$ and

$$z = \langle \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5} + \alpha_4, \frac{1}{5} + \alpha_5 \rangle$$

- $P = \{1, 2, 3, 4, 5, 6\}$. We need to find the payoff vector $y \in \mathbb{R}^6$ so that (P, y) is an objection. $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \dots, \frac{1}{5} + \alpha_6 \rangle$.

The conditions for (P, y) to be an objection are the following:

- each agent is as well off as in x : $\alpha > -\frac{1}{5}$, $\alpha_i \geq 0$
- y is feasible for coalition P : $\sum_{i=2}^6 \left(\alpha_i + \frac{1}{5} \right) + \alpha \leq 1$.

w.l.o.g $0 \leq \alpha_2 \leq \alpha_3 \leq \alpha_4 \leq \alpha_5 \leq \alpha_6$.

$$\text{Then } \sum_{i=2}^6 \left(\frac{1}{5} + \alpha_i \right) + \alpha = \frac{5}{5} + \sum_{i=2}^6 \alpha_i + \alpha = 1 + \sum_{i=2}^6 \alpha_i + \alpha \leq 1.$$

$$\text{Then } \sum_{i=2}^6 \alpha_i \leq -\alpha < \frac{1}{5}.$$

- We need to find a counter-objection for (P, y) .

claim: we can choose $Q = \{2, 3, 4, 5\}$ and

$$z = \langle \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5} + \alpha_4, \frac{1}{5} + \alpha_5 \rangle$$

$$z(Q) = \frac{1}{5} + \alpha_2 + \frac{1}{5} + \alpha_3 + \frac{1}{5} + \alpha_4 + \frac{1}{5} + \alpha_5 = \frac{4}{5} + \sum_{i=2}^5 \alpha_i \leq 1 \text{ since } \sum_{i=2}^5 \alpha_i \leq \sum_{i=2}^6 \alpha_i < \frac{1}{5} \text{ so } z \text{ is feasible.}$$

- $P = \{1, 2, 3, 4, 5, 6\}$. We need to find the payoff vector $y \in \mathbb{R}^6$ so that (P, y) is an objection. $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \dots, \frac{1}{5} + \alpha_6 \rangle$.

The conditions for (P, y) to be an objection are the following:

- each agent is as well off as in x : $\alpha > -\frac{1}{5}, \alpha_i \geq 0$
- y is feasible for coalition P : $\sum_{i=2}^6 \left(\alpha_i + \frac{1}{5} \right) + \alpha \leq 1$.

w.l.o.g $0 \leq \alpha_2 \leq \alpha_3 \leq \alpha_4 \leq \alpha_5 \leq \alpha_6$.

$$\text{Then } \sum_{i=2}^6 \left(\frac{1}{5} + \alpha_i \right) + \alpha = \frac{5}{5} + \sum_{i=2}^6 \alpha_i + \alpha = 1 + \sum_{i=2}^6 \alpha_i + \alpha \leq 1.$$

$$\text{Then } \sum_{i=2}^6 \alpha_i \leq -\alpha < \frac{1}{5}.$$

- ⇒ We need to find a counter-objection for (P, y) .

claim: we can choose $Q = \{2, 3, 4, 5\}$ and

$$z = \langle \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5} + \alpha_4, \frac{1}{5} + \alpha_5 \rangle$$

$$z(Q) = \frac{1}{5} + \alpha_2 + \frac{1}{5} + \alpha_3 + \frac{1}{5} + \alpha_4 + \frac{1}{5} + \alpha_5 = \frac{4}{5} + \sum_{i=2}^5 \alpha_i \leq 1 \text{ since } \sum_{i=2}^5 \alpha_i \leq \sum_{i=2}^6 \alpha_i < \frac{1}{5} \text{ so } z \text{ is feasible.}$$

It is clear that $\forall i \in Q, z_i \geq x_i$ ✓ and that $\forall i \in Q \cap P, z_i \geq y_i$ ✓

- $P = \{1, 2, 3, 4, 5, 6\}$. We need to find the payoff vector $y \in \mathbb{R}^6$ so that (P, y) is an objection. $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \dots, \frac{1}{5} + \alpha_6 \rangle$.

The conditions for (P, y) to be an objection are the following:

- each agent is as well off as in x : $\alpha > -\frac{1}{5}$, $\alpha_i \geq 0$
- y is feasible for coalition P : $\sum_{i=2}^6 \left(\alpha_i + \frac{1}{5} \right) + \alpha \leq 1$.

w.l.o.g $0 \leq \alpha_2 \leq \alpha_3 \leq \alpha_4 \leq \alpha_5 \leq \alpha_6$.

$$\text{Then } \sum_{i=2}^6 \left(\frac{1}{5} + \alpha_i \right) + \alpha = \frac{5}{5} + \sum_{i=2}^6 \alpha_i + \alpha = 1 + \sum_{i=2}^6 \alpha_i + \alpha \leq 1.$$

$$\text{Then } \sum_{i=2}^6 \alpha_i \leq -\alpha < \frac{1}{5}.$$

- ⇒ We need to find a counter-objection for (P, y) .

claim: we can choose $Q = \{2, 3, 4, 5\}$ and

$$z = \langle \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5} + \alpha_4, \frac{1}{5} + \alpha_5 \rangle$$

$$z(Q) = \frac{1}{5} + \alpha_2 + \frac{1}{5} + \alpha_3 + \frac{1}{5} + \alpha_4 + \frac{1}{5} + \alpha_5 = \frac{4}{5} + \sum_{i=2}^5 \alpha_i \leq 1 \text{ since } \sum_{i=2}^5 \alpha_i \leq \sum_{i=2}^6 \alpha_i < \frac{1}{5} \text{ so } z \text{ is feasible.}$$

It is clear that $\forall i \in Q$, $z_i \geq x_i$ ✓ and that $\forall i \in Q \cap P$, $z_i \geq y_i$ ✓

Hence, (Q, z) is a counter-objection. ✓

- $P = \{1, 2, 3, 4, 5\}$.

- $P = \{1, 2, 3, 4, 5\}$. The vector
 $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5} + \alpha_4, \frac{1}{5} + \alpha_5 \rangle$ is an objection when
 $\alpha > -\frac{1}{5}, \alpha_i \geq 0, \sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha \leq 1$

- $P = \{1, 2, 3, 4, 5\}$. The vector $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5} + \alpha_4, \frac{1}{5} + \alpha_5 \rangle$ is an objection when

$$\alpha > -\frac{1}{5}, \alpha_i \geq 0, \sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha \leq 1$$

This time, we have
$$\sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha = \frac{4}{5} + \sum_{i=2}^5 \alpha_i + \alpha \leq 1$$

- $P = \{1, 2, 3, 4, 5\}$. The vector $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5} + \alpha_4, \frac{1}{5} + \alpha_5 \rangle$ is an objection when

$$\alpha > -\frac{1}{5}, \alpha_i \geq 0, \sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha \leq 1$$

This time, we have
$$\sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha = \frac{4}{5} + \sum_{i=2}^5 \alpha_i + \alpha \leq 1$$

then
$$\sum_{i=2}^5 \alpha_i \leq 1 - \frac{4}{5} - \alpha = \frac{1}{5} - \alpha \text{ and finally } \sum_{i=2}^5 \alpha_i \leq \frac{1}{5} - \alpha < \frac{2}{5}.$$

- $P = \{1, 2, 3, 4, 5\}$. The vector $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5} + \alpha_4, \frac{1}{5} + \alpha_5 \rangle$ is an objection when

$$\alpha > -\frac{1}{5}, \alpha_i \geq 0, \sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha \leq 1$$

This time, we have
$$\sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha = \frac{4}{5} + \sum_{i=2}^5 \alpha_i + \alpha \leq 1$$

then
$$\sum_{i=2}^5 \alpha_i \leq 1 - \frac{4}{5} - \alpha = \frac{1}{5} - \alpha \text{ and finally } \sum_{i=2}^5 \alpha_i \leq \frac{1}{5} - \alpha < \frac{2}{5}.$$

⇒ We need to find a counter-objection to (P, y)

- $P = \{1, 2, 3, 4, 5\}$. The vector $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5} + \alpha_4, \frac{1}{5} + \alpha_5 \rangle$ is an objection when

$$\alpha > -\frac{1}{5}, \alpha_i \geq 0, \sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha \leq 1$$

This time, we have
$$\sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha = \frac{4}{5} + \sum_{i=2}^5 \alpha_i + \alpha \leq 1$$

then
$$\sum_{i=2}^5 \alpha_i \leq 1 - \frac{4}{5} - \alpha = \frac{1}{5} - \alpha \text{ and finally } \sum_{i=2}^5 \alpha_i \leq \frac{1}{5} - \alpha < \frac{2}{5}.$$

- ⇒ We need to find a counter-objection to (P, y)

claim: we can choose $Q = \{2, 3, 6, 7\}$, $z = \langle \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5}, \frac{1}{5} \rangle$

- $P = \{1, 2, 3, 4, 5\}$. The vector $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5} + \alpha_4, \frac{1}{5} + \alpha_5 \rangle$ is an objection when $\alpha > -\frac{1}{5}$, $\alpha_i \geq 0$, $\sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha \leq 1$

This time, we have $\sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha = \frac{4}{5} + \sum_{i=2}^5 \alpha_i + \alpha \leq 1$

then $\sum_{i=2}^5 \alpha_i \leq 1 - \frac{4}{5} - \alpha = \frac{1}{5} - \alpha$ and finally $\sum_{i=2}^5 \alpha_i \leq \frac{1}{5} - \alpha < \frac{2}{5}$.

⇒ We need to find a counter-objection to (P, y)

claim: we can choose $Q = \{2, 3, 6, 7\}$, $z = \langle \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5}, \frac{1}{5} \rangle$

It is clear that $\forall i \in Q$, $z_i \geq x_i$ ✓ and $\forall i \in P \cap Q$ $z_i \geq y_i$ (for agent 2 and 3).

- $P = \{1, 2, 3, 4, 5\}$. The vector $y = \langle \alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5} + \alpha_4, \frac{1}{5} + \alpha_5 \rangle$ is an objection when $\alpha > -\frac{1}{5}$, $\alpha_i \geq 0$, $\sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha \leq 1$

This time, we have $\sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha = \frac{4}{5} + \sum_{i=2}^5 \alpha_i + \alpha \leq 1$

then $\sum_{i=2}^5 \alpha_i \leq 1 - \frac{4}{5} - \alpha = \frac{1}{5} - \alpha$ and finally $\sum_{i=2}^5 \alpha_i \leq \frac{1}{5} - \alpha < \frac{2}{5}$.

⇒ We need to find a counter-objection to (P, y)

claim: we can choose $Q = \{2, 3, 6, 7\}$, $z = \langle \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5}, \frac{1}{5} \rangle$

It is clear that $\forall i \in Q$, $z_i \geq x_i$ ✓ and $\forall i \in P \cap Q$ $z_i \geq y_i$ (for agent 2 and 3).

$z(Q) = \frac{1}{5} + \alpha_2 + \frac{1}{5} + \alpha_3 + \frac{1}{5} + \frac{1}{5} = \frac{4}{5} + \alpha_2 + \alpha_3$. We have $\alpha_2 + \alpha_3 < \frac{1}{5}$, otherwise, we would have $\alpha_2 + \alpha_3 \geq \frac{1}{5}$ and since

the α_i are ordered, we would then have $\sum_{i=2}^5 \alpha_i \geq \frac{2}{5}$, which is

not possible. Hence $z(Q) \leq 1$ which proves z is feasible ✓

Using similar arguments, we find a counter-objection for each other objections (you might want to fill in the details at home).

- $P = \{1, 2, 3, 4\}$, $y = \langle \alpha, \frac{1}{5} + \alpha_1, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3 \rangle$, $\alpha > -\frac{1}{5}$, $\alpha_i \geq 0$,
 $\sum_{i=2}^4 \alpha_i + \alpha \leq \frac{2}{5} \Rightarrow \sum_{i=2}^4 \alpha_i \leq \frac{2}{5} - \alpha < \frac{3}{5}$.
- ⇒ $Q = \{2, 5, 6, 7\}$, $z = \langle \frac{1}{5} + \alpha_2, \frac{1}{5}, \frac{1}{5}, \frac{1}{5} \rangle$ since $\alpha_2 \leq \frac{1}{5}$
- $|P| \leq 3$ $P = \{1, 2, 3\}$, $v(P) = 0$, $y = \langle \alpha, \alpha_1, \alpha_2 \rangle$, $\alpha > -\frac{1}{5}$,
 $\alpha_i \geq 0$, $\alpha_1 + \alpha_2 \leq -\alpha < \frac{1}{5}$
- ⇒ $Q = \{4, 5, 6, 7\}$, $z = \langle \frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5} \rangle$ will be a counter argument
 (1 cannot provide more than $\frac{1}{5}$ to any other agent).
- For each possible objection of 1, we found a counter-objection. Using similar arguments, we can find a counter-objection to any objection of player 7 against player 1.
- ⇒ $x \in preBS(N, v, S)$. ✓

Using similar arguments, we find a counter-objection for each other objections (you might want to fill in the details at home).

- $P = \{1, 2, 3, 4\}$, $y = \langle \alpha, \frac{1}{5} + \alpha_1, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3 \rangle$, $\alpha > -\frac{1}{5}$, $\alpha_i \geq 0$,
 $\sum_{i=2}^4 \alpha_i + \alpha \leq \frac{2}{5} \Rightarrow \sum_{i=2}^4 \alpha_i \leq \frac{2}{5} - \alpha < \frac{3}{5}$.
- ⇒ $Q = \{2, 5, 6, 7\}$, $z = \langle \frac{1}{5} + \alpha_2, \frac{1}{5}, \frac{1}{5}, \frac{1}{5} \rangle$ since $\alpha_2 \leq \frac{1}{5}$
- $|P| \leq 3$ $P = \{1, 2, 3\}$, $v(P) = 0$, $y = \langle \alpha, \alpha_1, \alpha_2 \rangle$, $\alpha > -\frac{1}{5}$,
 $\alpha_i \geq 0$, $\alpha_1 + \alpha_2 \leq -\alpha < \frac{1}{5}$
- ⇒ $Q = \{4, 5, 6, 7\}$, $z = \langle \frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5} \rangle$ will be a counter argument
 (1 cannot provide more than $\frac{1}{5}$ to any other agent).
- For each possible objection of 1, we found a counter-objection. Using similar arguments, we can find a counter-objection to any objection of player 7 against player 1.
- ⇒ $x \in preBS(N, v, S)$. ✓

Using similar arguments, we find a counter-objection for each other objections (you might want to fill in the details at home).

- $P = \{1, 2, 3, 4\}$, $y = \langle \alpha, \frac{1}{5} + \alpha_1, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3 \rangle$, $\alpha > -\frac{1}{5}$, $\alpha_i \geq 0$,
 $\sum_{i=2}^4 \alpha_i + \alpha \leq \frac{2}{5} \Rightarrow \sum_{i=2}^4 \alpha_i \leq \frac{2}{5} - \alpha < \frac{3}{5}$.
- ⇒ $Q = \{2, 5, 6, 7\}$, $z = \langle \frac{1}{5} + \alpha_2, \frac{1}{5}, \frac{1}{5}, \frac{1}{5} \rangle$ since $\alpha_2 \leq \frac{1}{5}$
- $|P| \leq 3$ $P = \{1, 2, 3\}$, $v(P) = 0$, $y = \langle \alpha, \alpha_1, \alpha_2 \rangle$, $\alpha > -\frac{1}{5}$,
 $\alpha_i \geq 0$, $\alpha_1 + \alpha_2 \leq -\alpha < \frac{1}{5}$
- ⇒ $Q = \{4, 5, 6, 7\}$, $z = \langle \frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5} \rangle$ will be a counter argument
 (1 cannot provide more than $\frac{1}{5}$ to any other agent).
- For each possible objection of 1, we found a counter-objection. Using similar arguments, we can find a counter-objection to any objection of player 7 against player 1.
- ⇒ $x \in preBS(N, v, S)$. ✓

Bargaining set

In the example, agent 1 gets $-\frac{1}{5}$ when $v(\mathcal{C}) \geq 0$ for all coalition $\mathcal{C} \subseteq N$! This shows that the pre-bargaining set may **not** be individually rational.

Let $I(N, v, \mathcal{S}) = \{x \in X_{(N, v, \mathcal{S})} \mid x_i \geq v(\{i\}) \forall i \in N\}$ be the **set of individually rational payoff vector** in $X_{(N, v, \mathcal{S})}$.

Lemma

If a game is weakly superadditive, $I(N, v, \mathcal{S}) \neq \emptyset$.

Definition (Bargaining set)

Let (N, v, \mathcal{S}) a game in coalition structure.

The **bargaining set** (**BS**) is defined by

$$BS(N, v, \mathcal{S}) = I(N, v, \mathcal{S}) \cap preBS(N, v, \mathcal{S}).$$

Lemma

We have $Core(N, v, \mathcal{S}) \subseteq BS(N, v, \mathcal{S})$.

Theorem

Let (N, v, \mathcal{S}) a game with coalition structure. Assume that $I(N, v, \mathcal{S}) \neq \emptyset$. Then the bargaining set $BS(N, v, \mathcal{S}) \neq \emptyset$.

Proof

It is possible to give a direct proof of this theorem (a bit long, (Section 4.2 in **Introduction to the Theory of Cooperative Games**)).

We will show this result in a different way in the lecture about the nucleolus next week. □

B. Peleg and P. Sudhölter **Introduction to the Theory of Cooperative Games**, Springer, 2007.

Definition (weighted voting games)

A game $(N, w_{i \in N}, q, v)$ is a **weighted voting game** when v satisfies unanimity, monotonicity and the valuation function is defined as

$$v(S) = \begin{cases} 1 & \text{when } \sum_{i \in S} w_i \geq q \\ 0 & \text{otherwise} \end{cases}$$

We note such a game by $(q : w_1, \dots, w_n)$

Let (N, v) be the game associated with the 6-player weighted majority game $(3:1,1,1,1,1,0)$.

Agent 6 is a null/dummy player since its weight is 0.

Nevertheless $\langle \frac{1}{7}, \dots, \frac{1}{7}, \frac{2}{7} \rangle \in BS(N, v)$.

Proof

This will be part of homework 2



Agent 6 is a dummy, however, it receives a payoff of $\frac{2}{7}$, which is larger than agents who are not dummy!

Remember: $mc_i^{max} = \max_{\mathcal{C} \subseteq N \setminus \{i\}} v(\mathcal{C} \cup \{i\}) - v(\mathcal{C})$

x is **reasonable from above** if $\forall i \in N \ x^i < mc_i^{max}$

$\Leftrightarrow mc_i^{max}$ is the strongest **threat** that an agent can use against a coalition.

The bargaining set is not **Reasonable from above**: the dummy agent gets more than $\max_{\mathcal{C} \subseteq N \setminus \{6\}} (v(\mathcal{C} \cup \{6\}) - v(\mathcal{C})) = 0$. **✗**

Lemma

The core is reasonable for above and from below.

Proof

Since the core satisfies IR, it must be reasonable from below. Let (N, v) be a game, $x \in \text{Core}(N, v)$ and $i \in N$. Then $x(N) = v(N)$ and $x(N \setminus \{i\}) \geq v(N \setminus \{i\})$. Then $x_i = v(N) - x(N \setminus \{i\}) \leq v(N) - v(N \setminus \{i\}) \leq mc_i^{max}$.

□

Summary

- We introduced the bargaining set, and looked at some examples.
 - pros:** it is guaranteed to be non-empty, when the core is non-empty, it is contained in the bargaining set.
 - cons:** it may not be reasonable from above.

Coming next

- We will consider the Nucleolus. It can also be defined in terms of objections and counter objections, but the nature of the objection is different from the bargaining set.