



Commitment under uncertainty: Two-stage stochastic matching problems

Irit Katriel, Claire Kenyon-Mathieu, Eli Upfal *

Computer Science Department, Brown University, Box 1910, Providence, RI 02906, United States

ARTICLE INFO

Keywords:

Stochastic combinatorial optimization
Matching algorithm

ABSTRACT

We define and study two versions of the bipartite matching problem in the framework of two-stage stochastic optimization with recourse. In one version, the uncertainty is in the second stage costs of the edges, and in the other version, the uncertainty is in the set of vertices that needs to be matched. We prove lower bounds, and analyze efficient strategies for both cases. These problems model real-life stochastic integral planning problems, such as commodity trading, reservation systems and scheduling under uncertainty.

© 2008 Elsevier B.V. All rights reserved.

1. Introduction

Two-stage stochastic optimization with recourse is a popular model for hedging against uncertainty. Typically, part of the input to the problem is only known probabilistically in the first stage, when decisions have a low cost. In the second stage, the actual input is known, but the costs of the decisions are higher. We then face a delicate tradeoff between speculating at a low cost vs. waiting for the uncertainty to be resolved.

This model has been studied extensively for problems that can be modeled by linear programming (sometimes using techniques such as Sample Average Approximation (SAA) when the linear program (LP) is too large). Recently there has been a growing interest in 2-stage stochastic combinatorial optimization problems [1,2,6,12,19–22,24]. Since an LP relaxation does not guarantee an integer solution in general, one can either try to find an efficient rounding technique [11], or develop a purely combinatorial approach [5,8]. In order to develop successful algorithmic paradigms in this setting, there is an ongoing research program focusing on classical combinatorial optimization problems [23]: set cover, minimum spanning tree, Steiner tree, maximum weight matching, facility location, bin packing, multicommodity flow, minimum multicut, knapsack, and others. In this paper, we aim to enrich this research program by adding a basic combinatorial optimization problem to the list: the minimum cost maximum bipartite matching problem. The task is to buy edges of a bipartite graph, which together contain a maximum-cardinality matching in the graph. We examine two variants of this problem. In the first, the uncertainty is in the second stage edge-costs, that is, the cost of an edge can either grow or shrink in the second stage. In the second variant, all edges become more expensive in the second stage, but the set of nodes that need to be matched is not known.

Here are some features of minimum cost maximum bipartite matching that make this problem particularly interesting. First, it is not subadditive: the union of two feasible solutions is not necessarily a solution for the union of the two instances. In contrast, most previous work focused on subadditive structures, with the notable exception of Gupta and Pál's work on stochastic Steiner Tree [9]. Second, the solutions to two partial instances may interfere with one another in a way that seems to preclude the possibility of applying cost-sharing techniques associated with the scenario-sampling based algorithms [9, 10]. This intuitively makes the problem resistant to routine attempts, and indeed, we confirm this intuition by proving a

* Corresponding author.

E-mail addresses: irit@cs.brown.edu (I. Katriel), claire@cs.brown.edu (C. Kenyon-Mathieu), eli@cs.brown.edu (E. Upfal).




Fig. 1. An example in which buying edges speculatively can help.

lower bound which is stronger than what is known¹ for the sub-additive problems: in [Theorem 5](#), we prove a hardness of approximation result in the setting where the second-stage scenarios are generated by choosing vertices independently. It is therefore natural that our algorithms yield upper bounds which are either rather weak ([Theorem 2](#), Part 1) or quite specialized ([Theorem 7](#)). To address this issue, we relax the constraint that the output be a maximum matching, and consider bicriteria results, where there is a tradeoff between the cost of the edges bought and the size of the resulting matching ([Theorem 2](#), Part 2, and [Theorem 8](#)). This approach may be a way to circumvent hardness for other stochastic optimization problems, as well.

Although the primary focus of this work is stochastic optimization, another popular objective for the prudent investor is to minimize, not just the expected future cost, but the maximum future cost, over all possible future scenarios: that is the goal of robust optimization. We also prove a bicriteria result for robust optimization ([Theorem 3](#).) Guarding oneself against the worst case is more delicate than just working with expectations. The solution requires a different idea: preventing undesirable high-variance events by explicitly deciding, against the advice of the LP solution, to not buy expensive edges (To analyze this, the proof of [Theorem 3](#) involves some careful rounding.) This general idea might be applicable to other problems as well.

We note that within two-stage stochastic optimization with recourse, matching has been studied before [[14](#)]. However, the problem studied here is very different: there, the goal was to construct a maximum weight matching instead of the competing objective of large size and small cost; moreover the set of edges bought by the algorithm had to form exactly a matching instead of just contain a matching. In [Fig. 1](#), we give an example illustrating the difference between requiring equality with a matching or containment of a matching. If we are allowed to buy edges without committing to use them later, then the correct strategy is to buy edges $\{a, c\}$ and $\{a, d\}$ in the first stage, and then either $\{b, c\}$ or $\{b, d\}$ in the second stage, depending on which scenario materializes. The total cost is 0. If buying an edge commits to use it in the matching, then the correct strategy is to buy edge $\{b, d\}$ in the first stage and $\{a, c\}$ in the second stage, for a total cost of 200: the two problems have completely different structures.

Our main goal in this paper is to further fundamental understanding of the theory of stochastic optimization; however, we note that a conceivable application of this problem is commodity transactions, which can be viewed as a matching between supply and demand. When the commodity is indivisible, the set of possible transactions can be modeled as a weighted bipartite graph matching problem, where the weight of an edge represents the cost or profit of that transaction (including transportation cost when applicable). A trader tries to maximize profits or minimize total costs, depending on their position in the transaction. A further tool that a commodity trader may employ to improve their income is timing the transaction. We model timing as a two-stage stochastic optimization problem with recourse: The trader can limit their risk by buying an option for a transaction at current information, or they can assume the risk and defer decisions to the second stage. Two common uncertainties in commodity transactions, price uncertainty and supply and demand uncertainty, correspond to the two stochastic two-stage matching problems mentioned above: finding minimum weight maximum matching with uncertain edge costs, and finding maximum matching with uncertain matching vertices. Similar decision scenarios involving matchings also show up in a variety of other applications, such as scheduling and reservation systems.

Our results are summarized in the following table. We first prove ([Theorem 1](#)) that, with explicit scenarios, the uncertain matching vertices case is, in fact, a special case of the uncertain edge costs case. Then, it suffices to prove upper bounds for the more general variant and lower bounds for the restricted one. For the problem of minimizing the expected cost of the solution, we show an approximability lower bound of $\Omega(\log n)$. We then describe an algorithm that finds a maximum matching in the graph at a cost which is an n^2 -approximation for the optimum. We then show that by relaxing the demand that the algorithm constructs a maximum matching, we can “beat” the lower bound: at a cost of at most $1/\beta$ times the optimum, we can match at least $n(1 - \beta)$ vertices. Furthermore, we show that a similar bicriteria result holds also for the robust version of the problem, i.e., when we wish to minimize the worst-case cost.

With independent choices in the second-stage scenarios, our main contribution is the lower bound. The reduction of [Theorem 1](#) does not apply, but we prove, for both types of uncertainty, that it is NP-hard to approximate the problem within better than a certain constant factor. We also prove an upper bound for a special case of the uncertain matching vertices variant.

¹ To the best of our knowledge, all previous hardness results hold only when the second stage scenarios are given explicitly, i.e., when only certain combinations of parameter settings are possible.

Input:	Explicit Scenarios		Independent Choices
Criteria:	Expected Cost	Worst-Case Cost	Expected Cost
Uncertain edge costs	<ul style="list-style-type: none"> • n^2-approximation of the cost to get a maximum matching [Theorem 2, part 1] • $1/\beta$-approximation of the cost to match at least $n(1 - \beta)$ vertices [Theorem 2, part 2] <ul style="list-style-type: none"> • Same hardness results as below [Theorem 1] 	factor $1/\beta$ approximation of the cost to match at least $n(1 - \beta)$ vertices [Theorem 3]	NP-hard to approximate within a certain constant [Theorem 6]
Uncertain matching vertices	<ul style="list-style-type: none"> • $\Omega(\log n)$ approximability lower bound [Theorem 4, Part 1] • NP-hard already for two scenarios [Theorem 4, Part 2] • Same upper bounds as above [Theorem 1] 	As above [Theorem 1]	<ul style="list-style-type: none"> • As above [Theorem 5] • approximation for a special case [Theorem 7]

2. Explicit scenarios

In this section, we assume that we have an explicit list of possible scenarios for the second stage.

Uncertain edge costs. Given a bipartite graph $G = (A, B, E)$, we can buy edge e in the first stage at cost $C_e \geq 0$, or we can buy it in the second stage at cost $C_e^s \geq 0$ determined by the scenario s . The input has an explicit list of scenarios, and known edge costs (c_e^s) in scenario s . For uncertain edge costs, without loss of generality, we can assume that $|A| = |B| = n$ and that G has a perfect matching. Indeed, there is an easy reduction from the case where the maximum matching has size k : just create a new graph by adding a set A' of $n - k$ vertices on the left side, a set B' of $n - k$ vertices on the right side, and edges between all vertex pairs in $A' \times B$ and in $A \times B'$, with cost 0.

In the *stochastic optimization* setting, the algorithm also has a known second stage distribution: scenario s occurs with probability $\Pr(s)$. The goal is, in time polynomial in both the size of the graph and the number of scenarios, to minimize the *expected cost*; if E_1 denotes the set of edges bought in the first stage and E_2^s the set of edges bought in the second stage under scenario s , then:

$$\text{OPT}_1 = \min_{E_1, E_2^s} \left\{ \sum_{s \in S} \Pr(s) \left(\sum_{e \in E_1} C_e + \sum_{e \in E_2^s} C_e^s \right) : \forall s, E_1 \cup E_2^s \text{ contains a perfect matching} \right\}.$$

Stochastic optimization with uncertain edge costs has been studied for many problems, see for example [10,17].

In the *robust optimization* setting, the goal is to minimize the maximum cost (instead of the expected cost):

$$\text{OPT}_2 = \min_{E_1, E_2^s} \left\{ \max_{s \in S} \left(\sum_{e \in E_1} C_e + \sum_{e \in E_2^s} C_e^s \right) : \forall s, E_1 \cup E_2^s \text{ contains a perfect matching} \right\}.$$

Robust optimization with uncertain edge costs has also been studied for many problems, see for example [4].


Uncertain activated vertices. In this variant of the problem, there is a known distribution over scenarios s , each being defined by a set $B_s \subset B$ of *active vertices* that are allowed to be matched in that scenario. Each edge costs c_e today (before B_s is known) and τc_e tomorrow, where $\tau > 1$ is the *inflation parameter*. As in Expression (1), the goal is to minimize the expected cost, i.e.,

$$\text{OPT}_3 = \left\{ C(E_1) + \tau \sum_{s \in S} \Pr(s) C(E_2^s) : \forall s, E_1 \cup E_2^s \text{ contains a maximum matching of } (A, B_s, E \cap (A \times B_s)) \right\}.$$

Stochastic optimization with uncertain activated vertices has also been previously studied for many problems, see for example [9]. There is a similar expression for robust optimization with uncertain activated vertices.

Theorem 1 (Reduction). *The two-stage stochastic matching problem with uncertain activated vertices and explicit second-stage scenarios (OPT_3) reduces to the case of uncertain edge costs and explicit second-stage scenarios (OPT_1).*

Proof. We give an approximation preserving reduction. Given an instance with stochastic matching vertices, we transform it to an instance of the problem with stochastic edge-costs, as follows. Assume that our input graph is $G = (A, B, E)$ where $A = \{a_1, \dots, a_{|A|}\}$ and $B = \{b_1, \dots, b_{|B|}\}$. We first add a set $A' = \{a'_1, \dots, a'_{|B|}\}$ of $|B|$ new vertices to A , and connect each a'_i with b_i by an edge. In other words, we generate the graph $G' = (A \cup A', B, E \cup \{(a'_i, b_i) : 1 \leq i \leq |B|\})$.



For the edges between A and B , edge costs are the same as in the original instance, in the first stage as well as the second stage. The costs on the edges between A' and B create the effect of selecting the activated vertices: for each (a'_i, b_i) , the first-stage cost is n^2W , and the second-stage cost is n^2W if b_i is active and 0 otherwise. Here, W is the maximum cost of an edge, nW is an upper bound on the cost of the optimal solution, and n^2W is large enough that any solution containing this edge cannot be an optimal, or even an n -approximate solution. Hence, a second-stage cost of 0 for (a'_i, b_i) allows b_i to be matched with a'_i for free, while a cost of nW forces b_i to be matched with a vertex from A . This concludes the reduction. \square

From [Theorem 1](#), it follows that our algorithms for uncertain edges costs ([Theorems 2 and 3](#)) imply corresponding algorithms for uncertain activated vertices, and that our lower bounds for uncertain activated vertices ([Theorem 4](#)) imply corresponding lower bounds for uncertain edge costs.

Theorem 2 (Stochastic Optimization Upper Bound). (1) There is a polynomial-time deterministic algorithm for stochastic matching (OPT_1) that constructs a perfect matching whose expected cost is at most $2n^2 \cdot \text{OPT}_1$.
 (2) Given $\beta \in (0, 1)$, there is a polynomial-time randomized algorithm for stochastic matching (OPT_1) that returns a matching whose cardinality, with probability $1 - e^{-n}$ (over the random choices of the algorithm), is at least $(1 - \beta)n$, and whose overall expected cost is $O(\text{OPT}_1/\beta)$.

In particular, for any $\epsilon > 0$ we get a matching of size $(1 - \epsilon)n$ and cost $O(\text{OPT}/\epsilon)$ in expectation. Note that by [Theorem 4](#), we have to relax the constraint on the size anyway if we wish to obtain a better-than-log n approximation on the cost, so Part 2 of the Theorem is, in a sense, our best option.

The proof follows the general paradigm applied to stochastic optimization in recent papers such as [11]: formulate the problem as an integer linear program; solve the linear relaxation and use it to guide the algorithm; and use LP duality (König's theorem, for our problem) for the analysis.

To define the integer program, let X_e indicate whether edge e is bought in the first stage, and for each scenario s , let Z_e^s (resp. Y_e^s) indicate whether edge e is bought in the first stage (resp. in the second stage) and ends up in the perfect matching when scenario s materializes. We obtain:

$$\min \sum_{s \in S} \Pr(s) \left(\sum_e C_e X_e + \sum_e C_e^s Y_e^s \right) \text{ s.t. } \begin{cases} \sum_{e: v \in e} (Z_e^s + Y_e^s) = 1 & \forall v \in A \cup B, s \in S \\ Z_e^s \leq X_e & \forall e \in E, s \in S \\ X_e, Y_e^s, Z_e^s \in \{0, 1\} & \forall e \in E, s \in S. \end{cases}$$

The algorithm solves the standard LP relaxation:

$$\min \sum_{s \in S} \Pr(s) \left(\sum_e C_e X_e + \sum_e C_e^s Y_e^s \right) \text{ s.t. } \begin{cases} \sum_{e: v \in e} (Z_e^s + Y_e^s) = 1 & \forall v \in A \cup B, s \in S \\ Z_e^s \leq X_e & \forall e \in E, s \in S \\ \leq X_e, Y_e^s, Z_e^s \leq 1 & \forall e \in E, s \in S. \end{cases} \quad (1)$$

Now the proof of the two parts of the theorem diverges:

Proof of part 1. Here is the algorithm.

Solve the Linear Program (1). Let (X_e, Z_e^s, Y_e^s) denote the optimal solution.

First stage: buy every edge e such that $X_e \geq 1/(2n^2)$.

Second stage: under scenario s , buy every edge e such that $Y_e^s \geq 1/(2n^2)$.

Output: a maximum matching of the set of edges bought.

For the analysis, we see that the expected cost is

$$\begin{aligned} \sum_{s \in S} \Pr(s) \left[\sum_{e: X_e \geq 1/(2n^2)} C_e + \sum_{e: Y_e^s \geq 1/(2n^2)} C_e^s \right] &\leq 2n^2 \sum_{s \in S} \Pr(s) \left[\sum_e C_e X_e + \sum_e C_e^s Y_e^s \right] \\ &= 2n^2 \text{OPT}. \end{aligned}$$

It only remains to prove that for every scenario s , the output is a perfect matching.

Lemma 1 (Hall's Theorem). *A bipartite graph with vertex set (A, B) contains a perfect matching if and only if every subset U of A has at least $|U|$ neighbors in B .*

To apply Hall's theorem, fix a subset U of A and let $N(U) = \{w \in B \mid \exists v \in U, P_{\{v,w\}} \geq 1/n^2\}$, where $P_e = Z_e^s + Y_e^s$. Note that if $P_{\{v,w\}} \geq 1/n^2$, then at least one of $Z_{\{v,w\}}^s$ or $Y_{\{v,w\}}^s$ must be greater than or equal to $1/(2n^2)$, and so, the algorithm must have bought edge $\{v, w\}$ under scenario s : thus $N(U)$ is contained in the set of neighbors of U in the graph. Now, since (P_e) is a fractional perfect matching, we have $\sum_{e \in U \times B} P_e = |U|$ and $\sum_{e \in U \times N(U)} P_e \leq |N(U)|$. By definition of $N(U)$, we have $\sum_{e \in U \times (B \setminus N(U))} P_e \leq |U|(n - |N(U)|)(1/n^2)$, and so

$$|U| \leq |N(U)| + \frac{|U|(n - |N(U)|)}{n^2} < |N(U)| + 1.$$

Since $|N(U)|$ is an integer, it must therefore be greater than or equal to $|U|$.

Hence by Hall's theorem there is a matching of size n .

Proof of part 2. Here is the algorithm.

Solve the Linear Program (1). Let (X_e, Z_e^s, Y_e^s) denote the optimal solution.
 Let $\alpha = 8 \ln(2)/\beta$.
First stage: buy every edge e independently with probability $1 - e^{-X_e \alpha}$.
Second stage: under scenario s , buy every edge e independently with probability $1 - e^{-Y_e^s \alpha}$.
Output: a maximum matching of the set of edges bought.

For the analysis, we see that the expected cost of the output is

$$\sum_e \left(C_e (1 - e^{-X_e \alpha}) + \sum_s \Pr(s) C_e^s (1 - e^{-Y_e^s \alpha}) \right).$$

Using the upper bound $1 - e^{-Z} \leq Z$, we deduce that this quantity is at most α times the objective function of our linear program, i.e. at most α times OPT.

Let $\beta' = \beta/2$ where we recall that the goal of the theorem is to have a matching of expected size $n - \beta n$. We will prove that with high probability, the output has cardinality at least $n(1 - \beta')$. Indeed, assume that the output has cardinality less than $n(1 - \beta')$.

Lemma 2 (König's Theorem [15]). *In a bipartite graph, the cardinality of a maximum matching equals the cardinality of a minimum vertex cover.*

We apply König's theorem: there exists a set of vertices, of cardinality less than $n(1 - \beta')$, which covers all of $E_1 \cup E_2^s$.

Fix a subset V of $A \cup B$ of cardinality less than $n(1 - \beta')$. For any edge e that remains uncovered by V , the probability that the algorithm does not buy e is $e^{-X_e \alpha} e^{-Y_e^s \alpha} \leq e^{-P_e \alpha}$, where $P_e = Z_e^s + Y_e^s$. Thus the probability that V is a vertex cover is bounded by $\prod_{e: e \cap V = \emptyset} e^{-P_e \alpha} = e^{-\sum_{e: e \cap V = \emptyset} P_e \alpha}$. By the LP constraints and the fact that G is bipartite, $(P_e)_e$ is a convex combination of perfect matchings, each of which has at most $|V|$ edges adjacent to V , hence has at least $\beta'n$ edges not covered by V . Thus the sum of P_e , over edges e left uncovered by V , is at least $\beta'n$. So, the probability that V is a vertex cover is bounded by $e^{-\alpha \beta' n}$.

By the union bound, the probability that there exists such a vertex cover is at most $2^{2n} e^{-\alpha \beta' n} = e^{-(\alpha \beta' - 2 \ln 2)n}$. Thus the output matching has size $n(1 - \beta')$ with probability at least $(1 - e^{-(\alpha \beta' - 2 \ln 2)n})$, and the expected size is at least $(1 - e^{-(\alpha \beta' - 2 \ln 2)n})(n(1 - \beta')) \geq n(1 - \beta')$. \square

Theorem 3 (Robust Optimization). *Given $\beta \in (0, 1)$, there is a polynomial-time randomized algorithm for robust matching (OPT₂) with t scenarios that returns a matching s.t. with probability at least $1 - 2/n$ (over the random choices of the algorithm), the following holds: In every scenario, the algorithm incurs cost $O(\text{OPT}_2(1 + \ln(t)/\ln(n))/\beta)$ and outputs a matching of cardinality at least $(1 - \beta)n$.*

Proof. We detail this proof, which is the most interesting one in this section. The integer programming formulation is similar to the one used to prove [Theorem 2](#). More specifically, let X_e indicate whether edge e is bought in the first stage, and for each scenario s , let Z_e^s (resp. Y_e^s) indicate whether edge e is bought in the first stage (resp. in the second stage) and ends up in the perfect matching when scenario s materializes. We obtain:

$$\min W \text{ s.t. } \begin{cases} \sum_{e:v \in e} (Z_e^s + Y_e^s) = 1 & \forall v \in A \cup B \text{ and } \forall s \in S \\ Z_e^s \leq X_e & \forall e \in E \text{ and } s \in S \\ \sum_e [C_e X_e + C_e^s Y_e^s] \leq W & \forall s \in S \\ X_e, Y_e^s, Z_e^s \in \{0, 1\} & \forall e \in E \text{ and } s \in S. \end{cases}$$

The algorithm solves the standard LP relaxation:

$$\min W \text{ s.t. } \begin{cases} \sum_{e:v \in e} (Z_e^s + Y_e^s) = 1 & \forall v \in A \cup B \text{ and } \forall s \in S \\ Z_e^s \leq X_e & \forall e \in E \text{ and } s \in S \\ \sum_e [C_e X_e + C_e^s Y_e^s] \leq W & \forall s \in S \\ 0 \leq X_e, Y_e^s, Z_e^s \leq 1 & \forall e \in E \text{ and } s \in S. \end{cases} \quad (2)$$

Solve the Linear Program (2). Let $w, (x_e), (y_e^s), (z_e^s)$ denote the optimal solution.

Let $\alpha = 8 \ln(2)/\beta$ and $T = 3 \ln n$.

First stage:

Relabel the edges so that $c_1 \geq c_2 \geq \dots$.

Let t_1 be maximum such that $x_1 + x_2 + \dots + x_{t_1} \leq T$.

For every $j > t_1$, buy edge j with probability $1 - e^{-x_j \alpha}$.

(Do not buy any edge $j \leq t_1$.)

Second stage: under scenario s ,

Relabel the remaining edges so that $c_1^s \geq c_2^s \geq \dots$.

Let t_2 be maximum such that $y_1^s + y_2^s + \dots + y_{t_2}^s \leq T$.

For every $j > t_2$, buy edge j with probability $1 - e^{-y_j^s \alpha}$.

(Do not buy any edge $j \leq t_2$.)

Output: a maximum matching of the set of edges bought.

We note that this construction and the rounding used in the analysis are almost identical to the construction used in strip-packing [13]. The analysis of the cost of the edges bought is the difficult part. We first do a slight change of notation. The cost can be expressed as the sum of at most $2m$ random variables (at most m in each stage). Let $a_1 \geq a_2 \geq \dots$ be the multiset $\{c_e\} \cup \{c_e^s\}$, along with the corresponding probabilities p_i ($p_i = 1 - e^{-x_e \alpha}$ if $a_i = c_e$ is a first-stage cost, and $p_i = 1 - e^{-y_e^s \alpha}$ if $a_i = c_e^s$ is a second-stage cost.) Let X_i be the binary variable with expectation p_i . Clearly, the cost incurred by the algorithm can be bounded above by $X = \sum_{i \geq t^*} a_i X_i$, where t^* is maximum such that $p_1 + \dots + p_{t^*} \leq T$.

To prove a high-probability bound on X , we will partition $[1, 2m]$ into intervals to define groups. The first group is just $[1, t^*]$, and the subsequent groups are defined in greedy fashion, with group $[j, \ell]$ defined by choosing ℓ maximum so that $\sum_{i \in [j, \ell]} p_i \leq T$. Let G_1, G_2, \dots, G_r be the resulting groups. We have:

$$X \leq \sum_{\ell \geq 2} \sum_{i \in G_\ell} a_i X_i \leq \sum_{\ell \geq 2} \sum_{i \in G_\ell} (\max_{G_\ell} a_i) X_i \leq \sum_{\ell \geq 2} \sum_{i \in G_\ell} (\min_{G_{\ell-1}} a_i) X_i \leq \sum_{\ell \geq 1} (\min_{G_\ell} a_i) \sum_{i \in G_{\ell+1}} X_i.$$

On the other hand, (using the inequality $1 - e^{-Z} \leq Z$), the optimal value OPT^* of the LP relaxation satisfies:

$$\alpha \text{OPT}^* \geq \sum_i a_i p_i \geq \sum_{\ell \geq 1} \sum_{i \in G_\ell} (\min_{G_\ell} a_i) p_i \geq \sum_{\ell \geq 1} (\min_{G_\ell} a_i) (T - 1).$$

It remains, for each group G_ℓ , to apply a standard Chernoff bound to bound the sum of the X_i 's in G_ℓ , and use union bounds to put these results together and yield the statement of the theorem, as follows.

Lemma 3 (Chernoff). Let $X = \sum_{1 \leq i \leq N} X_i$ be a sum of independent binary random variables, with with expected value $E(X_i)$ for all i , and let σ^2 denote the variance of X . Then $\Pr(X - E(X) \geq k\sigma) \leq e^{-k^2/4}$ for any $k \in [0, 2\sigma]$.

We apply the Chernoff bound separately for each group with $k = 2\sqrt{3 \ln n + \ln t}$, where t is the number of scenarios. For a given group G_ℓ , we have $\sigma^2 = \sum_{i \in G_\ell} p_i(1 - p_i) \leq \sum p_i \leq T$; thus, the event that $\sum_{i \in G_\ell} X_i$ is less than $2\sqrt{T(3 \ln n + \ln t)}$ has probability at least $1 - 1/(n^3 t)$. The total number of groups is at most $2m \leq n^2$, so with probability at least $1 - 1/(nt)$, the event holds for every group; then we have

$$X \leq \sum_{\ell \geq 1} (\min_{G_\ell} a_i) 2\sqrt{T(3 \ln n + \ln t)} \leq 2\alpha \text{OPT}^* \left(1 + \frac{\ln t}{3 \ln n}\right) \frac{T}{T - 1} \leq 3\alpha \text{OPT}^* \left(1 + \frac{\ln t}{3 \ln n}\right).$$




Fig. 2. Reduction from minimum set cover.

In other words, with probability at least $1 - 1/(nt)$ the cost is bounded by $3\alpha \text{OPT}^* (1 + \frac{\ln t}{3 \ln n})$. Now, recall that t is the total number of scenarios. By the union bound, it holds that with probability at least $1 - 1/n$, we have that, for every scenario, the cost of the algorithm is bounded by $3\alpha \text{OPT}^* (1 + \frac{\ln t}{3 \ln n})$. But recall that, by assumption in this section, the number of scenarios is polynomial in n : then $\ln(t) = O(\ln n)$, and the part of the theorem about the cost follows.

For analysis of the size, it is easy to extend the proof of Theorem 2 so as to show that the output matching has size at least $n(1 - \beta/2)$ with probability at least $1 - e^{-(\alpha\beta/2 - \ln 4)n + 2\alpha T}$. Since $\alpha = o(n/\ln n)$, we have $2\alpha T = o(n \ln 4)$. From our lower bound on $\alpha\beta$, it follows easily that this probability is at least $1 - 1/(nt)$, where t is the number of scenarios. Using the union bound over all scenarios proves that with probability at least $1 - 1/n$, we have that for every scenario the size of the matching satisfies the stated bound of the theorem. \square

We note that the proof of Theorem 3 can also be extended to the setting of Theorem 2 to prove a high probability result: for scenario s , with probability at least $1 - 2/n$ over the random choices of the algorithm, the algorithm incurs cost $O(\text{OPT}_s/\beta)$ and outputs a matching of cardinality at least $(1 - \beta)n$, where $\text{OPT}_s = \sum_{E_1} C_e + \sum_{E_2} C_e^s$.

Finally, we can show two hardness of approximation results for the explicit scenario case.

Theorem 4 (Stochastic Optimization Lower Bound). (1) There exists a constant $c > 0$ such that Expression OPT_3 (Eq. (1)) is NP-hard to approximate within a factor of $c \ln n$.

(2) Expression OPT_3 (Eq. (1)) is NP-hard to compute, even when there are only two scenarios and τ is bounded.

Proof. Part 1. We will prove that when $\tau \geq n^2$, Expression (1) is at least as hard to approximate as Minimum Set Cover: given a universe $S = \{s_1, \dots, s_n\}$ of elements and a collection $C = \{c_1, \dots, c_k\}$ of subsets of S , find a minimum-cardinality subset \mathcal{SC} of C such that for every $1 \leq i \leq n$, $s_i \in c_j$ for some $c_j \in \mathcal{SC}$. It is known that there exists a constant $c > 0$ such that approximating Minimum Set-Cover to within a factor of $c \ln n$ is NP-hard [18].

Given an instance $(S = \{s_1, \dots, s_n\}; C = \{c_1, \dots, c_k\})$ of Minimum Set-Cover, we construct an instance of the two-stage matching problem with stochastic matching vertices as follows. The construction is illustrated in Fig. 2. The graph contains $|S| + 3|C|$ vertices: for every element $s_i \in S$ there is a vertex u_i ; for every set $c_j \in C$, there are three vertices x_j, y_j , and z_j connected by a path $(x_j, y_j), (y_j, z_j)$. For every set c_j and element s_i which belongs to c_j , we have the edge (z_j, u_i) . It is easy to see that the graph is bipartite. The first-stage edge costs are 1 for an (x_i, y_i) edge costs and 0 for the other edges. The second-stage costs are equal to the first-stage costs, multiplied by τ . There are n equally likely second-stage scenarios: in scenario i the vertices in $\{y_1, \dots, y_k\} \cup \{u_i\}$ are active.

Consider a set cover \mathcal{SC} for the input instance. In the first stage, buy edge (x_j, y_j) for each set $c_j \in \mathcal{SC}$. In the second stage, let i be the scenario and let c_j be a set in the set cover that contains the element s_i . Buy the edge (z_j, u_i) and every edge (y_j, z_j) for $j \neq j$. The edges bought contain a matching that matches all active vertices, and the total cost equals $|\mathcal{SC}|$.

On the other hand, assume that the edges bought in the first stage do not correspond to a set cover of the input instance. Let i be an element which is not covered. Then in scenario i , the algorithm will have to match u_i with some z_j such that $u_i \in c_j$, and then it will have to buy the edge (x_j, y_j) at cost n^2 . Thus the expected cost is at least $n^2/n = n$. Thus the minimum of Expression (1) is exactly equal to the cardinality of the minimum set cover of the input instance.

Part 2. By reduction from the NP-hard Simultaneous Matchings problem [7]: We are given a bipartite graph $G = (X, Y, E)$ and two subsets Z_1, Z_2 of X , such that G has a matching that matches every vertex of Z_i for each $i = 1, 2$. We need to find a minimum cardinality edge-set $M \subseteq E$ such that for each $i = 1, 2$, the restriction of M to $Z_i \times Y$ is a Z_i -perfect matching.

Given an instance of the Simultaneous Matchings problem, we create an instance of our problem as follows. The graph is $G' = (Y, X, E)$. Each edge has cost 1 in the first stage τ in the second stage. There are two equally-likely second-stage scenarios: in scenario i , the vertices of Z_i are active. We show that the instance we created has a solution of cost $\leq |X|$ if and only if the Simultaneous Matching instance has a solution of cardinality $|X|$.

For the first direction, assume that the Simultaneous Matching instance has a solution M of cardinality $|X|$, and consider an algorithm that buys the edges of M in the first stage. The bought edges contain a matching for each second-stage scenario, so the total cost is equal to the first-stage cost, i.e., $|X|$. Conversely, assume that the maximum simultaneous matching in the graph has cardinality smaller than $|X|$ and assume that an algorithm bought the edge-set \mathcal{M}_1 in the first stage. If $|\mathcal{M}_1| = |X|$, then in at least one of the second-stage scenarios, the matching we can create with the edges of \mathcal{M}_1 does not match all of




Fig. 3. The graph obtained from the 3-set-cover(2) instance $\{s_1, s_2, s_3\}, \{s_1, s_3, s_4\}, \{s_2, s_5, s_6\}, \{s_4, s_5, s_6\}$.

the active vertices, so with probability $1/2$ we will have to buy at least one edge in the second stage, at cost τ . Finally, if $|\mathcal{M}1| < |X|$ then there are $|X| - |\mathcal{M}1|$ vertices in X that are not incident on any edge in $\mathcal{M}1$. In the second stage, each of these vertices will be active with probability at least $1/2$, and in this case we will buy an edge matching it at cost τ . The total cost, then, is at least $|\mathcal{M}1| + (|X| - |\mathcal{M}1|)\tau/2 > |X|$. \square

3. Implicit scenarios

Instead of having an explicit list of scenarios for the second stage, it is common to have instead an implicit description: in the case of uncertain activated vertices, a natural stochastic model is the one in which each vertex is active in the second stage with some probability p , independently of the status of the other nodes. Due to independence, we get that although the total number of possible scenarios can be exponentially large, there is a succinct description consisting of simply specifying the activation probability of each node. In this case, we can no longer be certain that the second-stage graph contains a perfect matching, even if the input graph does, so the requirement is, as stated above, to find the largest possible matching. We first prove an interesting lower bound.

3.1. Lower bounds

Theorem 5. *Stochastic optimization with uncertain vertex set is NP-hard to approximate within a certain constant, even with independent vertex activation.*

Proof. We detail this proof, which is the most interesting of our lower bounds. We will use a reduction from Minimum 3-Set-Cover(2), the special case of Minimum Set-Cover where each set has cardinality 3 and each element belongs to two sets [16]. This variant is NP-hard to approximate to within a factor of $100/99$ [3]. We will prove that approximating Expression (1) to within a factor of β is at least as hard as approximating 3-set-cover(2) to within a factor of $\gamma = \beta(1 + (3p^2(1-p) + 2p^3)\tau)$. The theorem follows by setting p to be a constant in the interval $[0, 0.0033]$ and $\tau = 1/p$, because then $3p(1-p) + 2p^2 < 1/99$.

Given an instance $(S = \{s_1, \dots, s_n\}; C = \{c_1, \dots, c_k\})$ of 3-set-cover(2), we construct an instance of the two-stage matching problem with uncertain activated vertices as follows (see Fig. 3). The graph contains $2|S| + 3|C|$ vertices: for every element $s_i \in S$ there are two vertices u_i, u'_i connected by an edge; for every set $c_j \in C$, there are three vertices x_j, y_j , and z_j connected by a path $(x_j, y_j), (y_j, z_j)$. For every set c_j and element s_i which belongs to c_j , we have the edge (z_j, u_i) . It is easy to see that the graph is bipartite. The first-stage edge costs are 1 for an (x_j, y_j) or (u_i, u'_i) edge and 0 for the other edges. The second-stage costs are equal to the first-stage costs, multiplied by τ . In the second-stage scenarios, each vertex u_i is active with probability p and each y_j is active with probability 1.

If $p > 1/\tau$, then buying all (u_i, u'_i) edges in the first stage at cost n is optimal. To see why, assume that an algorithm spends $n' < n$ in the first stage. In the second stage, the expected number of active vertices that cannot be matched is at least $(n - n')p$ and the expected cost of matching them is $\tau(n - n')p > (n - n')$. We assume in the following that $p \leq 1/\tau$.

Consider a minimum set cover \mathcal{SC} of the input instance. Assume that in the first stage we buy (at cost 1) the edge (x_j, y_j) for every set $c_j \in \mathcal{SC}$. In the second stage, let I be the set of active vertices and find, in a way to be described shortly, a matching M_I between a subset I' of I and the vertex-set $\{z_j : c_j \in \mathcal{SC}\}$, using (z_j, u_i) -edges from the graph. Buy the edges in M_I (at cost 0). For every $i \in I \setminus I'$, buy the edge (u_i, u'_i) at cost τ . Now, all active u_i vertices are matched, and it remains to ensure that the y -vertices are matched as well. Assume that y_j is unmatched. If z_j is matched with some u_i node, this is because $c_j \in \mathcal{SC}$, so we bought the edge (x_j, y_j) in the first stage and can now use it at no additional cost. Otherwise, we buy the edge (y_j, z_j) at cost 0. The second stage has cost equal to τ times the cardinality of $I \setminus I'$ and the first stage has cost equal to the cardinality of the set cover. The matching M_I is found in a straightforward manner: given \mathcal{SC} , each element chooses exactly one set among the sets covering it, and, if it turns out to be active, will only try to be matched to that set. Each set in the set cover will be matched with one element, chosen arbitrarily among the active vertices who try to be matched with it.

To calculate the expected cost of matching the vertices of $I \setminus I'$, consider a set in \mathcal{SC} . It has 3 elements, and is chosen by at most 3 of them. Assume that it is chosen by all 3. With probability $(1 - p)^3 + 3p(1 - p)^2$, at most one of them is active and no cost is incurred in the second stage. With probability $3p^2(1 - p)$, two of them are active and a cost of τ is incurred.

With probability p^3 , all three of them are active and a cost of 2τ is incurred, for an expected cost of $(3p^2(1-p) + 2p^3)\tau$. If the set is chosen by two elements, the expected cost is at most $p^2\tau$, and if it is chosen by fewer, the expected cost is 0. Thus in all cases the expected cost of matching $I \setminus I'$ is bounded by $|\mathcal{SC}|(3p^2(1-p) + 2p^3)\tau$. With a cost of $|\mathcal{SC}|$ for the first stage, we get that the total cost of the solution is at most $|\mathcal{SC}|(1 + (3p^2(1-p) + 2p^3)\tau)$.

On the other hand, let \mathcal{M}_1 be the set of cost-1 edges bought in the first stage. Let an (x_i, y_i) edge represent the set c_i and let a (u_i, u'_i) edge represent the singleton set $\{s_i\}$. Now, assume that \mathcal{M}_1 does not correspond to a set cover of the input instance. Let x be the number of elements which are not covered by the sets corresponding to \mathcal{M}_1 and let X be the number of active elements among those x . In the second stage, the algorithm will have to match each uncovered element vertex u_i , either by its (u_i, u'_i) edge (at cost n) or by a (z_j, u_i) edge for some set c_j where $s_i \in c_j$. In the latter case, it would have to buy the edge (x_i, y_i) , again at cost n . The second stage cost, therefore, is at least Xn . But the expected value of X is x/n , thus the total expected cost is at least $|\mathcal{M}_1| + x$. Since we could complete \mathcal{M}_1 into a set cover by adding at most one set per uncovered element, we have $x + |\mathcal{M}_1| \geq |\mathcal{SC}|$.

In summary, we get that Expression (1) satisfies

$$|\mathcal{SC}| \leq \text{OPT} \leq |\mathcal{SC}|(1 + (3p^2(1-p) + 2p^3)\tau).$$

This means that if we can approximate our problem within a factor of β , then we can approximate Minimum 3-Set-Cover(2) within a factor of $\gamma = \beta(1 + (3p^2(1-p) + 2p^3)\tau)$, and the theorem follows. \square

Using similar ideas, we prove the following related result.

Theorem 6. *The case of uncertain, independent, edge costs is NP-hard to approximate within a certain constant.*

Proof. Once again, we use a reduction from 3-set-cover(2). Consider an instance of 3-set-cover(2). We define a graph that, for every element $s_i \in S$, contains two vertices u_i and u'_i joined by an edge (u_i, u'_i) . For every set $c_j \in C$, it contains three vertices x_j, y_j , and z_j and the path $(x_j, y_j), (y_j, z_j)$. Additionally, for every set c_j and element s_i such that $s_i \in c_j$, we have the edge (z_j, u_i) .

The edge-costs are as follows. In the first stage, the (x_j, y_j) and (u_i, u'_i) edges have cost 1 and all other edges have cost 0. In the second stage, an (x_j, y_j) edge has cost τ , and each (u_i, u'_i) edge has cost τ with probability p and 0 with probability $1-p$. The other edges have a second-stage cost of 0. Note that when the edge (u_i, u'_i) has cost 0 in the second stage, then u_i can be matched at cost 0. This is equivalent to u_i being inactive in the uncertain matching targets case, and therefore the rest of the proof is similar to the proof of Theorem 5. \square

3.2. Upper bound in a special case

We can show that when $c_e = 1$ for all $e \in E$, it is possible to construct a perfect matching cheaply when the graph has certain properties. We study the case in which B is significantly larger than A .

Theorem 7. *Assume that the graph contains n vertex-disjoint stars s_1, \dots, s_n such that star s_i contains $d = \max\{1, \ln(\tau p)\}/\ln(1/(1-p)) + 1$ vertices from B and is centered at some vertex of A . Then there is an algorithm whose running time is polynomial in n , and which returns a maximum-cardinality matching of the second stage graph, whose expected cost is $O(\text{OPT}_3 \cdot \min\{1, \ln(\tau p)\})$.*

To prove this theorem, let $A = \{a_1, \dots, a_n\}$ and $B = \{b_1, \dots, b_m\}$. Let B_2 be the vertices which are active in the second stage. Here is the algorithm.

First stage:

If $\tau p \leq e$ then
buy nothing
else

buy E_1 , the set of nd edges in the stars.

Second stage: complete the set of edges into a perfect matching in the cheapest way possible.

Output: the perfect matching.

To analyze the algorithm, we say that a_i is *miserable* if none of the vertices in s_i are active and that it is *poor* if exactly one vertex in s_i is active. Let A_m be the set of miserable vertices and A_p the set of poor vertices. The following Lemma is the key of the analysis to constructing a perfect matching, and so we give its proof in detail.

Lemma 4. *There exists a maximum-cardinality matching M^* in G_2 such that*

$$|M^* \setminus E_1| \leq 2|A_m| + |A_p|.$$




Fig. 4. Illustration of the proof of Lemma 4.

Proof. Let M^* be a maximum matching in G_2 that has the maximum number of edges from E_1 . Let M be a maximum matching that uses only edges from E_1 . The edge-set $M \oplus M^*$ is a collection of vertex-disjoint odd-length paths, each of which connects a vertex a_i of A with a vertex b_j of B and is denoted $P(a_i, b_j)$; both a_i and b_j are unmatched in M . Since vertex a_i is unmatched in M , it must be that it is miserable. For each other vertex $a_k \in A \cap P(a_i, b_j)$, let (a_k, b_k) be the matching edge in M and (a_k, b_{k+1}) be the matching edge in M^* . If a_k is not poor, then there is another vertex $b_{k'}$ in the star centered on a_k , which is active but not matched in M . If $b_{k'}$ is not matched in M^* , then $(M^* \setminus \{(a_k, b_{k+1})\}) \cup \{(a_k, b_{k'})\}$ would be another maximum matching in G_2 with one more edge from E_1 , contradicting the definition of M^* . Thus $b_{k'}$ is matched in M^* , but not in M . Let $P(a_{i'}, b_{k'})$ be the path of $M \oplus M^*$ that $b_{k'}$ belongs to: $a_{i'}$ is miserable. In this way, we can associate every rich A -vertex that lies on an alternating path with a unique miserable node. We get that the total number of vertices of A which are along the paths of $M \oplus M^*$ is at most $|A_m| + |A_p| + |A_m|$, hence the lemma.

Fig. 4 illustrates the proof. It shows an alternating path starting at the unmatched, miserable vertex $a \in A$ to a rich vertex $d \in A$. For every rich A -vertex along the path (except for the last), such as the vertex b in the example, there is another alternating path that ends in this node's star. Hence, we can charge the miserable vertex at the head of that path (f in the example) for this rich internal node. \square

The following two lemmas are not difficult.

Lemma 5. If $\tau p \geq e$, then the expected number of miserable vertices is $E|A_m| = n(1-p)/(\tau p)$, and the expected number of poor vertices is $E|A_p| = nd/\tau$. Otherwise, the expected number of miserable vertices is $E(|A_m|) = n(1-p)/e$.

Proof. If $\tau p \geq e$, then a vertex in A is miserable with probability $(1-p)^d = (1-p)e^{-\ln(\tau p)} = (1-p)/(\tau p)$. Hence, the expected number of miserable vertices is n times that quantity. Similarly, a vertex in A is poor with probability $dp(1-p)^{d-1} = dp/(\tau p) = d/\tau$. If $\tau p < e$, then a vertex in A is miserable with probability $(1-p)/e$. The lemma follows. \square

Lemma 6. The optimal cost is at least $(n - E(|A_m|)) \min(\tau, 1/p)$.

Proof. Because of the disjoint star structure, the cardinality Z_2 of the maximum-cardinality matching in G_2 is certainly at least $n - E(|A_m|)$ in expectation. Let F be the set of edges bought by OPT in the first stage and B_F denote the set of endpoints of those edges on the B side. The number of edges of F which can be used in the maximum matching is certainly at most $\sum_{b \in B_F} \chi(b \text{ active})$, and so, the cost paid by OPT in the second stage is at least $\tau(Z_2 - \sum_{b \in B_F} \chi(b \text{ active}))$. Thus:

$$\text{OPT} \geq \min\{|F| + \tau(n - E(|A_m|)) - \tau p|F|, |F|\}.$$

If $\tau p \leq 1$ then this expression is minimized for $|F| = 0$, when its value is $\text{OPT} \geq \tau(n - E(|A_m|))$. Otherwise, the expression is minimized for $|F| = (n - E(|A_m|))/p$. \square

Assume that $\tau p > e$. From Lemmas 4 and 5, it follows that the algorithm has average cost $n(d + 2(1-p)/p + d)$. From Lemmas 5 and 6, it follows that the optimum cost is at least $n(1 - 1/(\tau p))/p$. It follows that the approximation ratio is bounded by

$$\frac{p(d + 2(1-p)/p + d)}{(1 - 1/(\tau p))} = O(1 + dp) = O\left(\ln(\tau p) \frac{p}{\ln(1/(1-p))}\right) = O(\ln(\tau p)).$$

On the other hand, assume that $\tau p \leq e$. Then the algorithm has cost at most $n\tau$ and OPT has cost at least $(n - E(|A_m|))\tau/2$. Since $E|A_m| = n/e$, this is $\Omega(n\tau)$ and so the approximation ratio is $O(1)$.

3.3. Generalization: The black box model

With independently activated vertices, the number of scenarios is extremely large, and so solving an LP of the kind described in previous sections is prohibitively time-consuming. However, in such a situation there is often a *black box* sampling procedure that provides, in polynomial time, an unbiased sample of scenarios; then one can use the SAA method to simulate the explicit scenarios case, and, if the edge cost distributions have bounded second moment, one can extend the analysis so as to obtain a similar approximation guarantee. The main observation is that the value of the LP defined by taking a polynomial number of samples of scenarios tightly approximates the value of the LP defined by taking all possible scenarios. An analysis similar to [5] gives:

Theorem 8. *Consider a two-stage edge stochastic matching problem with (1) a polynomial time unbiased sampling procedure, and (2) edge cost distributions whose second moment is bounded. For any constants $\epsilon > 0$ and $\delta, \beta \in (0, 1)$, there is a polynomial-time randomized algorithm that outputs a matching whose cardinality is at least $(1 - \beta)n$ and, with probability at least $1 - \delta$ (over the choices of the black box and of the algorithm), incurs expected cost $O(\text{OPT}/\beta)$ (where the expectation is over the space of scenarios).*

Acknowledgements

First author was supported in part by NSF awards DMI-0600384 and ONR Award N000140610607. Third author was supported in part by NSF awards CCR-0121154 and DMI-0600384, and ONR Award N000140610607.

References

- [1] J. Birge, F. Louveaux, *Introduction to Stochastic Programming*, Springer, 1997.
- [2] M. Charikar, C. Chekuri, M. Pal, Sampling bounds for stochastic optimization, in: APPROX-RANDOM, 2005, pp. 257–269.
- [3] M. Chlebík, J. Chlebíková, Inapproximability results for bounded variants of optimization problems, in: FCT, in: LNCS, vol. 2751, 2003, pp. 27–38.
- [4] K. Dhamdhere, V. Goyal, R. Ravi, M. Singh, How to pay, come what may: Approximation algorithms for demand-robust covering problems, in: FOCS, 2005, pp. 367–378.
- [5] K. Dhamdhere, R. Ravi, M. Singh, On two-stage stochastic minimum spanning trees, in: IPCO, in: LNCS, vol. 3509, 2005, pp. 321–334.
- [6] S. Dye, L. Stougie, A. Tomasgard, The stochastic single resource service-provision problem, *Naval Research Logistics* 50 (2003) 257–269.
- [7] K.M. Elbassioni, I. Katriel, M. Kutz, M. Mahajan, Simultaneous matchings, in: ISAAC, in: LNCS, vol. 3827, 2005, pp. 106–115.
- [8] A.D. Flaxman, A.M. Frieze, M. Krivelevich, On the random 2-stage minimum spanning tree, in: SODA, 2005, pp. 919–926.
- [9] A. Gupta, M. Pál, Stochastic steiner trees without a root, in: ICALP, in: LNCS, vol. 3580, 2005, pp. 1051–1063.
- [10] A. Gupta, M. Pál, R. Ravi, A. Sinha, Boosted sampling: Approximation algorithms for stochastic optimization, in: STOC, ACM, 2004, pp. 417–426.
- [11] A. Gupta, R. Ravi, A. Sinha, An edge in time saves nine: LP rounding approx. algorithms for stochastic network design, in: FOCS, 2004, pp. 218–227.
- [12] N. Immerlica, D. Karger, M. Minkoff, V.S. Mirrokni, On the costs and benefits of procrastination: Approximation algorithms for stochastic combinatorial optimization problems, in: SODA, 2004, pp. 691–700.
- [13] C. Kenyon, E. Rémila, A near-optimal solution to a two-dimensional cutting stock problem, *Math. Oper. Res.* 25 (4) (2000) 645–656.
- [14] N. Kong, A.J. Schaefer, A factor 1/2 approximation algorithm for two-stage stochastic matching problems, *Eur. J. Oper. Res.* 172 (2006) 740–746.
- [15] E. Lawler, *Combinatorial Optimization: Networks and Matroids*, Holt, Rinehart, Winston, 1976.
- [16] C.H. Papadimitriou, M. Yannakakis, Optimization, approximation, and complexity classes, *J. Comput. System Sci.* 43 (1991) 425–440.
- [17] R. Ravi, A. Sinha, Hedging uncertainty: Approximation algorithms for stochastic optimization problems, in: IPCO, in: LNCS, vol. 3064, 2004, pp. 101–115.
- [18] R. Raz, S. Safra, A sub-constant error-prob. low-degree test, and a sub-constant error-prob. PCP characterization of NP, in: STOC, 1997, pp. 475–484.
- [19] D.B. Shmoys, M. Sozio, Approximation algorithms for 2-stage stochastic scheduling problems, in: IPCO, 2007.
- [20] D.B. Shmoys, C. Swamy, The Sample Average Approximation Method for 2-Stage Stochastic Optimization, 2004.
- [21] D.B. Shmoys, C. Swamy, Stochastic optimization is almost as easy as deterministic optimization, in: FOCS, 2004, pp. 228–237.
- [22] C. Swamy, D.B. Shmoys, The sampling-based approximation algorithms for multi-stage stochastic optimization, in: FOCS, 2005, pp. 357–366.
- [23] C. Swamy, D.B. Shmoys, Algorithms column: Approximation algorithms for 2-stage stochastic optimization problems, *ACM SIGACT News* 37 (1) (2006) 33–46.
- [24] B. Verweij, S. Ahmed, A.J. Kleywegt, G. Nemhauser, A. Shapiro, The sample average approximation method applied to stochastic routing problems: A computational study, *Comput. Optim. Appl.* 24 (2003) 289–333.