

CAHIER DU LAMSADE

351

Mars 2014

A note on the asymmetric part
of an outranking relation

Denis Bouyssou, Marc Pirlot

A note on the asymmetric part of an outranking relation

Denis Bouyssou¹ Marc Pirlot²

19 March 2014

Abstract

This note discusses the properties of the asymmetric part of an outranking relation à la ELECTRE. Such relations have been used in the pseudo-disjunctive version of ELECTRE TRI-B. We show that they have properties that are somewhat different from the ones of outranking relations. Indeed, contrary to outranking relations, they allow to have at the same time veto and bonus effects. We suggest that this explains the difficulty in analyzing the properties of the pseudo-disjunctive version of ELECTRE TRI-B and the complexity of the elicitation methods of its parameters. We give conditions that characterize such relations.

Key Words: Decisions with multiple attributes, Outranking relations, Concordance, Discordance, Axiomatic analysis.

¹ CNRS, LAMSADE, UMR 7243 & Université Paris Dauphine, Place du Maréchal de Lattre de Tassigny, F-75775 Paris Cedex 16, France, tel: +33 1 44 05 48 98, fax: +33 1 44 05 40 91, e-mail: bouyssou@lamsade.dauphine.fr.

² Université de Mons, Faculté Polytechnique, 9, rue de Houdain, B-7000 Mons, Belgium, tel: +32 65 374682, fax: +32 65 374689, e-mail: marc.pirlot@umons.ac.be.

Contents

1	Introduction	1
2	Definitions and notation	2
2.1	Binary relations	2
2.2	Product sets	2
2.3	Outranking relations	3
3	Background	6
3.1	Traces	6
3.1.1	Traces on differences	6
3.1.2	Traces on levels	7
3.1.3	Relations between traces	9
3.2	Characterization of outranking relations	10
3.2.1	Concordance relations	10
3.2.2	Outranking relations	12
4	The asymmetric part of an outranking relation	13
4.1	Introduction	13
4.2	Properties of an AP-R-CDR-AT	16
4.3	Additional properties of an AP-R-CDR-AT	18
4.4	Vetoes and bonuses	24
4.5	The main result	26
4.6	Examples	30
5	Discussion	34
	References	35

1 Introduction

Most outranking methods, including the well known ELECTRE methods (Roy, 1968; Roy and Bertier, 1973), compare alternatives evaluated on several attributes using the *concordance / non-discordance* principle. It leads to accepting the proposition that an alternative is “superior” to another if the coalition of attributes supporting it is “sufficiently important” (concordance condition) and there is no attribute that “strongly rejects” it (non-discordance condition). The fact that an alternative is “superior” to another means at least two different things. In the ELECTRE methods, superior means “not worse”. Such methods aim at building a *reflexive* preference relation that is interpreted as an “at least as good as” relation. In other methods, like the TACTIC method (Vansnick, 1986), superior means “strictly better than”. Such methods build an *asymmetric* relation that is interpreted as strict preference.

We have studied, from an axiomatic point of view, both reflexive outranking relations (Bouyssou and Pirlot, 2005a, 2007, 2009) and asymmetric outranking relations (Bouyssou and Pirlot, 2002b, 2012). We have also recently unified and generalized the analysis of both cases in Bouyssou and Pirlot (2013). For other approaches, we refer to Greco, Matarazzo, and Słowiński (2001) and Dubois, Fargier, Perny, and Prade (2003).

These works leave the following problem open. What are the properties of the asymmetric part of a reflexive outranking relation? This question is not only of theoretical interest. Indeed, the pseudo-disjunctive version (also known as “optimistic”, we use here the terminology of Almeida-Dias, Figueira, and Roy, 2010 and Roy, 2002) of ELECTRE TRI-B (Roy and Bouyssou, 1993, Ch. 6) makes use of such a relation, contrary to the pseudo-conjunctive version (also known as “pessimistic”) that uses a reflexive outranking relation. We will show that the asymmetric part of a reflexive outranking relation is a more complex object than an outranking relation since it allows both veto and bonus effects. The idea of veto is classical in outranking methods and refers to a deleted preference due to an excessively large negative difference of performance on some criterion. The idea of bonus is more uncommon. It refers to a preference induced by a large positive difference of performance on some criterion. This idea was already introduced in Bouyssou and Pirlot (2013). We will use it here in a slightly different manner: while the bonus effects in Bouyssou and Pirlot (2013) were unconditional, the ones used here will only occur in specific cases.

The rest of the paper is organized as follows. Section 2 introduces our notation and definitions concerning binary relations, product sets and outranking relations. Section 3 presents background material on traces and our previous characterizations of outranking relations. Section 4 presents our main results. A final section concludes and discusses our findings.

2 Definitions and notation

2.1 Binary relations

A preference relation on a set X is in general denoted by \mathcal{R} . Omitting obvious quantifications, we say that a binary relation \mathcal{R} on X is *reflexive* if $a \mathcal{R} a$. It is *complete* if $a \mathcal{R} b$ or $b \mathcal{R} a$. It is *antisymmetric* if $[a \mathcal{R} b \text{ and } b \mathcal{R} a] \Rightarrow a = b$. It is *asymmetric* if $a \mathcal{R} b \Rightarrow \text{Not}[b \mathcal{R} a]$. It is *transitive* if $[a \mathcal{R} b \text{ and } b \mathcal{R} c] \Rightarrow a \mathcal{R} c$. It is *Ferrers* if $[a \mathcal{R} b \text{ and } c \mathcal{R} d] \Rightarrow [a \mathcal{R} d \text{ or } c \mathcal{R} b]$. It is *semi-transitive* if $[a \mathcal{R} b \text{ and } b \mathcal{R} c] \Rightarrow [a \mathcal{R} d \text{ or } d \mathcal{R} c]$.

A *weak order* is a complete transitive relation. A *semiorder* is a reflexive, Ferrers and semi-transitive relation. As first observed by Luce (1956), any semiorder \mathcal{R} on X induces a weak order $T(\mathcal{R})$ on X that is defined, for all $a, b \in X$,

$$a T(\mathcal{R}) b \text{ if } \forall c \in X, [b \mathcal{R} c \Rightarrow a \mathcal{R} c] \text{ and } [c \mathcal{R} a \Rightarrow c \mathcal{R} b]. \quad (1)$$

Conversely, if \mathcal{R} is reflexive and there is a weak order T such that, for all $a, b \in X$, $a T b \Rightarrow \forall c \in X, [b \mathcal{R} c \Rightarrow a \mathcal{R} c] \text{ and } [c \mathcal{R} a \Rightarrow c \mathcal{R} b]$, then \mathcal{R} is a semiorder.

The asymmetric (resp. symmetric) part of \mathcal{R} is the binary relation \mathcal{R}^α (resp. \mathcal{R}^σ) such that, for all $a, b \in X$ $a \mathcal{R}^\alpha b$ if $[a \mathcal{R} b \text{ and } \text{Not}[b \mathcal{R} a]]$ (resp. $a \mathcal{R}^\sigma b$ if $[a \mathcal{R} b \text{ and } b \mathcal{R} a]$).

2.2 Product sets

The set of alternatives will be denoted by X . As is usual in conjoint measurement this set will be identified with the Cartesian product $\prod_{i=1}^n X_i$ of $n \geq 2$ sets X_i . The set X_i gathers all possible evaluations that an alternative can have on attribute i . The sets X_i are not assumed to be sets of numbers, not even to be ordered sets. The set $\{1, 2, \dots, n\}$ of all attributes will be denoted by N .

For any nonempty subset J of the set of attributes N , we denote by X_J (resp. X_{-J}) the set $\prod_{i \in J} X_i$ (resp. $\prod_{i \in N \setminus J} X_i$). With customary abuse of notation, (x_J, y_{-J}) will denote the element $w \in X$ such that $w_i = x_i$ if $i \in J$ and $w_i = y_i$ otherwise. We sometimes omit braces around sets. For instance, when $J = \{i\}$ we write X_{-i} and (x_i, y_{-i}) .

If \mathcal{R} is a binary relation on X , we say that attribute $i \in N$ is *influential* for \mathcal{R} if there are $x_i, y_i, z_i, w_i \in X_i$ and $x_{-i}, y_{-i} \in X_{-i}$ such that $(x_i, x_{-i}) \mathcal{R} (y_i, y_{-i})$ and $\text{Not}[(z_i, x_{-i}) \mathcal{R} (w_i, y_{-i})]$ and *degenerate* otherwise. A degenerate attribute has no influence whatsoever on the comparison of the elements of X and may be suppressed from N . As in Bouyssou and Pirlot (2005a), in order to avoid unnecessary minor complications, we suppose henceforth that *all attributes in N are influential* for \mathcal{R} . Our results in Section 4 will use the slightly stronger

hypothesis stating that all attributes are influential for the asymmetric part of \mathcal{R} (see Remark 26).

2.3 Outranking relations

The following definition of an outranking relation is taken from Bouyssou and Pirlot (2013). Notice that it is silent about the fact that the relation \mathcal{R} is reflexive or asymmetric.

Definition 1 (Concordance-discordance relations)

Let \mathcal{R} be a binary relation on $X = \prod_{i=1}^n X_i$. We say that \mathcal{R} is a concordance-discordance relation (CDR) if there are:

1. a complete binary relation S_i on each X_i ($i = 1, 2, \dots, n$) (with asymmetric part P_i and symmetric part I_i),
2. an asymmetric binary relation V_i on each X_i ($i = 1, 2, \dots, n$) such that $V_i \subseteq P_i$,
3. a binary relation \supseteq between subsets of N having N for union that is monotonic w.r.t. inclusion, i.e., for all $A, B, C, D \subseteq N$ with $A \cup B = N$ and $C \cup D = N$,

$$[A \supseteq B, C \supseteq A, B \supseteq D] \Rightarrow C \supseteq D, \quad (2)$$

such that, for all $x, y \in X$,

$$x \mathcal{R} y \Leftrightarrow [S[x, y] \supseteq S[y, x] \text{ and } V[y, x] = \emptyset], \quad (3)$$

where $S[x, y] = \{i \in N : x_i S_i y_i\}$ and $V[x, y] = \{i \in N : x_i V_i y_i\}$. We say that $\langle \supseteq, S_i, V_i \rangle$ is a representation of \mathcal{R} as a CDR.

A concordance relation (CR) is a CDR having a representation in which all relations V_i are empty.

A CDR with attribute transitivity (CDR-AT) is a CDR for which, for all $i \in N$,

- S_i is a semiorder
- V_i is the asymmetric part of a semiorder U_i
- (S_i, U_i) form a homogeneous chain of semiorders, i.e., there is a weak order T_i on X_i such that:

$$\begin{aligned} x_i T_i y_i &\Rightarrow \forall z_i \in X_i, [y_i S_i z_i \Rightarrow x_i S_i z_i] \text{ and } [z_i S_i x_i \Rightarrow z_i S_i y_i], \text{ and} \\ x_i T_i y_i &\Rightarrow \forall z_i \in X_i, [y_i U_i z_i \Rightarrow x_i U_i z_i] \text{ and } [z_i U_i x_i \Rightarrow z_i U_i y_i]. \end{aligned} \quad (4)$$

The asymmetric (resp. symmetric) part of \succeq will be denoted by \triangleright (resp. \trianglelefteq). It is easy to show that if \succeq satisfies (2), the same is true for \triangleright .

Remark 2

For the case of *asymmetric* outranking relations Bouyssou and Pirlot (2012) have introduced the following definition¹. Instead of (3) they have stated that

$$x \mathcal{R} y \Leftrightarrow [P[x, y] \succeq' P[y, x] \text{ and } V[y, x] = \emptyset],$$

with P_i an asymmetric binary relation on X_i and \succeq' a relation between disjoint subsets of N . This definition is clearly equivalent to the one used above. This is obvious letting $A \succeq' B \Leftrightarrow (N \setminus B) \succeq (N \setminus A)$ and observing that if \succeq' satisfies (2) then \succeq also satisfies (2) (and vice versa). •

Remark 3

Observe that, when S_i and U_i are complete, the existence of a weak order T_i satisfying (4) implies that both S_i and U_i are semiorders. The above definition could therefore omit to specify that S_i and U_i are semiorders. Moreover, it is not difficult to check that, when S_i and U_i are complete, the first part of (4) may equivalently be written as

$$x_i T_i y_i \Rightarrow \forall z_i \in X_i, [y_i P_i z_i \Rightarrow x_i P_i z_i] \text{ and } [z_i P_i x_i \Rightarrow z_i P_i y_i],$$

and the second part as

$$x_i T_i y_i \Rightarrow \forall z_i \in X_i, [y_i V_i z_i \Rightarrow x_i V_i z_i] \text{ and } [z_i V_i x_i \Rightarrow z_i V_i y_i],$$

where, as before, P_i (resp. V_i) is the asymmetric part of S_i (resp. U_i). This will be useful later. •

The following proposition shows that the above definition is flexible enough to cover both the case of a reflexive outranking relation à la ELECTRE and the case of an asymmetric outranking relation à la TACTIC.

Proposition 4 (Bouyssou and Pirlot, 2013, Proposition 8)

If \mathcal{R} is a CDR with representation $\langle \succeq, S_i, V_i \rangle$, then

1. \mathcal{R} is either reflexive or irreflexive. \mathcal{R} is reflexive iff $N \succeq N$. Otherwise, \mathcal{R} is irreflexive,
2. \mathcal{R} is asymmetric iff \succeq is asymmetric.

¹As noted in Bouyssou and Pirlot (2013), this definition also applies to reflexive outranking relations.

Let us illustrate the flexibility of Definition 1 with two well-known examples.

Example 5 (ELECTRE I, Roy, 1968)

Outranking relations in ELECTRE I are built as follows. Using a real-valued function u_i defined on X_i , and a pair of non-negative thresholds pt_i and vt_i , with $pt_i \leq vt_i$, we define the semiorders S_i' and U_i' letting:

$$\begin{aligned} x_i S_i' y_i &\Leftrightarrow u_i(x_i) \geq u_i(y_i) - pt_i, \\ x_i U_i' y_i &\Leftrightarrow u_i(x_i) \geq u_i(y_i) - vt_i. \end{aligned} \quad (5)$$

The relation V_i' is the asymmetric part of U_i' , i.e., $x_i V_i' y_i \Leftrightarrow [x_i U_i' y_i \text{ and } \text{Not}[y_i U_i' x_i]] \Leftrightarrow u_i(x_i) > u_i(y_i) + vt_i$

In ELECTRE I, the outranking relation \mathcal{R} is determined using positive weights w_i attached to each attribute and a threshold s (with $1/2 \leq s \leq 1$), such that, for all $x, y \in X$,

$$x \mathcal{R} y \Leftrightarrow \frac{\sum_{i \in S'[x,y]} w_i}{\sum_{j \in N} w_j} \geq s \text{ and } V'[y, x] = \emptyset,$$

where $S'[x, y] = \{i \in N : x_i S_i' y_i\}$, and $V'[x, y] = \{i \in N : x_i V_i' y_i\}$.

It is easy to see that a relation \mathcal{R} built with ELECTRE I is a reflexive CDR (i.e., an R-CDR) with attribute transitivity (i.e., an R-CDR-AT) according to Definition 1. Indeed, it suffices to take, for all $i \in N$, $S_i = S_i'$, $V_i = V_i'$, and, for all $A, B \in 2^N$ with $A \cup B = N$, $A \supseteq B \Leftrightarrow (\sum_{i \in A} w_i) / (\sum_{j \in N} w_j) \geq s$. \diamond

Example 6 (TACTIC, Vansnick, 1986)

Outranking relations in TACTIC are built as follows. Using a real-valued function u_i defined on X_i , and a pair of non-negative thresholds pt_i and vt_i , with $pt_i \leq vt_i$, we define the semiorders S_i' and U_i' as in (5). The relation P_i' is defined as the asymmetric part of S_i' , i.e., we have $x_i P_i' y_i \Leftrightarrow [x_i S_i' y_i \text{ and } \text{Not}[y_i S_i' x_i]] \Leftrightarrow u_i(x_i) > u_i(y_i) + vt_i$. The relation V_i' is the asymmetric part of U_i' , i.e., $x_i V_i' y_i \Leftrightarrow [x_i U_i' y_i \text{ and } \text{Not}[y_i U_i' x_i]] \Leftrightarrow u_i(x_i) > u_i(y_i) + vt_i$

An outranking relation \mathcal{R} is defined letting, for all $x, y \in X$,

$$x \mathcal{R} y \Leftrightarrow \left[\sum_{i \in P'[x,y]} w_i > \rho \sum_{j \in P'[y,x]} w_j + \varepsilon \text{ and } V'[y, x] = \emptyset \right],$$

where w_i is a positive weight assigned to attribute i , ρ is a multiplicative threshold with $\rho \geq 1$, ε is a nonnegative additive threshold, $P'[x, y] = \{i \in N : x_i P_i' y_i\}$ and $V'[x, y] = \{i \in N : x_i V_i' y_i\}$.

It is easy to see that a relation \mathcal{R} built with TACTIC is an asymmetric CDR (i.e., an A-CDR) with attribute transitivity (i.e., an A-CDR-AT) according to Definition 1. Indeed, it suffices to take, for all $i \in N$, $S_i = S_i'$, $V_i = V_i'$, and, for all $A, B \in 2^N$ such that $A \cup B = N$, $A \supseteq B \Leftrightarrow \sum_{i \in N \setminus B} w_i > \rho \sum_{j \in N \setminus A} w_j + \varepsilon$. \diamond

3 Background

3.1 Traces

We first recall a few structural definitions taken from Bouyssou and Pirlot (2002a, 2004a,b).

3.1.1 Traces on differences

Our first definition gives conditions under which it is possible to define *complete* traces on differences.

Definition 7

We say that \mathcal{R} satisfies

$$\begin{aligned}
 RC1_i \text{ if } & \left. \begin{array}{c} (x_i, a_{-i}) \mathcal{R} (y_i, b_{-i}) \\ \text{and} \\ (z_i, c_{-i}) \mathcal{R} (w_i, d_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{c} (x_i, c_{-i}) \mathcal{R} (y_i, d_{-i}) \\ \text{or} \\ (z_i, a_{-i}) \mathcal{R} (w_i, b_{-i}), \end{array} \right. \\
 RC2_i \text{ if } & \left. \begin{array}{c} (x_i, a_{-i}) \mathcal{R} (y_i, b_{-i}) \\ \text{and} \\ (y_i, c_{-i}) \mathcal{R} (x_i, d_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{c} (z_i, a_{-i}) \mathcal{R} (w_i, b_{-i}) \\ \text{or} \\ (w_i, c_{-i}) \mathcal{R} (z_i, d_{-i}), \end{array} \right.
 \end{aligned}$$

for all $x_i, y_i, z_i, w_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$.

We say that \mathcal{R} satisfies $RC1$ (resp. $RC2$) if it satisfies $RC1_i$ (resp. $RC2_i$) for all $i \in N$.

Condition $RC1_i$ amounts to say that all preference differences (x_i, y_i) on X_i can be weakly ordered. Condition $RC2_i$ establishes a link between opposite differences of preferences such as (x_i, y_i) and (y_i, x_i) . Note that $RC2$ entails that \mathcal{R} is independent (i.e., that, for all $i \in N$ all $x_i, y_i \in X_i$ and all $a_{-i}, b_{-i} \in X_{-i}$, $(x_i, a_{-i}) \mathcal{R} (x_i, b_{-i}) \Leftrightarrow (y_i, a_{-i}) \mathcal{R} (y_i, b_{-i})$). Since they will be useful in the sequel, we recall the precise definition of the weak orders induced on each attribute X_i as well as the main properties linking them to conditions $RC1$ and $RC2$. It is easy to check that all CDR satisfy conditions $RC1$ and $RC2$.

Definition 8

Let \mathcal{R} be a binary relation on a set $X = \prod_{i=1}^n X_i$. We define the binary relations \succsim_i^* and \succsim_i^{**} on X_i^2 letting, for all $x_i, y_i, z_i, w_i \in X_i$,

$$\begin{aligned}
 (x_i, y_i) \succsim_i^* (z_i, w_i) & \Leftrightarrow \\
 \forall a_{-i}, b_{-i} \in X_{-i}, & [(z_i, a_{-i}) \mathcal{R} (w_i, b_{-i}) \Rightarrow (x_i, a_{-i}) \mathcal{R} (y_i, b_{-i})], \\
 (x_i, y_i) \succsim_i^{**} (z_i, w_i) & \Leftrightarrow [(x_i, y_i) \succsim_i^* (z_i, w_i) \text{ and } (w_i, z_i) \succsim_i^* (y_i, x_i)].
 \end{aligned}$$

These relations allow to give a precise meaning to the comparison of preference differences on each attribute (see Bouyssou and Pirlot, 2002a, for more detail). By construction, both relations \succsim_i^* and \succsim_i^{**} are transitive. The impact of $RC1_i$ and $RC2_i$ is to ensure their completeness, as shown below.

Lemma 9 (Bouyssou and Pirlot, 2002a, Lemma 1)

1. $RC1_i \Leftrightarrow [\succsim_i^* \text{ is complete}]$,
2. $RC2_i \Leftrightarrow$
 $[for\ all\ x_i, y_i, z_i, w_i \in X_i, Not[(x_i, y_i) \succsim_i^* (z_i, w_i)] \Rightarrow (y_i, x_i) \succsim_i^* (w_i, z_i)],$
3. $[RC1_i \text{ and } RC2_i] \Leftrightarrow [\succsim_i^{**} \text{ is complete}]$.

Traces on differences are well-behaved, as detailed below.

Lemma 10 (Bouyssou and Pirlot, 2005b, Lemma 3.7)

For all $x, y \in X$ and all $z_i, w_i \in X_i$,

$$\begin{aligned}
& [(z_i, w_i) \sim_i^* (x_i, y_i) \text{ for all } i \in N] \Rightarrow [x \mathcal{R} y \Leftrightarrow z \mathcal{R} w], \\
& [x \mathcal{R} y \text{ and } (z_i, w_i) \succsim_i^* (x_i, y_i)] \Rightarrow (z_i, x_{-i}) \mathcal{R} (w_i, y_{-i}), \\
& [x \mathcal{R}^\alpha y \text{ and } (z_i, w_i) \succsim_i^{**} (x_i, y_i)] \Rightarrow (z_i, x_{-i}) \mathcal{R}^\alpha (w_i, y_{-i}), \\
& [(z_i, w_i) \sim_i^{**} (x_i, y_i) \text{ for all } i \in N] \Rightarrow \begin{cases} [x \mathcal{R} y \Leftrightarrow z \mathcal{R} w] \\ \text{and} \\ [x \mathcal{R}^\alpha y \Leftrightarrow z \mathcal{R}^\alpha w]. \end{cases}
\end{aligned}$$

3.1.2 Traces on levels

The next definition gives conditions under which it is possible to define *complete* traces on the elements of each attribute.

Definition 11

We say that \mathcal{R} satisfies

$$\begin{aligned}
AC1_i \text{ if } & \left. \begin{array}{c} (x_i, a_{-i}) \mathcal{R} (y_i, b_{-i}) \\ \text{and} \\ (z_i, c_{-i}) \mathcal{R} (w_i, d_{-i}) \end{array} \right\} \Rightarrow \begin{cases} (z_i, a_{-i}) \mathcal{R} (y_i, b_{-i}) \\ \text{or} \\ (x_i, c_{-i}) \mathcal{R} (w_i, d_{-i}), \end{cases} \\
AC2_i \text{ if } & \left. \begin{array}{c} (x_i, a_{-i}) \mathcal{R} (y_i, b_{-i}) \\ \text{and} \\ (z_i, c_{-i}) \mathcal{R} (w_i, d_{-i}) \end{array} \right\} \Rightarrow \begin{cases} (x_i, a_{-i}) \mathcal{R} (w_i, b_{-i}) \\ \text{or} \\ (z_i, c_{-i}) \mathcal{R} (y_i, d_{-i}), \end{cases} \\
AC3_i \text{ if } & \left. \begin{array}{c} (x_i, a_{-i}) \mathcal{R} (y_i, b_{-i}) \\ \text{and} \\ (y_i, c_{-i}) \mathcal{R} (w_i, d_{-i}) \end{array} \right\} \Rightarrow \begin{cases} (x_i, a_{-i}) \mathcal{R} (z_i, b_{-i}) \\ \text{or} \\ (z_i, c_{-i}) \mathcal{R} (w_i, d_{-i}), \end{cases}
\end{aligned}$$

for all $x_i, y_i, z_i, w_i \in X_i$, all $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$.

We say that \mathcal{R} satisfies $AC1$ (resp. $AC2$, $AC3$) if it satisfies $AC1_i$ (resp. $AC2_i$, $AC3_i$) for all $i \in N$.

These conditions are related to the existence of linear arrangements of the elements of X_i . $AC1_i$ suggests that the elements of X_i can be ordered relatively to “upward dominance”: if x_i “upward dominates” z_i , then $(z_i, c_{-i}) \mathcal{R} (w_i, d_{-i})$ entails $(x_i, c_{-i}) \mathcal{R} (w_i, d_{-i})$. $AC2_i$ has a similar interpretation regarding “downward dominance”. $AC3_i$ ensures that the upward and downward dominance orders are not incompatible. It is easy to check that all CDR-AT satisfy conditions $AC1$, $AC2$, and $AC3$.

The following gives a precise definition of the upward and downward dominance relations.

Definition 12

Let \mathcal{R} be a binary relation on a set $X = \prod_{i=1}^n X_i$. We define the binary relations \succsim_i^+ , \succsim_i^- and \succsim_i^\pm on X_i letting, for all $x_i, y_i \in X_i$,

$$x_i \succsim_i^+ y_i \Leftrightarrow \forall a_{-i} \in X_{-i}, b \in X, [(y_i, a_{-i}) \mathcal{R} b \Rightarrow (x_i, a_{-i}) \mathcal{R} b], \quad (6)$$

$$x_i \succsim_i^- y_i \Leftrightarrow \forall a \in X, b_{-i} \in X_{-i}, [a \mathcal{R} (x_i, b_{-i}) \Rightarrow a \mathcal{R} (y_i, b_{-i})], \quad (7)$$

$$x_i \succsim_i^\pm y_i \Leftrightarrow x_i \succsim_i^+ y_i \text{ and } x_i \succsim_i^- y_i. \quad (8)$$

By definition, \succsim_i^+ , \succsim_i^- and \succsim_i^\pm are transitive relations. Conditions $AC1_i$, $AC2_i$ and $AC3_i$ ensure that they are complete, as shown below.

Lemma 13 (Bouyssou and Pirlot, 2004a, Lemma 3)

Let \mathcal{R} be a binary relation on a set $X = \prod_{i=1}^n X_i$. \mathcal{R} satisfies:

1. $AC1_i \Leftrightarrow \succsim_i^+$ is complete,
2. $AC2_i \Leftrightarrow \succsim_i^-$ is complete,
3. $AC3_i \Leftrightarrow [Not[x_i \succsim_i^+ y_i] \Rightarrow y_i \succsim_i^- x_i] \Leftrightarrow [Not[x_i \succsim_i^- y_i] \Rightarrow y_i \succsim_i^+ x_i]$,
4. $[AC1_i, AC2_i \text{ and } AC3_i] \Leftrightarrow \succsim_i^\pm$ is complete.

Traces on levels are also well-behaved, as detailed below.

Lemma 14 (Bouyssou and Pirlot, 2005b, Lemma 3.5)

For all $i \in N$ and $x, y, z, w \in X$:

$$[x \mathcal{R} y, z_i \succsim_i^+ x_i] \Rightarrow (z_i, x_{-i}) \mathcal{R} y,$$

$$[x \mathcal{R} y, y_i \succsim_i^- w_i] \Rightarrow x \mathcal{R} (w_i, y_{-i}),$$

$$[z_i \succsim_i^\pm x_i, y_i \succsim_i^\pm w_i] \Rightarrow \begin{cases} x \mathcal{R} y \Rightarrow (z_i, x_{-i}) \mathcal{R} (w_i, y_{-i}), \text{ and} \\ x \mathcal{R}^\alpha y \Rightarrow (z_i, x_{-i}) \mathcal{R}^\alpha (w_i, y_{-i}), \end{cases}$$

$$[z_i \sim_i^\pm x_i, y_i \sim_i^\pm w_i, \text{ for all } i \in N] \Rightarrow \begin{cases} x \mathcal{R} y \Leftrightarrow z \mathcal{R} w, \text{ and} \\ x \mathcal{R}^\alpha y \Leftrightarrow z \mathcal{R}^\alpha w. \end{cases}$$

3.1.3 Relations between traces

Traces on differences and on levels will be of central importance in what follows. We summarize below the main links between the relations \succsim_i^* , \succsim_i^{**} , \succsim_i^\pm , \succsim_i^+ and \succsim_i^- below.

Lemma 15 (Bouyssou and Pirlot, 2005b, Lemma 3.8)

For all $i \in N$ and all $x_i, y_i \in X_i$,

1. $x_i \succsim_i^+ y_i \Leftrightarrow [(x_i, w_i) \succsim_i^* (y_i, w_i), \text{ for all } w_i \in X_i]$,
2. $x_i \succsim_i^- y_i \Leftrightarrow [(w_i, y_i) \succsim_i^* (w_i, x_i), \text{ for all } w_i \in X_i]$,
3. $x_i \succsim_i^\pm y_i \Leftrightarrow [(x_i, w_i) \succsim_i^{**} (y_i, w_i), \text{ for all } w_i \in X_i]$,
4. $[\ell_i \succsim_i^+ x_i \text{ and } (x_i, y_i) \succsim_i^* (z_i, w_i)] \Rightarrow (\ell_i, y_i) \succsim_i^* (z_i, w_i)$,
5. $[y_i \succsim_i^- \ell_i \text{ and } (x_i, y_i) \succsim_i^* (z_i, w_i)] \Rightarrow (x_i, \ell_i) \succsim_i^* (z_i, w_i)$,
6. $[z_i \succsim_i^+ \ell_i \text{ and } (x_i, y_i) \succsim_i^* (z_i, w_i)] \Rightarrow (x_i, y_i) \succsim_i^* (\ell_i, w_i)$,
7. $[\ell_i \succsim_i^- w_i \text{ and } (x_i, y_i) \succsim_i^* (z_i, w_i)] \Rightarrow (x_i, y_i) \succsim_i^* (z_i, \ell_i)$,
8. $[x_i \sim_i^+ z_i \text{ and } y_i \sim_i^- w_i] \Rightarrow (x_i, y_i) \sim_i^* (z_i, w_i)$,
9. $[x_i \sim_i^\pm z_i \text{ and } y_i \sim_i^\pm w_i] \Rightarrow (x_i, y_i) \sim_i^{**} (z_i, w_i)$.

Bouyssou and Pirlot (2002a, 2004a,b) have shown how the conditions introduced so far enable to obtain various numerical representations of \mathcal{R} even though it may not be complete or transitive.

3.2 Characterization of outranking relations

3.2.1 Concordance relations

The strategy of Bouyssou and Pirlot (2002b, 2005a, 2007, 2009), first presented in Bouyssou, Pirlot, and Vincke (1997) and Bouyssou and Pirlot (1999), to characterize outranking relations is to introduce conditions limiting the number of equivalence classes of \succsim_i^* and \succsim_i^{**} . Indeed, it is easy to see that an outranking relation generates a relation \succsim_i^* that has at most four equivalence classes. The first one corresponds to a strict preference. The second one to an indifference. The third one to a reverse strict preference that is not a veto. The fourth one to a reverse strict preference that is veto. When this relation is a concordance relation, the fourth class must be empty. This is the intuition behind the following two conditions.

Definition 16

We say that \mathcal{R} satisfies

$$\begin{aligned}
 MM1_i \text{ if } & \left. \begin{array}{c} (x_i, a_{-i}) \mathcal{R} (y_i, b_{-i}) \\ \text{and} \\ (z_i, a_{-i}) \mathcal{R} (w_i, b_{-i}) \\ \text{and} \\ (z_i, c_{-i}) \mathcal{R} (w_i, d_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{c} (y_i, a_{-i}) \mathcal{R} (x_i, b_{-i}) \\ \text{or} \\ (w_i, a_{-i}) \mathcal{R} (z_i, b_{-i}) \\ \text{or} \\ (x_i, c_{-i}) \mathcal{R} (y_i, d_{-i}), \end{array} \right. \\
 MM2_i \text{ if } & \left. \begin{array}{c} (x_i, a_{-i}) \mathcal{R} (y_i, b_{-i}) \\ \text{and} \\ (w_i, a_{-i}) \mathcal{R} (z_i, b_{-i}) \\ \text{and} \\ (y_i, c_{-i}) \mathcal{R} (x_i, d_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{c} (y_i, a_{-i}) \mathcal{R} (x_i, b_{-i}) \\ \text{or} \\ (z_i, a_{-i}) \mathcal{R} (w_i, b_{-i}) \\ \text{or} \\ (z_i, c_{-i}) \mathcal{R} (w_i, d_{-i}), \end{array} \right.
 \end{aligned}$$

for all $x_i, y_i, z_i, w_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$.

We say that $MM1$ (resp. $MM2$) holds if $MM1_i$ (resp. $MM2_i$) holds for all $i \in N$.

$MM1_i$ without its second premise is called $M1_i$. $MM1_i$ without its second conclusion is called $Maj1_i$. $MM1_i$ without both its second premise and its second conclusion is called UC_i .

$MM2_i$ without its second premise is called $M2_i$. $MM2_i$ without its second conclusion is called $Maj2_i$. $MM2_i$ without both its second premise and its second conclusion is called LC_i .

$MM1_i$ is clearly a weaker condition than both $M1_i$ and $Maj1_i$. Both $M1_i$ and $Maj1_i$ are implied by UC_i . Similarly, $MM2_i$ is clearly a weaker condition than both $M2_i$ and $Maj2_i$. Both $M2_i$ and $Maj2_i$ are implied by LC_i .

It is not difficult to check that a concordance relation (CR) satisfies *UC* and *LC*. The need for the weaker conditions introduced above is to ensure that all conditions remain independent from the conditions used to impose the existence of complete traces. We have:

Lemma 17 (Bouyssou and Pirlot, 2013, Lemma 41)

The following implications hold, for all $i \in N$:

1. $MM1_i$ and $RC1_i$ entail $M1_i$,
2. $MM1_i$ and $RC2_i$ entail $Maj1_i$,
3. $MM2_i$ and $RC2_i$ entail $M2_i$,
4. $MM2_i$ and $RC1_i$ entail $Maj2_i$.

Under $RC1_i$ and $RC2_i$, we have:

1. $MM1_i \Leftrightarrow M1_i \Leftrightarrow Maj1_i \Leftrightarrow UC_i$,
2. $MM2_i \Leftrightarrow M2_i \Leftrightarrow Maj2_i \Leftrightarrow LC_i$.

Moreover we have:

Lemma 18 (Bouyssou and Pirlot, 2013, Lemma 24)

1. *If \mathcal{R} satisfies $RC2_i$, then \mathcal{R} satisfies $M1_i \Leftrightarrow$
[for all $x_i, y_i, z_i, w_i \in X_i$, $Not[(y_i, x_i) \succsim_i^* (x_i, y_i)] \Rightarrow (x_i, y_i) \succsim_i^* (z_i, w_i)$],*
2. *If \mathcal{R} satisfies $RC1_i$, then \mathcal{R} satisfies $M2_i \Leftrightarrow$
[for all $x_i, y_i, z_i, w_i \in X_i$, $Not[(y_i, x_i) \succsim_i^* (x_i, y_i)] \Rightarrow (z_i, w_i) \succsim_i^* (y_i, x_i)$].*

The above two lemmas show that, in presence of *RC1* and *RC2*, conditions *MM1* and *MM2* imply that there are only one type of positive differences (corresponding to strict preference) and one type of negative differences (corresponding to reverse strict preference). This is a characteristic of outranking relations in which there is no veto, i.e., of concordance relations. We have:

Theorem 19 (Bouyssou and Pirlot, 2013, Theorems 43 & 52)

Let \mathcal{R} be a reflexive binary relation on $X = \prod_{i=1}^n X_i$.

1. *\mathcal{R} is a concordance relation (CR) iff it satisfies $RC1$, $RC2$, $MM1$ and $MM2$.*
2. *\mathcal{R} is a concordance relation with attribute transitivity (CR-AT) iff it satisfies $RC1$, $RC2$, $AC1$, $AC3$, $MM1$, $MM2$.*
3. *In the class of reflexive relations and in the class of asymmetric relations, conditions $RC1$, $RC2$, $AC1$, $AC3$, $MM1$, $MM2$ are independent.*

The above result generalizes the ones obtained in Bouyssou and Pirlot (2005a, 2007, 2012).

3.2.2 Outranking relations

When veto effect are allowed, there can be two types of negative differences: negative difference corresponding to a “normal” reverse preference and negative differences corresponding to a “large” reverse preference, i.e., a negative difference acting as a veto. Taking veto effect into account implies relaxing condition $MM2$. This motivates the condition below.

Definition 20

We say that \mathcal{R} satisfies

$$MM3_i \text{ if } \left. \begin{array}{l} (x_i, a_{-i}) \mathcal{R} (y_i, b_{-i}) \\ \text{and} \\ (w_i, a_{-i}) \mathcal{R} (z_i, b_{-i}) \\ \text{and} \\ (y_i, c_{-i}) \mathcal{R} (x_i, d_{-i}) \\ \text{and} \\ (z_i, e_{-i}) \mathcal{R} (w_i, f_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (y_i, a_{-i}) \mathcal{R} (x_i, b_{-i}) \\ \text{or} \\ (z_i, a_{-i}) \mathcal{R} (w_i, b_{-i}) \\ \text{or} \\ (z_i, c_{-i}) \mathcal{R} (w_i, d_{-i}), \end{array} \right. \quad (9)$$

for all $x_i, y_i, z_i, w_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i}, e_{-i}, f_{-i} \in X_{-i}$.

We say that \mathcal{R} satisfies $MM3$ if it satisfies $MM3_i$, for all $i \in N$.

$M3_i$ is the same condition as $MM3_i$ except that the second premise has been removed. $Maj3_i$ is the same condition as $MM3_i$ except that the second conclusion has been removed.

Let us observe that $MM3_i$ only differs from $MM2_i$ by the adjunction of a premise, implying that $MM3_i$ is a weakening of $MM2_i$.

Let us interpret condition $M3_i$, under the hypothesis that $RC1_i$ holds. Assuming $RC1_i$ amounts to say that \succsim_i^* is complete. Hence if the first two premises of $M3_i$ hold and neither the first nor the third conclusion do, then we have $(x_i, y_i) \succ_i^* (y_i, x_i) \succ_i^* (z_i, w_i)$. In these circumstances, the second conclusion cannot be true, since this would imply that $(z_i, w_i) \succ_i^* (y_i, x_i)$, a contradiction with $(y_i, x_i) \succ_i^* (z_i, w_i)$. Hence, none of the three conclusions holds and $M3_i$ can only be satisfied if it never happens that $(z_i, e_{-i}) \mathcal{R} (w_i, f_{-i})$. This means that the ordered pair (z_i, w_i) represents an unacceptable preference difference, leading to a veto. We have:

Lemma 21 (Bouyssou and Pirlot, 2013, Lemma 55)

The following implications hold:

1. $MM3_i$ and $RC2_i$ entail $M3_i$,
2. $MM3_i$ and $RC1_i$ entail $Maj3_i$,

3. $M3_i$ and $RC1_i$ entail $Maj3_i$,

4. $Maj3_i$ and $RC2_i$ entail $M3_i$.

Under $RC1_i$ and $RC2_i$, we have:

$$MM3_i \Leftrightarrow M3_i \Leftrightarrow Maj3_i.$$

The conditions introduced so far are all what we need to characterize outranking relations.

Theorem 22 (Bouyssou and Pirlot, 2013, Theorems 57 & 61)

Let \mathcal{R} be a binary relation on $X = \prod_{i=1}^n X_i$.

1. The relation \mathcal{R} is a CDR iff it satisfies $RC1$, $RC2$, $AC1$, $MM1$ and $MM3$.
2. The relation \mathcal{R} is a CDR-AT iff it satisfies $RC1$, $RC2$, $AC1$, $AC2$, $AC3$, $MM1$ and $MM3$.
3. Conditions $RC1$, $RC2$, $AC1$, $AC2$, $AC3$, $MM1$ and $MM3$ are independent in the class of reflexive relations as well as in the class of asymmetric relations.

The above result generalizes the ones obtained in Bouyssou and Pirlot (2009, 2012).

We now have all the necessary background to study the properties of the asymmetric part of a reflexive CDR-AT (i.e., is an R-CDR-AT). Indeed, if the CDR-AT is already asymmetric (i.e., is an A-CDR-AT), there is little point studying its asymmetric part.

4 The asymmetric part of an outranking relation

4.1 Introduction

Let \mathcal{R} be a reflexive CDR (i.e., an R-CDR) with representation $\langle \succeq, S_i, V_i \rangle$. Using the definition of an R-CDR, its asymmetric part, denoted by \mathcal{R}^α , is such that, for all $x, y \in X$,

$$\begin{aligned} x \mathcal{R}^\alpha y &\Leftrightarrow [x \mathcal{R} y \text{ and } \text{Not}[y \mathcal{R} x]] \\ &\Leftrightarrow \begin{cases} [S[x, y] \succeq S[y, x] \text{ and } V[y, x] = \emptyset] \\ \text{and} \\ [\text{Not}[S[y, x] \succeq S[x, y]] \text{ or } V[x, y] \neq \emptyset] . \end{cases} \end{aligned}$$

Such a relation \mathcal{R}^α is the asymmetric part of an R-CDR. We say that \mathcal{R}^α is an AP-R-CDR. If, furthermore, the relation \mathcal{R} is an R-CDR-AT, we say that \mathcal{R}^α is an AP-R-CDR-AT.

It is easy to see that the above definition can equivalently be written as:

$$x \mathcal{R}^\alpha y \Leftrightarrow [S[x, y] \triangleright S[y, x] \text{ and } V[y, x] = \emptyset] \text{ or} \quad (10)$$

$$\left[S[x, y] \triangleq S[y, x] \text{ and } V[y, x] = \emptyset \text{ and } V[x, y] \neq \emptyset \right]. \quad (11)$$

The first of these conditions is similar to the one defining a CDR. The second one is more complex. In the asymmetric part of an R-CDR, the relation V_i may play the rôle of a veto (discarding a pair belonging to the asymmetric part of the concordance relation) or, on the contrary, play the rôle of a *bonus* (transforming an indifference, w.r.t. the concordance relation, into a strict preference). Note that the “bonus effect” just described is different from the one at work in the *concordance relations with bonus* studied in Bouyssou and Pirlot (2013, Section 4.7). In the latter, a bonus effect shows up and results in a preference as soon as there is a large enough difference of preference on a single attribute. In the present case, the bonus effect is conditional: it only shows up in case a pair of alternatives are indifferent w.r.t. the concordance relation and there is no veto against preference.

Remark 23

The relations we study in this paper have been defined in an indirect way, namely, as the asymmetric part of a concordance-discordance relation. A direct definition of such relations can easily be obtained from Definition 1. We just have to specify that the relation \mathcal{R} is asymmetric and to substitute the condition (3), for having $x \mathcal{R} y$, by conditions (10) and (11). •

Remark 24

It is easy to see that there are two simple cases in which the asymmetric part of a reflexive CDR-AT is an asymmetric CDR-AT.

The first case is trivial: if the reflexive CDR-AT is antisymmetric, i.e., when the relation \triangleq is empty, except that $N \triangleq N$, the asymmetric part of \mathcal{R} is identical to \mathcal{R} except for loops. It is then easy to check that the asymmetric part of \mathcal{R} is an asymmetric CDR-AT.

The second case is also trivial. When the relation \mathcal{R} is a reflexive CDR-AT has a representation in which all relations V_i are empty (i.e., is a CR-AT), it is easy to see that the asymmetric part of \mathcal{R} is an asymmetric CDR-AT (that is also a CR-AT). •

Indeed, in general, the asymmetric part of a reflexive CDR-AT is not an asymmetric CDR-AT. This is illustrated below.

Example 25

Let $N = \{1, 2, 3, 4\}$ and $X = \{x_1, y_1, z_1\} \times \{x_2, y_2\} \times \{x_3, y_3\} \times \{x_4, y_4\}$. Let \mathcal{R} on X be such that

$$x \mathcal{R} y \Leftrightarrow |S[x, y]| \geq |S[y, x]| \text{ and } V[y, x] = \emptyset,$$

where $S[x, y] = \{i \in N : x_i S_i y_i\}$ and $V[x, y] = \{i \in N : x_i V_i y_i\}$ with:

$$\begin{aligned} x_1 P_1 y_1 P_1 z_1, \\ x_2 P_2 y_2, x_3 P_3 y_3, x_4 P_4 y_4, \\ V_i = \emptyset, \text{ for } i \in \{2, 3, 4\} \text{ and } x_1 V_1 z_1. \end{aligned}$$

(where P_i denotes the asymmetric part of the complete relation S_i).

By construction, \mathcal{R} is a reflexive CDR-AT. Denoting by \mathcal{R}^α the asymmetric part of \mathcal{R} , it is easy to check that we have:

$$\begin{aligned} (x_1, x_2, x_3, y_4) \mathcal{R}^\alpha (z_1, x_2, x_3, x_4) \text{ and } \text{Not}[(x_1, x_2, x_3, y_4) \mathcal{R}^\alpha (y_1, x_2, x_3, x_4)], \\ (x_1, x_2, x_3, x_4) \mathcal{R}^\alpha (y_1, y_2, x_3, x_4) \text{ and } \text{Not}[(y_1, x_2, x_3, x_4) \mathcal{R}^\alpha (x_1, y_2, x_3, x_4)]. \end{aligned}$$

The first line implies that, w.r.t. \mathcal{R}^α , the difference (x_1, z_1) is strictly larger (w.r.t. \succsim_1^*) than the difference (x_1, y_1) . The second line shows that the difference (x_1, y_1) is strictly larger (w.r.t. \succsim_1^*) than its opposite (y_1, x_1) . It is easy to see that this is impossible in a CDR (see Bouyssou and Pirlot, 2012, Lemma 9.1, for details). \diamond

The problem we would like to study is the following. Suppose that we are given an asymmetric relation \mathcal{P} on X . Under what conditions is there a reflexive CDR-AT \mathcal{R} on X such that $\mathcal{R}^\alpha = \mathcal{P}$? In other words, under what conditions can we be sure that \mathcal{P} is the asymmetric part of a reflexive CDR-AT?

Some of the conditions characterizing an R-CDR-AT also hold for an AP-R-CDR-AT. Others will have to be weakened.

In the rest of this section, in order to avoid the superscript α , we use \mathcal{P} to denote an asymmetric relation. When \mathcal{P} is an AP-R-CDR-AT, the R-CDR-AT of which \mathcal{P} is the asymmetric part will be denoted by \mathcal{S} . In the rest of this section, unless otherwise specified, the relations \succsim_i^* , \succsim_i^{**} , \succsim_i^+ , \succsim_i^- and \succsim_i^\pm always refer to the relation \mathcal{P} .

Remark 26

For an AP-R-CDR \mathcal{P} that is the asymmetric part of an R-CDR \mathcal{S} , assuming that all attributes are influential for \mathcal{S} is in general insufficient for guaranteeing the corresponding property for \mathcal{S}^α . It is clear that if an attribute is influent for \mathcal{S}^α it must be influent for \mathcal{S} . Indeed the influence of $i \in N$ for \mathcal{S}^α implies that there are $x_i, y_i, z_i, w_i \in X_i$ and $x_{-i}, y_{-i} \in X_{-i}$ such that $(x_i, x_{-i}) \mathcal{S}^\alpha (y_i, y_{-i})$ and $\text{Not}[(z_i, x_{-i}) \mathcal{S}^\alpha (w_i, y_{-i})]$. Hence we have $(x_i, x_{-i}) \mathcal{S} (y_i, y_{-i})$ and $\text{Not}[(y_i, y_{-i}) \mathcal{S} (z_i, x_{-i})]$.

$(x_i, x_{-i})]$ together with either $\text{Not}[(z_i, x_{-i}) \mathcal{S} (w_i, y_{-i})]$ or $(w_i, y_{-i}) \mathcal{S} (z_i, x_{-i})$. In either case we easily conclude that $i \in N$ is influential for \mathcal{S} .

The converse is not true. Indeed, an attribute which only impact would be to transform some pairs belonging to the symmetric part \mathcal{S}^σ of \mathcal{S} into pairs that are incomparable w.r.t. \mathcal{S} , is influential for \mathcal{S} but is degenerate for \mathcal{S}^α .

The condition that all attributes are influential for \mathcal{S}^α is therefore slightly stronger than the one stating that all attributes are influential for \mathcal{S} . Our results on AP-R-CDR-AT will use this slightly stronger hypothesis. \bullet

4.2 Properties of an AP-R-CDR-AT

The main difference between an AP-R-CDR-AT and an R-CR-AT is that with an AP-R-CDR-AT there may exist bonus effects as well as veto effects. Bonus effects amount to creating a new class of positive preference differences. Hence it should not be a surprise that we have to relax condition *MM1*. However, as shown in the next lemma, all other conditions needed to characterize R-CDR-AT also hold for AP-R-CDR-AT.

Lemma 27

If \mathcal{P} is an AP-R-CDR-AT, then

1. \mathcal{P} satisfies *RC1* and *RC2*,
2. \mathcal{P} satisfies *AC1*, *AC2*, and *AC3*,
3. \mathcal{P} satisfies *Maj3* and, hence, *MM3*.

PROOF

The fact that \mathcal{P} is an AP-R-CDR means that there is an R-CDR, \mathcal{S} , with representation $\langle \succeq, S_i, V_i \rangle$ such that \mathcal{P} is the asymmetric part of \mathcal{S} . Alternatively, one may use the definition introduced in Remark 23.

Part 1. [RC1] Suppose that $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$, $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$. We know that $\text{Not}[y_i V_i x_i]$ and $\text{Not}[w_i V_i z_i]$. If $x_i V_i y_i$, since $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$, we must have $(x_i, c_{-i}) \mathcal{P} (y_i, d_{-i})$. Similarly, if $z_i V_i w_i$, since $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$, we must have $(z_i, a_{-i}) \mathcal{P} (w_i, b_{-i})$. If neither $x_i V_i y_i$ nor $z_i V_i w_i$, the desired conclusion easily follows from the monotonicity of \succeq .

[RC2] Suppose that $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$, $(y_i, c_{-i}) \mathcal{P} (x_i, d_{-i})$. This implies that neither $x_i V_i y_i$ nor $y_i V_i x_i$. If $z_i V_i w_i$, since $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$, we must have $(z_i, a_{-i}) \mathcal{P} (w_i, b_{-i})$. Similarly if $w_i V_i z_i$, since $(y_i, c_{-i}) \mathcal{P} (x_i, d_{-i})$, we must have $(w_i, c_{-i}) \mathcal{P} (z_i, d_{-i})$. If neither $z_i V_i w_i$ nor $w_i V_i z_i$, the desired conclusion easily follows from the completeness of S_i and the monotonicity of \succeq .

Part 2. By hypothesis, there is a weak order T_i on X_i such that

$$x_i T_i y_i \Rightarrow \forall z_i \in X_i, [y_i S_i z_i \Rightarrow x_i S_i z_i] \text{ and } [z_i S_i x_i \Rightarrow z_i S_i y_i], \text{ and} \\ x_i T_i y_i \Rightarrow \forall z_i \in X_i, [y_i U_i z_i \Rightarrow x_i U_i z_i] \text{ and } [z_i U_i x_i \Rightarrow z_i U_i y_i].$$

Let us show that, if $z_i T_i x_i$, then $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$ implies $(z_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$. By hypothesis, we have $S[(z_i, a_{-i}), (y_i, b_{-i})] \supseteq S[(x_i, a_{-i}), (y_i, b_{-i})]$, $S[(y_i, b_{-i}), (z_i, a_{-i})] \subseteq S[(y_i, b_{-i}), (x_i, a_{-i})]$, $V[(y_i, b_{-i}), (z_i, a_{-i})] \subseteq V[(y_i, b_{-i}), (x_i, a_{-i})]$, $V[(z_i, a_{-i}), (y_i, b_{-i})] \supseteq V[(x_i, a_{-i}), (y_i, b_{-i})]$.

By hypothesis, $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$ means that $(x_i, a_{-i}) \mathcal{S} (y_i, b_{-i})$ and $\text{Not}[(y_i, b_{-i}) \mathcal{S} (x_i, a_{-i})]$. Hence, using the monotonicity of \supseteq , because $(x_i, a_{-i}) \mathcal{S} (y_i, b_{-i})$, we know that $(z_i, a_{-i}) \mathcal{S} (y_i, b_{-i})$. Suppose now that $(y_i, b_{-i}) \mathcal{S} (z_i, a_{-i})$. Using the monotonicity of \supseteq , we conclude that $(y_i, b_{-i}) \mathcal{S} (x_i, a_{-i})$, a contradiction. Hence, we have $(z_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$.

Similarly, it is easy to show that, if $y_i T_i w_i$, then $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$ implies $(x_i, a_{-i}) \mathcal{P} (w_i, b_{-i})$.

Using these two facts and the completeness of T_i , it is easy to check that $AC1$, $AC2$, and $AC3$ hold.

Part 3. Suppose that $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$, $(w_i, a_{-i}) \mathcal{P} (z_i, b_{-i})$, $(y_i, c_{-i}) \mathcal{P} (x_i, d_{-i})$, and $(z_i, e_{-i}) \mathcal{P} (w_i, f_{-i})$. This implies that it is impossible to have $x_i V_i y_i$, $y_i V_i x_i$, $z_i V_i w_i$ and $w_i V_i z_i$. The desired conclusion easily follows from the monotonicity of \supseteq . \square

The following lemma gives a simple interpretation of *Maj3*. It is also useful for checking whether *Maj3* holds.

Lemma 28

Let \mathcal{P} be a binary relation on $X = \prod_{i=1}^n X_i$ satisfying *RC1* and *RC2*. Then \mathcal{P} satisfies *Maj3* iff, for all $i \in N$, for all $x_i, y_i, z_i, w_i, r_i, s_i \in X_i$, $[(x_i, y_i) \succ_i^* (y_i, x_i) \succ_i^* (z_i, w_i)] \Rightarrow (r_i, s_i) \lesssim_i^* (z_i, w_i)$ and $\text{Not}[(z_i, e_{-i}) \mathcal{P} (w_i, f_{-i})]$, for all $e_{-i}, f_{-i} \in X_{-i}$.

PROOF

Part $[\Rightarrow]$. Suppose that, for some $x_i, y_i, z_i, w_i, r_i, s_i \in X_i$, we have $(x_i, y_i) \succ_i^* (y_i, x_i) \succ_i^* (z_i, w_i)$ and $(z_i, w_i) \succ_i^* (r_i, s_i)$. This implies $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$, $\text{Not}[(y_i, a_{-i}) \mathcal{P} (x_i, b_{-i})]$, $(y_i, c_{-i}) \mathcal{P} (x_i, d_{-i})$, $\text{Not}[(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})]$ and $(z_i, e_{-i}) \mathcal{P} (w_i, f_{-i})$, for some $a_{-i}, b_{-i}, c_{-i}, d_{-i}, e_{-i}, f_{-i} \in X_{-i}$.

Since this implies $\text{Not}[(z_i, w_i) \lesssim_i^* (y_i, x_i)]$, we have, using *RC2* and Lemma 9.2, that $(w_i, z_i) \lesssim_i^* (x_i, y_i)$ so that $(w_i, a_{-i}) \mathcal{P} (z_i, b_{-i})$. Using *Maj3_i*, $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$, $(w_i, a_{-i}) \mathcal{P} (z_i, b_{-i})$, $(y_i, c_{-i}) \mathcal{P} (x_i, d_{-i})$, and $(z_i, e_{-i}) \mathcal{P} (w_i, f_{-i})$ imply $(y_i, a_{-i}) \mathcal{P} (x_i, b_{-i})$ or $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$, a contradiction. Note that the contradiction is obtained as soon as $(z_i, e_{-i}) \mathcal{P} (w_i, f_{-i})$, for some $e_{-i}, f_{-i} \in X_{-i}$. This proves the second part of the assertion.

Part $[\Leftarrow]$. To prove $Maj3_i$, we assume that the first three premises in $Maj3_i$ are true and the two conclusions are false. $Maj3_i$ will result if we prove that the fourth premise must be false. Using $RC1_i$, the first premise of $Maj3_i$ and the negation of the first conclusion imply $(x_i, y_i) \succ_i^* (y_i, x_i)$. Using $RC1_i$ again, the third premise and the negation of the second conclusion imply $(y_i, x_i) \succ_i^* (z_i, w_i)$. Since by hypothesis $[(x_i, y_i) \succ_i^* (y_i, x_i) \succ_i^* (z_i, w_i)] \Rightarrow \text{Not}[(z_i, e_{-i}) \mathcal{P}(w_i, f_{-i})]$, for all $e_{-i}, f_{-i} \in X_{-i}$, $Maj3_i$ is established. \square

4.3 Additional properties of an AP-R-CDR-AT

The following two conditions are clearly weaker than $MM1$. They shape the way bonus effects can occur in an AP-R-CDR-AT. The first condition amounts to saying that the inverse of a bonus must be a veto. The second one implies that there is only one type of bonus.

Definition 29

We say that \mathcal{P} satisfies

$$MM4_i \text{ if } \left. \begin{array}{l} (x_i, a_{-i}) \mathcal{P}(y_i, b_{-i}) \\ \text{and} \\ (z_i, a_{-i}) \mathcal{P}(w_i, b_{-i}) \\ \text{and} \\ (z_i, c_{-i}) \mathcal{P}(w_i, d_{-i}) \\ \text{and} \\ (w_i, e_{-i}) \mathcal{P}(z_i, f_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (y_i, a_{-i}) \mathcal{P}(x_i, b_{-i}) \\ \text{or} \\ (x_i, c_{-i}) \mathcal{P}(y_i, d_{-i}) \\ \text{or} \\ (w_i, a_{-i}) \mathcal{P}(z_i, b_{-i}), \end{array} \right. \quad (12)$$

$$MM5_i \text{ if } \left. \begin{array}{l} (x_i, a_{-i}) \mathcal{P}(y_i, b_{-i}) \\ \text{and} \\ (z_i, c_{-i}) \mathcal{P}(w_i, d_{-i}) \\ \text{and} \\ (z_i, a_{-i}) \mathcal{P}(w_i, b_{-i}) \\ \text{and} \\ (r_i, e_{-i}) \mathcal{P}(s_i, f_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (y_i, a_{-i}) \mathcal{P}(x_i, b_{-i}) \\ \text{or} \\ (x_i, c_{-i}) \mathcal{P}(y_i, d_{-i}), \\ \text{or} \\ (x_i, e_{-i}) \mathcal{P}(y_i, f_{-i}), \\ \text{or} \\ (z_i, e_{-i}) \mathcal{P}(w_i, f_{-i}), \\ \text{or} \\ (w_i, a_{-i}) \mathcal{P}(z_i, b_{-i}). \end{array} \right. \quad (13)$$

We say that $MM4$ (resp. $MM5$) holds if $MM4_i$ (resp. $MM5_i$) holds for all $i \in N$.

$MM4_i$ without its third conclusion is called $Maj4_i$. $MM5_i$ without its fifth conclusion is called $Maj5_i$.

The above two conditions clearly weaken condition $MM1$. Let us note the following relations between our conditions.

Lemma 30

We have

1. $Maj4_i \Rightarrow MM4_i$,
2. $MM1_i \Rightarrow MM4_i$,
3. $MM4_i$ and $RC2_i \Rightarrow Maj4_i$,
4. $Maj5_i \Rightarrow MM5_i$,
5. $MM1_i \Rightarrow MM5_i$,
6. $MM5_i$ and $RC2_i \Rightarrow Maj5_i$.

PROOF

The first two points are obvious. Indeed, $MM4_i$ adds to $Maj4_i$ an additional possible conclusions. Similarly, $MM4_i$ adds to $MM1_i$ an additional premise. We prove the third point. Suppose that $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$, $(z_i, a_{-i}) \mathcal{P} (w_i, b_{-i})$, $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$, $(w_i, e_{-i}) \mathcal{P} (z_i, f_{-i})$. If $\text{Not}[(w_i, a_{-i}) \mathcal{P} (z_i, b_{-i})]$, then one of the two possible conclusions of $Maj4_i$ holds. If $(w_i, a_{-i}) \mathcal{P} (z_i, b_{-i})$, using $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$ and $RC2_i$, we have either $(y_i, a_{-i}) \mathcal{P} (x_i, b_{-i})$ or $(x_i, c_{-i}) \mathcal{P} (y_i, d_{-i})$, which are the two possible conclusions of $Maj4_i$.

The fourth and fifth points are obvious. Indeed, $MM5_i$ adds to $Maj5_i$ an additional premise and an additional possible conclusion. Similarly, $MM5_i$ adds to $MM1_i$ an additional premise and two possible conclusions. We prove the sixth point. Suppose that $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$, $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$, $(z_i, a_{-i}) \mathcal{P} (w_i, b_{-i})$, $(r_i, e_{-i}) \mathcal{P} (s_i, f_{-i})$.

If $\text{Not}[(w_i, a_{-i}) \mathcal{P} (z_i, b_{-i})]$, we must have one of the four possible conclusions of $Maj5_i$. If $(w_i, a_{-i}) \mathcal{P} (z_i, b_{-i})$, then $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$ and $RC2_i$ imply either $(y_i, a_{-i}) \mathcal{P} (x_i, b_{-i})$ or $(x_i, c_{-i}) \mathcal{P} (y_i, d_{-i})$. Hence, one of the possible conclusions of $Maj5_i$ holds. \square

As shown below, conditions $MM4$ and $MM5$ are satisfied by an AP-R-CDR-AT.

Lemma 31

If \mathcal{P} is an AP-R-CDR then

1. *it satisfies $Maj4$ and, hence, $MM4$,*
2. *it satisfies $Maj5$ and, hence, $MM5$.*

PROOF

Part 1. Suppose that $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$, $(z_i, a_{-i}) \mathcal{P} (w_i, b_{-i})$, $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$, and $(w_i, e_{-i}) \mathcal{P} (z_i, f_{-i})$. This implies that it is impossible to have: $y_i V_i x_i$, $w_i V_i z_i$, and $z_i V_i w_i$. If $x_i V_i y_i$, since $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$, we must have $(x_i, c_{-i}) \mathcal{P} (y_i, d_{-i})$. If $\text{Not}[x_i V_i y_i]$, the desired conclusion easily follows from the monotonicity of \succeq .

Part 2. Suppose that $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$, $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$, $(z_i, a_{-i}) \mathcal{P} (w_i, b_{-i})$, and $(r_i, e_{-i}) \mathcal{P} (s_i, f_{-i})$. This implies that it is impossible to have: $y_i V_i x_i$, $w_i V_i z_i$, and $s_i V_i r_i$. If $x_i V_i y_i$, since $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$, we must have $(x_i, c_{-i}) \mathcal{P} (y_i, d_{-i})$. Similarly, if $z_i V_i w_i$, since $(r_i, e_{-i}) \mathcal{P} (s_i, f_{-i})$, we must have $(z_i, e_{-i}) \mathcal{P} (w_i, f_{-i})$. If neither $x_i V_i y_i$ nor $z_i V_i w_i$, the desired conclusion easily follows from the monotonicity of \succeq . \square

It is clear that UC_i implies $M1_i$. Hence, UC_i implies $MM1_i$ and, therefore, $MM4_i$ and $MM5_i$. This will be useful for checking that $MM4_i$ and $MM5_i$ hold. The following two lemmas give an interpretation of $Maj4$ and $Maj5$ in terms of the relations \succ_i^* . They can also be useful to check that these conditions hold.

Lemma 32

Let \mathcal{P} be a binary relation on $X = \prod_{i=1}^n X_i$, which satisfies $RC1$. Then \mathcal{P} satisfies $Maj4$ if and only if, for all $i \in N$, for all $x_i, y_i, z_i, w_i, r_i, s_i \in X_i$, $[(z_i, w_i) \succ_i^* (x_i, y_i) \succ_i^* (y_i, x_i)] \Rightarrow (r_i, s_i) \succ_i^* (w_i, z_i)$ and $\text{Not}[(w_i, e_{-i}) \mathcal{P} (z_i, f_{-i})]$, for all $e_{-i}, f_{-i} \in X_{-i}$.

PROOF

Part $[\Rightarrow]$. Suppose that $(z_i, w_i) \succ_i^* (x_i, y_i) \succ_i^* (y_i, x_i)$. This implies $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$ and $\text{Not}[(x_i, c_{-i}) \mathcal{P} (y_i, d_{-i})]$, $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$ and $\text{Not}[(y_i, a_{-i}) \mathcal{P} (x_i, b_{-i})]$, for some $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$. Suppose furthermore that $(w_i, e_{-i}) \mathcal{P} (z_i, f_{-i})$, for some $e_{-i}, f_{-i} \in X_{-i}$. Because $RC1$ holds, we know that \succ_i^* is complete. Hence $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$ implies $(z_i, a_{-i}) \mathcal{P} (w_i, b_{-i})$. Applying now $Maj4_i$ leads to either $(x_i, c_{-i}) \mathcal{P} (y_i, d_{-i})$ or $(y_i, a_{-i}) \mathcal{P} (x_i, b_{-i})$, a contradiction.

Part $[\Leftarrow]$. To prove $Maj4_i$, we assume that the first three premises of $Maj4_i$ are true and the two conclusions are false. $Maj4_i$ will result if we prove that the fourth premise must be false. Using $RC1_i$, the first premise of $Maj4_i$ and the negation of the first conclusion imply $(x_i, y_i) \succ_i^* (y_i, x_i)$. Using $RC1_i$ again, the third premise and the negation of the second conclusion imply $(z_i, w_i) \succ_i^* (x_i, y_i)$. Since by hypothesis $[(z_i, w_i) \succ_i^* (x_i, y_i) \succ_i^* (y_i, x_i)] \Rightarrow \text{Not}[(w_i, e_{-i}) \mathcal{P} (z_i, f_{-i})]$, for all $e_{-i}, f_{-i} \in X_{-i}$, $Maj4_i$ is established. \square

Lemma 33

Let \mathcal{P} be a binary relation on $X = \prod_{i=1}^n X_i$ satisfying $RC1$. Then \mathcal{P} satisfies $Maj5$ iff, for all $i \in N$, for all $x_i, y_i, z_i, w_i, r_i, s_i \in X_i$, $[(z_i, w_i) \succ_i^* (x_i, y_i) \succ_i^* (y_i, x_i)] \Rightarrow (z_i, w_i) \succ_i^* (r_i, s_i)$, for all $r_i, s_i \in X_i$.

PROOF

Part $[\Rightarrow]$. Suppose that $(z_i, w_i) \succ_i^* (x_i, y_i) \succ_i^* (y_i, x_i)$. This implies $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$ and $\text{Not}[(x_i, c_{-i}) \mathcal{P} (y_i, d_{-i})]$, $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$ and $\text{Not}[(y_i, a_{-i}) \mathcal{P} (x_i, b_{-i})]$, for some $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$. Suppose furthermore that $(r_i, e_{-i}) \mathcal{P} (s_i, f_{-i})$, for some $r_i, s_i \in X_i$ and $e_{-i}, f_{-i} \in X_{-i}$. We must show that we have $(z_i, e_{-i}) \mathcal{P} (w_i, f_{-i})$.

Because $RC1$ is assumed, we have $(z_i, w_i) \succsim_i^* (x_i, y_i)$, hence, $(z_i, a_{-i}) \mathcal{P} (w_i, b_{-i})$. We have $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$, $(z_i, c_{-i}) \mathcal{P} (w_i, d_{-i})$, $(z_i, a_{-i}) \mathcal{P} (w_i, b_{-i})$, and $(r_i, e_{-i}) \mathcal{P} (s_i, f_{-i})$, while $\text{Not}[(y_i, a_{-i}) \mathcal{P} (x_i, b_{-i})]$ and $\text{Not}[(x_i, c_{-i}) \mathcal{P} (y_i, d_{-i})]$. Using $Maj5_i$, we obtain either $(x_i, e_{-i}) \mathcal{P} (y_i, f_{-i})$ or $(z_i, e_{-i}) \mathcal{P} (w_i, f_{-i})$. Since $(z_i, w_i) \succsim_i^* (x_i, y_i)$, the former consequence implies the latter, yielding the desired conclusion.

Part $[\Leftarrow]$. To prove $Maj5_i$, we assume that the four premises of $Maj5_i$ are true and the first three conclusions are false. $Maj5_i$ will result if we prove that the fourth conclusion must be true. Using $RC1_i$, the first premise of $Maj5_i$ and the negation of the first conclusion imply $(x_i, y_i) \succ_i^* (y_i, x_i)$. Using $RC1_i$ again, the second premise together with the negation of the second conclusion imply $(z_i, w_i) \succ_i^* (x_i, y_i)$. Since by hypothesis $[(z_i, w_i) \succ_i^* (x_i, y_i) \succ_i^* (y_i, x_i)] \Rightarrow [(z_i, w_i) \succ_i^* (r_i, s_i)]$ for all $r_i, s_i \in X_i$, the fourth conclusion in $Maj5_i$ is true, hence $Maj5_i$ is established. \square

Our last condition (the “bonus condition”) requires that the effect of one bonus is not altered by adding a second bonus on another attribute.

Definition 34

Let $i, j \in N$ with $i \neq j$. We say that \mathcal{P} satisfies

$$BC_{ij} \text{ if } \left. \begin{array}{l} (x_i, a_{-i}) \mathcal{P} (y_i, b_{-i}) \\ \text{and} \\ (x'_i, c_{-i}) \mathcal{P} (y'_i, d_{-i}) \\ \text{and} \\ (z_j, e_{-j}) \mathcal{P} (w_j, f_{-j}) \\ \text{and} \\ (z'_j, g_{-j}) \mathcal{P} (w'_j, h_{-j}) \\ \text{and} \\ (x_i, z_j, \alpha_{-ij}) \mathcal{P} (y_i, w_j, \beta_{-ij}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (x'_i, a_{-i}) \mathcal{P} (y'_i, b_{-i}) \\ \text{or} \\ (y'_i, c_{-i}) \mathcal{P} (x'_i, d_{-i}) \\ \text{or} \\ (z'_j, e_{-j}) \mathcal{P} (w'_j, f_{-j}) \\ \text{or} \\ (w'_j, g_{-j}) \mathcal{P} (z'_j, h_{-j}) \\ \text{or} \\ (x'_i, z_j, \alpha_{-ij}) \mathcal{P} (y'_i, w_j, \beta_{-ij}), \end{array} \right.$$

for all $x_i, x'_i, y_i, y'_i \in X_i$, all $z_j, z'_j, w_j, w'_j \in X_j$, all $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$, all $e_{-j}, f_{-j}, g_{-j}, h_{-j} \in X_{-j}$ and all $\alpha_{-ij}, \beta_{-ij} \in X_{-ij}$.

We say that BC holds if BC_{ij} holds for all $i, j \in N$ with $i \neq j$.

Condition BC ensures that the combination of bonuses has no different effect from that of a single bonus. In order to motivate this condition, suppose that

all the premises of the condition hold. Suppose furthermore that the first four conclusions are false. This implies that the differences (x_i, y_i) and (z_j, w_j) both act as bonuses, as we shall see in Lemma 38. The last conclusion therefore implies that, replacing a bonus on both $i, j \in N$ by a bonus on only one of these two attributes has no effect since at least one bonus is present.

The following lemma shows that BC holds for an AP-R-CDR.

Lemma 35

If \mathcal{P} is an AP-R-CDR then it satisfies BC .

PROOF

Let $\langle \succeq, S_i, V_i \rangle$ be a representation of \mathcal{P} . Suppose that all the premises of BC_{ij} hold and that the first four conclusions do not hold. Using the direct definition of an AP-R-CDR (Remark 23), it is readily seen that the above imply: $x'_i P_i y'_i$, $\text{Not}[x'_i V_i y'_i]$, $x_i P_i y_i$, $x_i V_i y_i$, $z'_j P_j w'_j$, $\text{Not}[z'_j V_j w'_j]$, $z_j P_j w_j$, $z_j V_j w_j$. Let $A = S[(x_i, z_j, \alpha_{-ij}), (y_i, w_j, \beta_{-ij})]$ and $B = S[(y_i, w_j, \beta_{-ij}), (x_i, z_j, \alpha_{-ij})]$. It is clear that $A = S[(x'_i, z_j, \alpha_{-ij}), (y'_i, w_j, \beta_{-ij})]$ and $B = S[(y'_i, w_j, \beta_{-ij}), (x'_i, z_j, \alpha_{-ij})]$. Because $(x_i, z_j, \alpha_{-ij}) \mathcal{P} (y_i, w_j, \beta_{-ij})$, we have $A \succeq B$. If $A \succ B$, we must have $(x'_i, z_j, \alpha_{-ij}) \mathcal{P} (y'_i, w_j, \beta_{-ij})$. Suppose that $A \trianglelefteq B$. Since $V[(x'_i, z_j, \alpha_{-ij}), (y'_i, w_j, \beta_{-ij})] \neq \emptyset$, we have $(x'_i, z_j, \alpha_{-ij}) \mathcal{P} (y'_i, w_j, \beta_{-ij})$. \square

The following three lemmas will help checking that BC holds.

Lemma 36

If \mathcal{P} satisfies UC_i then, for all $j \neq i$ it satisfies BC_{ij} .

PROOF

Suppose that $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$ and $(x'_i, c_{-i}) \mathcal{P} (y'_i, d_{-i})$. If the second conclusion of BC_{ij} is not verified, we have $\text{Not}[(y'_i, c_{-i}) \mathcal{P} (x'_i, d_{-i})]$. This implies $\text{Not}[(y'_i, x'_i) \succ_i^* (x'_i, y'_i)]$ and UC_i implies $(x'_i, y'_i) \succ_i^* (a_i, b_i)$, for all $a_i, b_i \in X_i$. Hence, we have $(x'_i, a_{-i}) \mathcal{P} (y'_i, b_{-i})$, so that the first conclusion of BC_{ij} holds. \square

Lemma 37

If \mathcal{P} satisfies $RC1_i$, $RC2_i$, and $MM1_i$ then, for all $j \neq i$, it satisfies BC_{ij} .

PROOF

The proof follows from combining Lemma 36 with Lemma 17. \square

Lemma 38

Let $i, j \in N$ with $i \neq j$. Let \mathcal{P} be a relation satisfying $RC1_i$ and $RC1_j$. Then \mathcal{P} satisfies BC_{ij} if and only if, for all $x_i, x'_i, y_i, y'_i \in X_i$, and, for all $z_j, z'_j, w_j, w'_j \in X_j$, we have that

$$\left. \begin{array}{l} (x_i, y_i) \succ_i^* (x'_i, y'_i) \succ_i^* (y'_i, x'_i) \\ (z_j, w_j) \succ_j^* (z'_j, w'_j) \succ_j^* (w'_j, z'_j) \\ (x_i, z_j, \alpha_{-ij}) \mathcal{P} (y_i, w_j, \beta_{-ij}) \end{array} \right\} \Rightarrow (x'_i, z_j, \alpha_{-ij}) \mathcal{P} (y'_i, w_j, \beta_{-ij}). \quad (14)$$

PROOF

Conditionally on the fact that \succsim_i^* and \succsim_j^* are complete relations (a consequence of $RC1_i$ and $RC1_j$), the conjunction of the five premises of BC_{ij} and the negation of the first four consequences is logically equivalent with the premises of (14). Applying BC_{ij} yields the consequence of (14). Conversely, using (14) yields the fifth consequence of BC_{ij} . \square

An easy consequence of $RC1$ and BC is the following property saying that a bonus on attribute i has the same effect as a bonus on attribute j .

Lemma 39

Let \mathcal{P} be a relation satisfying $RC1$ and BC . Then, for all $i, j \in N$ with $i \neq j$, we have:

$$\left. \begin{array}{l} (x_i, a_{-i}) \mathcal{P} (y_i, b_{-i}) \\ \text{and} \\ (x'_i, c_{-i}) \mathcal{P} (y'_i, d_{-i}) \\ \text{and} \\ (z_j, e_{-j}) \mathcal{P} (w_j, f_{-j}) \\ \text{and} \\ (z'_j, g_{-j}) \mathcal{P} (w'_j, h_{-j}) \\ \text{and} \\ (x_i, z'_j, \alpha_{-ij}) \mathcal{P} (y_i, w'_j, \beta_{-ij}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (x'_i, a_{-i}) \mathcal{P} (y'_i, b_{-i}) \\ \text{or} \\ (y'_i, c_{-i}) \mathcal{P} (x'_i, d_{-i}) \\ \text{or} \\ (z'_j, e_{-j}) \mathcal{P} (w'_j, f_{-j}) \\ \text{or} \\ (w'_j, g_{-j}) \mathcal{P} (z'_j, h_{-j}) \\ \text{or} \\ (x'_i, z_j, \alpha_{-ij}) \mathcal{P} (y'_i, w_j, \beta_{-ij}), \end{array} \right. \quad (15)$$

for all $x_i, x'_i, y_i, y'_i \in X_i$, all $z_j, z'_j, w_j, w'_j \in X_j$, all $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$, all $e_{-j}, f_{-j}, g_{-j}, h_{-j} \in X_{-j}$, and all $\alpha_{-ij}, \beta_{-ij} \in X_{-ij}$.

Assuming $RC1$, property (15) is equivalent with

$$\left. \begin{array}{l} (x_i, y_i) \succ_i^* (x'_i, y'_i) \succ_i^* (y'_i, x'_i) \\ (z_j, w_j) \succ_j^* (z'_j, w'_j) \succ_j^* (w'_j, z'_j) \\ (x_i, z'_j, \alpha_{-ij}) \mathcal{P} (y_i, w'_j, \beta_{-ij}) \end{array} \right\} \Rightarrow (x'_i, z_j, \alpha_{-ij}) \mathcal{P} (y'_i, w_j, \beta_{-ij}). \quad (16)$$

PROOF

Assume that the five premises of (15) are satisfied and none of the first four conclusions are. Using $RC1_j$, we infer that $(z_j, w_j) \succ_j^* (z'_j, w'_j)$. Hence, from the fifth premise in (15), we deduce that $(x_i, z_j, \alpha_{-ij}) \mathcal{P} (y_i, w_j, \beta_{-ij})$, which is the fifth premise of BC . Applying BC we get $(x'_i, z_j, \alpha_{-ij}) \mathcal{P} (y'_i, w_j, \beta_{-ij})$, which establishes property (15).

The proof that (15) is equivalent with (16) is similar to that of Lemma 38 (assuming that \mathcal{P} satisfies $RC1$). \square

Condition (15) can be interpreted in the following way. Suppose that all the premises of the condition hold. Suppose furthermore that the first four conclusions

are false. This is easily seen to imply that the differences (x_i, y_i) and (z_j, w_j) both act as bonuses. The last conclusion therefore implies that, replacing a bonus on $i \in N$ by a relation P_i (that may or not be a bonus) and replacing a relation P_j on $j \in N$ (that may or not be a bonus) by a bonus on j has no effect since at least one bonus is present. Property (15) ensures that bonuses can be exchanged between attributes. A bonus on $i \in N$ has a similar effect to that of a bonus on $j \in N$.

4.4 Vetoes and bonuses

We define on each $i \in N$ a relation S_i letting, for all $x_i, y_i \in X_i$,

$$x_i S_i y_i \Leftrightarrow (x_i, y_i) \succsim_i^* (y_i, x_i). \quad (17)$$

The asymmetric (resp. symmetric) part of S_i will be denoted by P_i (resp. I_i). This is standard definition of S_i already used for the study of CDR-AT in Bouyssou and Pirlot (2009). We have:

Lemma 40

Let \mathcal{P} be a binary relation on $X = \prod_{i=1}^n X_i$ satisfying RC1_i. The relation S_i on X_i defined by (17) is complete. Moreover, if RC2_i holds and $x_i I_i y_i$ and $z_i I_i w_i$, then $(x_i, y_i) \sim_i^ (z_i, w_i) \sim_i^* (y_i, x_i) \sim_i^* (w_i, z_i) \sim_i^* (a_i, a_i)$, for all $a_i \in X_i$. The relation P_i is nonempty.*

PROOF

RC1 implies that \succsim_i^* is complete. Hence S_i is complete.

Using the definition of S_i , $x_i I_i y_i$ and $z_i I_i w_i$ is equivalent to $(x_i, y_i) \sim_i^* (y_i, x_i)$ and $(z_i, w_i) \sim_i^* (w_i, z_i)$. The conclusion follows from Lemma 9.2.

If P_i is empty, we must have for all $x_i, y_i, z_i, w_i \in X_i$, $x_i I_i y_i$ and $z_i I_i w_i$. This implies $(x_i, y_i) \sim_i^* (z_i, w_i)$, violating the fact that attribute $i \in N$ is influential. \square

The definition of the relation V_i requires more care. Indeed, it includes at the same time cases of vetoes and cases of bonuses. The relation \mathcal{V}_i will model vetoes. The relation \mathcal{B}_i will model bonuses. The necessity to define separately the relations \mathcal{V}_i and \mathcal{B}_i comes from the fact that one can build degenerate examples in which \mathcal{B}_i is nonempty while \mathcal{V}_i is empty (in that case all negative differences act as a veto). Similarly there are cases in which \mathcal{B}_i is empty but \mathcal{V}_i is not (e.g., when the relation \triangleq only contains $N \triangleq N$.)

Define the relation \mathcal{V}_i on X_i letting, for all $x_i, y_i \in X_i$

$$x_i \mathcal{V}_i y_i \Leftrightarrow (z_i, w_i) \succ_i^* (w_i, z_i) \succ_i^* (y_i, x_i) \text{ for some } z_i, w_i \in X_i. \quad (18)$$

We have:

Lemma 41

Let \mathcal{P} be a binary relation on $X = \prod_{i=1}^n X_i$ satisfying *RC1*, *RC2*, and *Maj3*. We have:

1. $\mathcal{V}_i \subseteq P_i$,
2. $[x_i \mathcal{V}_i y_i \text{ and } z_i \mathcal{V}_i w_i] \Rightarrow (y_i, x_i) \sim_i^* (w_i, z_i)$,
3. $[x_i P_i y_i, z_i P_i w_i, \text{Not}[x_i \mathcal{V}_i y_i] \text{ and } \text{Not}[z_i \mathcal{V}_i w_i]] \Rightarrow (y_i, x_i) \sim_i^* (w_i, z_i)$.

PROOF

Part 1. Suppose that $x_i \mathcal{V}_i y_i$, so that $(z_i, w_i) \succ_i^* (w_i, z_i) \succ_i^* (y_i, x_i)$, for some $z_i, w_i \in X_i$. Suppose that $\text{Not}[x_i P_i y_i]$ so that $(y_i, x_i) \lesssim_i^* (x_i, y_i)$. Using *RC1* and *RC2*, it is easy to check that $(z_i, w_i) \succ_i^* (w_i, z_i)$ implies $(z_i, w_i) \lesssim_i^* (a_i, a_i) \lesssim_i^* (w_i, z_i)$, for all $a_i \in X_i$. We therefore obtain $(a_i, a_i) \succ_i^* (y_i, x_i) \lesssim_i^* (x_i, y_i)$. This contradicts *RC2*, using Lemma 9.2.

Part 2. Suppose that $x_i \mathcal{V}_i y_i$ and $z_i \mathcal{V}_i w_i$. In view of (18), Lemma 28 implies that $(w_i, z_i) \lesssim_i^* (y_i, x_i)$ and $(y_i, x_i) \lesssim_i^* (w_i, z_i)$ so that $(w_i, z_i) \sim_i^* (y_i, x_i)$.

Part 3. By definition, we have $(x_i, y_i) \succ_i^* (y_i, x_i)$ and $(z_i, w_i) \succ_i^* (w_i, z_i)$. Suppose that $(y_i, x_i) \succ_i^* (w_i, z_i)$. This would imply $(x_i, y_i) \succ_i^* (y_i, x_i) \succ_i^* (w_i, z_i)$, contradicting the fact that $\text{Not}[z_i \mathcal{V}_i w_i]$. Similarly it is impossible that $(w_i, z_i) \succ_i^* (y_i, x_i)$. Hence, we have $(y_i, x_i) \sim_i^* (w_i, z_i)$. \square

Remark 42

Assuming *RC1*, *RC2* and *Maj3*, observe that, if \mathcal{V}_i is nonempty, then there exists $z_i, w_i \in X_i$ such that $z_i P_i w_i$ and $\text{Not}[z_i \mathcal{V}_i w_i]$.

Indeed, since \mathcal{V}_i is nonempty, we know that there are $x_i, y_i \in X_i$ such that $(z_i, w_i) \succ_i^* (w_i, z_i) \succ_i^* (y_i, x_i)$. Hence, we know that $z_i P_i w_i$. Suppose now that $z_i \mathcal{V}_i w_i$, so that there are $r_i, s_i \in X_i$ such that $(r_i, s_i) \succ_i^* (s_i, r_i) \succ_i^* (w_i, z_i)$. This implies $(w_i, z_i) \succ_i^* (y_i, x_i)$, violating Lemma 28. \bullet

Define the relation \mathcal{B}_i on X_i letting, for all $x_i, y_i \in X_i$

$$x_i \mathcal{B}_i y_i \Leftrightarrow (x_i, y_i) \succ_i^* (z_i, w_i) \succ_i^* (w_i, z_i) \text{ for some } z_i, w_i \in X_i. \quad (19)$$

Lemma 43

Let \mathcal{P} be a binary relation on $X = \prod_{i=1}^n X_i$ satisfying *RC1*, *RC2*, *Maj3*, *Maj4*, and *Maj5*. We have:

1. $\mathcal{B}_i \subseteq P_i$.
2. $z_i \mathcal{B}_i w_i$ and $x_i \mathcal{B}_i y_i$ imply $(z_i, w_i) \sim_i^* (x_i, y_i)$.
3. $z_i \mathcal{B}_i w_i$ and $x_i \mathcal{B}_i y_i$ imply $(w_i, z_i) \sim_i^* (y_i, x_i)$.

4. $z_i P_i w_i$, $\text{Not}[z_i \mathcal{B}_i w_i]$, $x_i P_i y_i$ and $\text{Not}[x_i \mathcal{B}_i y_i]$, imply $(z_i, w_i) \sim_i^* (x_i, y_i)$.

5. $z_i \mathcal{B}_i w_i$, $x_i P_i y_i$, $\text{Not}[x_i \mathcal{B}_i y_i]$ imply $(z_i, w_i) \succ_i^* (x_i, y_i)$.

PROOF

Part 1. Suppose that $x_i \mathcal{B}_i y_i$, so that $(x_i, y_i) \succ_i^* (z_i, w_i) \succ_i^* (w_i, z_i)$, for some $z_i, w_i \in X_i$. Suppose that $\text{Not}[x_i P_i y_i]$ so that $(y_i, x_i) \succsim_i^* (x_i, y_i)$. Using *RC1* and *RC2*, it is easy to check that $(z_i, w_i) \succ_i^* (w_i, z_i)$ implies $(z_i, w_i) \succsim_i^* (a_i, a_i) \succsim_i^* (w_i, z_i)$, for all $a_i \in X_i$. We therefore obtain $(y_i, x_i) \succsim_i^* (x_i, y_i) \succ_i^* (a_i, a_i)$. This contradicts *RC2* (Lemma 9.2).

Part 2. Suppose that $z_i \mathcal{B}_i w_i$ and $x_i \mathcal{B}_i y_i$. In view of (19), Lemma 33 implies $(z_i, w_i) \succsim_i^* (r_i, s_i)$ and $(x_i, y_i) \succsim_i^* (r_i, s_i)$, for all $r_i, s_i \in X_i$. This implies $(z_i, w_i) \sim_i^* (x_i, y_i)$.

Part 3. Suppose that $z_i \mathcal{B}_i w_i$ and $x_i \mathcal{B}_i y_i$. Lemma 32 implies $(r_i, s_i) \succsim_i^* (w_i, z_i)$ and $(r_i, s_i) \succsim_i^* (y_i, x_i)$, for all $r_i, s_i \in X_i$. This implies $(w_i, z_i) \sim_i^* (y_i, x_i)$.

Part 4. Suppose that $z_i P_i w_i$, $\text{Not}[z_i \mathcal{B}_i w_i]$, $x_i P_i y_i$ and $\text{Not}[x_i \mathcal{B}_i y_i]$. This implies that $(z_i, w_i) \succ_i^* (w_i, z_i)$ and $(x_i, y_i) \succ_i^* (y_i, x_i)$. If $(z_i, w_i) \succ_i^* (x_i, y_i)$, we obtain $z_i \mathcal{B}_i w_i$, a contradiction. Similarly, it is impossible that $(x_i, y_i) \succ_i^* (z_i, w_i)$. Hence, we must have $(x_i, y_i) \sim_i^* (z_i, w_i)$.

Part 5. Suppose that $z_i \mathcal{B}_i w_i$, $x_i P_i y_i$, $\text{Not}[x_i \mathcal{B}_i y_i]$. This implies $(z_i, w_i) \succ_i^* (a_i, b_i) \succ_i^* (b_i, a_i)$, for some $a_i, b_i \in X_i$, and $(x_i, y_i) \succ_i^* (y_i, x_i)$. If $(x_i, y_i) \succsim_i^* (z_i, w_i)$, we obtain $(x_i, y_i) \succ_i^* (a_i, b_i) \succ_i^* (b_i, a_i)$, implying that $x_i \mathcal{B}_i y_i$, a contradiction. \square

Remark 44

Observe that if \mathcal{B}_i is nonempty then there exist $z_i, w_i \in X_i$ such that $z_i P_i w_i$ and $\text{Not}[z_i \mathcal{B}_i w_i]$.

Assuming that \mathcal{B}_i is nonempty, we know that there are $x_i, y_i \in X_i$ such that $(x_i, y_i) \succ_i^* (z_i, w_i) \succ_i^* (w_i, z_i)$. Hence, we know that $z_i P_i w_i$. It is clearly impossible that $z_i \mathcal{B}_i w_i$, since by Lemma 33, this would imply that $(z_i, w_i) \succ_i^* (r_i, s_i)$, for all $r_i, s_i \in X_i$, contrary to the fact that $(x_i, y_i) \succ_i^* (z_i, w_i)$. \bullet

Let us now define the relation V_i on X_i , letting, for all $x_i, y_i \in X_i$,

$$x_i V_i y_i \Leftrightarrow [x_i \mathcal{B}_i y_i \text{ or } x_i \mathcal{V}_i y_i]. \quad (20)$$

Combining Lemma 43.1 with Lemma 41.1 shows that $V_i \subseteq P_i$.

4.5 The main result

The main result of this note is the following.

Theorem 45 (Characterization of AP-R-CDR)

Let \mathcal{P} be a binary relation on $X = \prod_{i=1}^n X_i$ and assume that all attributes are influential for \mathcal{P} . Then \mathcal{P} is an AP-R-CDR-AT iff \mathcal{P} is an asymmetric relation satisfying *RC1*, *RC2*, *MM3*, *MM4*, *MM5*, *BC*, *AC1*, *AC2*, and *AC3*.

PROOF

Necessity was shown above. We show sufficiency.

Define the relation S_i and V_i on X_i as above (formulas (17), (18), (19), (20)). We have shown above that S_i is complete, that P_i is nonempty and that V_i is included in P_i .

Consider any two subsets $A, B \subseteq N$ such that $A \cup B = N$ and let:

$$A \triangleright B \Leftrightarrow$$

$$[x \mathcal{P} y, \text{ for all } x, y \in X \text{ such that } S[x, y] = A, S[y, x] = B, \text{ and } V[y, x] = \emptyset].$$

Similarly, Consider any two subsets $A, B \subseteq N$ such that $A \cup B = N$ and let:

$$A \trianglelefteq B \Leftrightarrow$$

$$[x \mathcal{P} y, \text{Not}[z \mathcal{P} w], \text{ for some } x, y, z, w \in X \text{ such that}$$

$$S[x, y] = S[z, w] = A, S[y, x] = S[w, z] = B, V[y, x] = V[w, z] = \emptyset].$$

Let us show that, with the previous definitions of S_i , V_i and \triangleright , (10–11) holds for \mathcal{P} .

Part $[\Rightarrow]$ of (10–11). Suppose that $x \mathcal{P} y$. Using Lemma 28, we know that it is impossible that $y_i \mathcal{V}_i x_i$, for some $i \in N$. Similarly, using Lemma 32, it is impossible that $y_i \mathcal{B}_i x_i$, for some $i \in N$. In view of (20), this implies $V(y, x) = \emptyset$. Let $A = S[x, y]$ and $B = S[y, x]$. We must show that $A \triangleright B$ or that $[A \trianglelefteq B$ and $V[x, y] \neq \emptyset]$. We distinguish two cases.

1. Suppose first that, for all $z, w \in X$, $A = S[z, w]$, $B = S[w, z]$ and $V[w, z] = \emptyset$ implies $z \mathcal{P} w$. By definition, this implies $A \triangleright B$. Hence, in this case, we have $S[x, y] \triangleright S[y, x]$ and $V[y, x] = \emptyset$.
2. Suppose now that, for some $z, w \in X$, $A = S[z, w]$, $B = S[w, z]$ and $V[w, z] = \emptyset$ and $\text{Not}[z \mathcal{P} w]$. Since $x \mathcal{P} y$, this implies $A \trianglelefteq B$. Let us show that we must have $x_i \mathcal{B}_i y_i$, for some $i \in N$.

We have $A = S[x, y] = S[z, w]$, $B = S[y, x] = S[w, z]$ and $V[y, x] = V[w, z] = \emptyset$. We have $x_i I_i y_i \Leftrightarrow z_i I_i w_i$. For all $i \in N$ such that $x_i I_i y_i$, we have, using Lemma 40, $(x_i, y_i) \sim_i^{**} (z_i, w_i)$. We have $y_i P_i x_i \Leftrightarrow w_i P_i z_i$. Because we know that $V[y, x] = V[w, z] = \emptyset$, Lemma 40 implies that, for all $i \in N$ such that $y_i P_i x_i$, we have $(x_i, y_i) \sim_i^* (z_i, w_i)$. We have $x_i P_i y_i \Leftrightarrow z_i P_i w_i$. Suppose that there is no $j \in N$ such that $x_j \mathcal{B}_j y_j$. Using Parts 4 and 5 of Lemma 43, this implies that $(z_i, w_i) \succsim_i^* (x_i, y_i)$, for all $i \in N$ such that $x_i P_i y_i$.

We have $(z_i, w_i) \succsim_i^* (x_i, y_i)$, for all $i \in N$. This is contradictory since $x \mathcal{P} y$ and $\text{Not}[z \mathcal{P} w]$ (see Lemma 10). Hence, we must have $x_j \mathcal{B}_j y_j$, for some $j \in N$. We therefore have $S[x, y] \trianglelefteq S[y, x]$, $V[y, x] = \emptyset$ and $V[x, y] \neq \emptyset$.

Part $[\Leftarrow]$ of (10–11). Let $A, B \subseteq N$ such that $A \cup B = N$. Because P_i is nonempty (since all attributes are influential for \mathcal{P}), we know that there are $x, y \in X$ such that $A = S[x, y]$ and $B = S[y, x]$.

If $A \triangleright B$ and $V[y, x] = \emptyset$, we have, by construction, $x \mathcal{P} y$.

Suppose now that $A \trianglelefteq B$, $V[y, x] = \emptyset$ and $V[x, y] \neq \emptyset$ and $\text{Not}[x \mathcal{P} y]$. By construction, we know that there are $z, w \in X$ such that $z \mathcal{P} w$, $V[w, z] = \emptyset$, $A = S[z, w]$ and $B = S[w, z]$.

If $V[x, y] \supseteq V[z, w]$, we know that $(x_i, y_i) \succsim_i^* (z_i, w_i)$, for all $i \in N$. Since $\text{Not}[x \mathcal{P} y]$ and $z \mathcal{P} w$, this is contradictory (see Lemma 10).

Otherwise, we repeatedly use BC to build alternatives z' and w' such that $A = S[z', w']$, $B = S[w', z']$, $z' \mathcal{P} w'$ and $V[w', z']$ contains a single attribute. Suppose for definiteness that $\{k\} = V[w', z']$.

If $k \in V[x, y]$, we have $(x_i, y_i) \succsim_i^* (z'_i, w'_i)$, for all $i \in N$, contradicting the fact that $\text{Not}[x \mathcal{P} y]$ and $z \mathcal{P} w$. If $k \notin V[x, y]$, we use property (15), defined in Lemma 39, to build alternatives z'' and w'' such that $A = S[z', w']$, $B = S[w', z']$, $z'' \mathcal{P} w''$ and $\{\ell\} = V[w'', z''] \subseteq V[x, y]$. Hence, we have $(x_i, y_i) \succsim_i^* (z''_i, w''_i)$, for all $i \in N$, contradicting the fact that $\text{Not}[x \mathcal{P} y]$ and $z \mathcal{P} w$.

This shows that, with our definitions, (10–11) hold.

Let us now show that the relation $\triangleright = \triangleright \cup \trianglelefteq$ defined on the set of subsets of N having N for union is monotonic, i.e., satisfies (2).

Suppose that $A \triangleright B$. It is easy to see that the proof used in Bouyssou and Pirlot (2012) can be used to show that \triangleright is monotonic. Indeed, the set of all $x, y \in X$ such that $V[y, x] = \emptyset$, $A = S[x, y]$ and $B = S[y, x]$ must contain alternatives $z, w \in X$ such that $V[w, z] = \emptyset$, $A = S[z, w]$, $B = S[w, z]$, and $V[z, w] = \emptyset$.

Suppose that $A \trianglelefteq B$. By construction, we know that there are $x, y \in X$ such that $A = S[x, y]$, $B = S[y, x]$, and $x \mathcal{P} y$. Suppose that $C \supseteq A$, $B \supseteq D$ such that $C \cup D = N$. Let us show that there are $z, w \in X$ such that $C = S[z, w]$, $D = S[w, z]$, and $z \mathcal{P} w$, which will complete the proof.

Let $E = C \setminus A$. We have $B \supseteq E$. We build $z, w \in X$ with $S[z, w] = C \supseteq S[w, z] = B$. We know that for all $a_i \in X_i$, we have $a_i I_i a_i$. Using such pairs, define $z', w' \in X$ as follows:

	A	E	$B \setminus E$
z'	x_i	a_i	x_i
w'	y_i	a_i	y_i

It is clear that $(z'_i, w'_i) \succsim_i^* (x_i, y_i)$, for all $i \in N$. Hence, we have $z' \mathcal{P} w'$.

Since $C \cup D = N$, we have $B \setminus D \subseteq C$. Let $F = B \setminus D$. For all $i \in N$, P_i is not empty so that we can take, for all $i \in F$, any $a_i, b_i \in X_i$ such that $a_i P_i b_i$. Using such pairs, define $z'', w'' \in X$ as follows:

$C \setminus F$	F	D
z''	z'_i	a_i
w''	w'_i	b_i
		z_i
		w_i

It is clear that $(z''_i, w''_i) \succsim_i^* (z_i, w_i)$, for all $i \in N$. Hence, we have $z'' \mathcal{P} w''$.

Let us finally show that (S_i, U_i) form a homogeneous chain of semiorders. We already know that both S_i and U_i are complete and that V_i is included in P_i . In view of Remark 3, all we have to show is that there is a weak order T_i on X_i such that:

$$\begin{aligned} x_i T_i y_i &\Rightarrow \forall z_i \in X_i, [y_i S_i z_i \Rightarrow x_i S_i z_i] \text{ and } [z_i S_i x_i \Rightarrow z_i S_i y_i], \text{ and} \\ x_i T_i y_i &\Rightarrow \forall z_i \in X_i, [y_i V_i z_i \Rightarrow x_i V_i z_i] \text{ and } [z_i V_i x_i \Rightarrow z_i V_i y_i]. \end{aligned}$$

Let us show that the weak order T_i can be taken to be \succsim_i^\pm , this relation being a weak order due to *AC1*, *AC2*, and *AC3*.

Suppose that $a_i S_i b_i$ so that $(a_i, b_i) \succsim_i^* (b_i, a_i)$. Suppose furthermore that $x_i \succsim_i^\pm a_i$. By construction $x_i \succsim_i^\pm a_i$ implies $x_i \succsim_i^+ a_i$, which is equivalent to saying that $(x_i, c_i) \succsim_i^* (a_i, c_i)$, for all $c_i \in X_i$. Taking $c_i = b_i$, we obtain $(x_i, b_i) \succsim_i^* (a_i, b_i)$. Similarly, $x_i \succsim_i^\pm a_i$ implies $x_i \succsim_i^- a_i$, which is equivalent to saying that $(c_i, a_i) \succsim_i^* (c_i, x_i)$, for all $c_i \in X_i$. Taking $c_i = b_i$, we obtain $(b_i, a_i) \succsim_i^* (b_i, x_i)$. Hence, we have $(x_i, b_i) \succsim_i^* (a_i, b_i) \succsim_i^* (b_i, a_i) \succsim_i^* (b_i, x_i)$, so that $(x_i, b_i) \succsim_i^* (b_i, x_i)$ and $x_i S_i b_i$. The proof that $b_i S_i a_i$ and $a_i \succsim_i^\pm x_i$ imply $b_i S_i x_i$ is similar.

Suppose now that $a_i V_i b_i$, so that either $a_i \mathcal{V}_i b_i$ or $a_i \mathcal{B}_i b_i$. Suppose that $a_i \mathcal{V}_i b_i$, so that $(c_i, d_i) \succ_i^* (d_i, c_i) \succ_i^* (b_i, a_i)$, for some $c_i, d_i \in X_i$. Suppose furthermore that $x_i \succsim_i^\pm a_i$. By construction, $x_i \succsim_i^\pm a_i$ implies $x_i \succsim_i^- a_i$, which is equivalent to saying that $(e_i, a_i) \succsim_i^* (e_i, x_i)$, for all $e_i \in X_i$. Taking $e_i = b_i$, we obtain $(b_i, a_i) \succsim_i^* (b_i, x_i)$. Hence, we have $(c_i, d_i) \succ_i^* (d_i, c_i) \succ_i^* (b_i, a_i) \succsim_i^* (b_i, x_i)$. This implies $x_i \mathcal{V}_i b_i$. The proof that $b_i \mathcal{V}_i a_i$ and $a_i \succsim_i^\pm x_i$ imply $b_i \mathcal{V}_i x_i$ is similar.

Suppose finally that $a_i \mathcal{B}_i b_i$, so that $(a_i, b_i) \succ_i^* (c_i, d_i) \succ_i^* (d_i, c_i)$, for some $c_i, d_i \in X_i$. Suppose furthermore that $x_i \succsim_i^\pm a_i$. By construction $x_i \succsim_i^\pm a_i$ implies $x_i \succsim_i^+ a_i$, which is equivalent to saying that $(x_i, c_i) \succsim_i^* (a_i, c_i)$, for all $c_i \in X_i$. Taking $c_i = b_i$, we obtain $(x_i, b_i) \succsim_i^* (a_i, b_i)$. Hence, we have $(x_i, b_i) \succsim_i^* (a_i, b_i) \succ_i^* (c_i, d_i) \succ_i^* (d_i, c_i)$, so that $x_i \mathcal{B}_i b_i$. The proof that $b_i \mathcal{B}_i a_i$ and $a_i \succsim_i^\pm x_i$ imply $b_i \mathcal{B}_i x_i$ is similar. This completes the proof. \square

Remark 46

An easy corollary of the above result is that \mathcal{P} is an AP-R-CDR iff it is asymmetric and satisfies *RC1*, *RC2*, *Maj3*, *Maj4*, *Maj5*, and *BC*. Indeed, in the above proof, conditions *AC1*, *AC2*, and *AC3* (via the fact that they imply that \succsim_i^\pm is a weak order) are only used to show that (S_i, U_i) is a homogeneous chain of semiorders. \bullet

The above result uses 9 conditions. It would be of minor interest if these 9 conditions were not independent. We show below that this is indeed the case in the class of asymmetric relations, i.e., the class of relations of interest for the study of AP-R-CDR-AT.

Proposition 47

In the class of asymmetric relations, conditions $RC1$, $RC2$, $AC1$, $AC2$, $AC3$, $MM3$, $MM4$, $MM5$, and BC are independent.

PROOF

We need 9 examples.

Bouyssou and Pirlot (2013) have shown that conditions $RC1$, $RC2$, $AC1$, $AC2$, $AC3$, $MM1$ and $MM3$ are independent in the class of asymmetric relations.

We know from Lemma 30 that $MM1$ implies $MM4$ and $MM5$. Moreover, we know from Lemma 37 that $RC1$, $RC2$, and $MM1$ imply BC . Hence, the four examples in Bouyssou and Pirlot (2013) concerning $MM3$, $AC1$, $AC2$, $AC3$ can be used here. This leaves us with 5 examples. We give them below in Section 4.6. \square

4.6 Examples

We give below the 5 examples needed to complete the proof of Proposition 47.

Example 48 (*Not*[BC])

Let $X = \prod_{i=1}^5 X_i$ with $X_i = \{x_i, y_i, z_i\}$, for $i = 1, 2$ and $X_j = \{x_j, y_j\}$ for $j = 3, 4, 5$. We define \mathcal{P} as follows. For all $a, b \in X$, we have $a \mathcal{P} b$ if $\sum_{i=1}^5 p_i(a_i, b_i) > 0$, where p_i is defined in the following table for all pairs (a_i, b_i) in $X_i \times X_i$.

p_i	X_1	X_2	X_j for $j = 3, 4, 5$
21		(x_2, z_2)	
11	(x_1, z_1)		
10	$(x_1, y_1), (y_1, z_1)$	$(x_2, y_2), (y_2, z_2)$	(x_j, y_j)
0	(a_1, a_1)	(a_2, a_2)	(a_j, a_j)
-10	$(y_1, x_1), (z_1, y_1)$	$(y_2, x_2), (z_2, y_2)$	(y_j, x_j)
-100	(z_1, x_1)	(z_2, x_2)	

It is easy to check that relation \mathcal{P} is asymmetric, mainly because the opposite (z_1, x_1) and (z_2, x_2) of the bonus pairs (x_1, z_1) and (x_2, z_2) play the rôle of vetoes. Indeed, due to the fact that $p_1(z_1, x_1) = p_2(z_2, x_2) = -100$, these pairs are incompatible with relation \mathcal{P} .

For $i = 1, 2, 3, 4, 5$, we have $\succsim_i^* = \succsim_i^{**}$, which are the weak orders induced by the values assigned to p_i in the above table. Hence \mathcal{P} satisfies $RC1$ and $RC2$. It is easy to check that $AC1$, $AC2$, and $AC3$ hold.

The only cases of super negative pairs, i.e., pairs (c_i, d_i) such that there are elements a_i, b_i with $(a_i, b_i) \succ_i^* (b_i, a_i) \succ_i^* (c_i, d_i)$, are again (z_1, x_1) and (z_2, x_2) . These pairs, as already observed, play the rôle of vetoes. They are therefore incompatible with relation \mathcal{P} , which, in view of Lemma 28, entails that \mathcal{P} satisfies *Maj3*.

Since (z_1, x_1) and (z_2, x_2) are the opposite of the only two bonus pairs and are never involved in preferences, using Lemma 32 shows that \mathcal{P} satisfies *Maj4*. Using Lemma 33, it is also clear that *Maj5* holds. Hence, *MM4* and *MM5* hold.

In view of Lemma 38, property BC_{ij} is trivially satisfied for all $\{i, j\} \neq \{1, 2\}$. Using the same lemma, we see that this example violates BC_{12} since we have $(x_1, z_1) \succ_1^* (x_1, y_1) \succ_1^* (y_1, x_1)$, $(x_2, z_2) \succ_2^* (x_2, y_2) \succ_2^* (y_2, x_2)$, and $(x_1, x_2, y_3, y_4, y_5) \mathcal{P} (z_1, z_2, x_3, x_4, x_5)$ but $\text{Not}[(x_1, x_2, y_3, y_4, y_5) \mathcal{P} (z_1, y_2, x_3, x_4, x_5)]$. \diamond

Example 49 (*Not[MM5]*)

Let $X = \prod_{i=1}^3 X_i$ with $X_1 = \{x_1, y_1, z_1\}$ and $X_j = \{x_j, y_j\}$ for $j = 2, 3$. The relation \mathcal{P} is defined as follows. For all $a, b \in X$, we have $a \mathcal{P} b$ if $\sum_{i=1}^3 p_i(a_i, b_i) > 0$, where p_i is defined in the following table for all pairs (a_i, b_i) in $X_i \times X_i$.

p_i	X_1	X_2	X_3
3	(x_1, z_1)		
2	(x_1, y_1)		
1	(y_1, z_1)	(x_2, y_2)	(x_3, y_3)
0	(a_1, a_1)	(a_2, a_2)	(a_3, a_3)
-1	(z_1, y_1)	(y_2, x_2)	(y_3, x_3)
-10	$(y_1, x_1), (z_1, x_1)$		

The relation \mathcal{P} is asymmetric, since the opposite (y_1, x_1) and (z_1, x_1) of the bonus pairs (x_1, y_1) and (x_1, z_1) are never involved in pairs belonging to \mathcal{P} .

For all i , \succ_i^* is the weak order induced by p_i in the above table. For $i = 2, 3$, $\succ_i^* = \succ_i^{**}$ while \succ_1^{**} separates the bottom class of \succ_1^* in two classes since $(y_1, x_1) \succ_1^{**} (z_1, x_1)$. Hence \mathcal{P} satisfies *RC1* and *RC2*. It is easy to check that *AC1*, *AC2*, and *AC3* hold.

MM3_i is trivially satisfied for $i = 2, 3$ (in view of Lemma 28). For $i = 1$, the only super negative pairs are (y_1, x_1) and (z_1, x_1) and they are never involved in \mathcal{P} . Therefore, using Lemma 28, we have that *Maj3₁* and, hence, *MM3_i*, hold.

Maj4_i is trivially satisfied for $i = 2, 3$. For $i = 1$, the only bonus pairs are (x_1, y_1) and (x_1, z_1) and their opposite pairs are never involved in \mathcal{P} . Therefore, applying Lemma 32 yields *Maj4₁* and, hence, *MM4*.

Maj5_i and, hence, *MM5_i* are trivially satisfied for $i = 2, 3$ but this example violates *Maj5₁*. Indeed, we have $(x_1, y_1) \succ_1^* (y_1, z_1) \succ_1^* (z_1, y_1)$ but there is a pair

above (x_1, y_1) in \succsim_1^* , namely, $(x_1, z_1) \succ_1^* (x_1, y_1)$. Since, $RC1$ and $RC2$ hold, this implies a violation of $MM5_1$ (Lemma 33).

In view of Lemma 38, property BC_{ij} is trivially satisfied for all $i \neq j$ since bonuses occur on the sole attribute X_1 . \diamond

Example 50 (Not[MM4])

Let $X = \prod_{i=1}^3 X_i$ with $X_1 = \{x_1, y_1, z_1\}$ and $X_j = \{x_j, y_j\}$ for $j = 2, 3$. The relation \mathcal{P} is defined as follows. For all $a, b \in X$, we have $a \mathcal{P} b$ if $\sum_{i=1}^3 p_i(a_i, b_i) > 0$, where p_i is defined in the following table for all pairs (a_i, b_i) in $X_i \times X_i$.

p_i	X_1	X_2	X_3
2	(x_1, z_1)		
1	$(x_1, y_1), (y_1, z_1)$	(x_2, y_2)	(x_3, y_3)
0	(a_1, a_1)	(a_2, a_2)	(a_3, a_3)
-1	$(y_1, x_1), (z_1, y_1), (z_1, x_1)$	(y_2, x_2)	(y_3, x_3)

When comparing a and b , provided (x_1, z_1) is neither (a_1, b_1) nor (b_1, a_1) , we have $\sum_{i=1}^3 p_i(b_i, a_i) = -\sum_{i=1}^3 p_i(a_i, b_i)$. Hence, at most one of the pairs (a, b) or (b, a) belongs to \mathcal{P} . If $a_1 = x_1$ and $b_1 = z_1$, there is only one case in which we have $b \mathcal{P} a$, namely $(z_1, x_2, x_3) \mathcal{P} (x_1, y_2, y_3)$. In this case, we do not have $a \mathcal{P} b$ since $\text{Not}[(x_1, y_2, y_3) \mathcal{P} (z_1, x_2, x_3)]$. Relation \mathcal{P} is thus asymmetric.

For all i , \succsim_i^* is the weak order induced by function p_i defined in the above table. For $i = 2, 3$, $\succsim_i^* = \succsim_i^{**}$ while \succsim_1^{**} separates the bottom class of \succsim_1^* in two classes since $[(y_1, x_1) \sim_1^{**} (z_1, y_1)] \succ_1^{**} (z_1, x_1)$. Hence \mathcal{P} satisfies $RC1$ and $RC2$. It is easy to check that $AC1$, $AC2$, and $AC3$ hold.

$Maj3_i$ and, hence, $MM3_i$ are trivially satisfied for all i since there are no super negative pairs at all (Lemma 28).

$Maj4_i$ and, hence, $MM4_i$ are trivially satisfied for $i = 2, 3$. $Maj4_1$ is violated for $i = 1$. Indeed, we have $(x_1, z_1) \succ_1^* (x_1, y_1) \succ_1^* (y_1, x_1)$ while $(z_1, x_2, x_3) \mathcal{P} (x_1, y_2, y_3)$, which, combined with Lemma 32, implies that $Maj4_1$ is not true. Since $RC1$ and $RC2$ hold, this implies a violation of $MM4_1$.

$Maj5_i$ is trivially satisfied for $i = 2, 3$. For $i = 1$, (x_1, z_1) is the only bonus and for all (r_1, s_1) , we have $(x_1, z_1) \succsim_1^* (r_1, s_1)$. Therefore, using Lemma 33, we have established that $Maj5_1$ and, hence, $MM5_i$ hold.

In view of Lemma 38, property BC_{ij} is trivially satisfied for all $i \neq j$ since a bonus occurs on the sole attribute X_1 . \diamond

Example 51 (Not[RC2])

Let $X = X_1 \times X_2$ with $X_1 = \{a, b\}$ and $X_2 = \{x, y\}$. Let \mathcal{P} be empty except that we have $(a, x) \mathcal{P} (a, y)$ and $(a, x) \mathcal{P} (b, y)$. The relation is clearly asymmetric.

On attribute 1, we have $[(a, a) \sim_1^* (a, b)] \succ_1^* [(b, a) \sim_1^* (b, b)]$. This shows that $RC1_1$ holds but that $RC2_1$ is violated. We have $a \succ_1^+ b$ and $a \succ_1^- b$. Hence, $AC1_1$,

$AC2_1$ and $AC3_1$ hold. It is easy to check that UC_1 and LC_1 hold so that $MM3_1$, $MM4_1$, $MM5_1$, and BC_{12} hold.

On attribute 2, we have $(x, y) \succ_2^* [(x, x) \sim_2^* (y, x) \sim_2^* (y, y)]$. This shows that $RC1_2$ and $RC2_2$ hold. We have $x \succ_2^+ y$ and $x \succ_2^- y$. Hence, $AC1_2$, $AC2_2$ and $AC3_2$ hold. It is easy to check that UC_2 and LC_2 hold so that $MM3_2$, $MM4_2$, $MM5_2$, and BC_{21} hold. \diamond

Example 52 (Not[RC1])

(This is also Example 92 in Bouyssou and Pirlot (2013)). Let $X = X_1 \times X_2 \times X_3$ with $X_1 = \{x, y, z, w\}$, $X_2 = \{a, b\}$ and $X_3 = \{p, q\}$. Let \mathcal{P} be such that $(\alpha, a, p) \mathcal{P} (\beta, a, q)$, $(\alpha, a, p) \mathcal{P} (\beta, b, p)$, $(\alpha, a, p) \mathcal{P} (\beta, b, q)$, $(\alpha, a, q) \mathcal{P} (\beta, b, q)$, $(\alpha, b, p) \mathcal{P} (\beta, b, q)$, for all $(\alpha, \beta) \in \Gamma_1 = X_1^2 \setminus \{(x, y), (z, y), (z, w)\}$.

We add the following pairs: $(x, a, p) \mathcal{P} (y, b, p)$, $(x, a, p) \mathcal{P} (y, b, q)$, $(x, a, q) \mathcal{P} (y, b, q)$, and $(z, a, p) \mathcal{P} (w, a, q)$, $(z, a, p) \mathcal{P} (w, b, q)$, $(z, b, p) \mathcal{P} (w, b, q)$.

This relation has a total of 71 ordered pairs. It is easy to check that it is asymmetric.

On attribute 2, we have $(a, b) \succ_2^* [(a, a) \sim_2^* (b, b)] \succ_2^* (b, a)$. This shows that $RC1_2$ and $RC2_2$ hold. We have $a \succ_2^+ b$ and $a \succ_2^- b$. Hence, $AC1_2$, $AC2_2$ and $AC3_2$ hold. It is easy to check that UC_2 and LC_2 hold so that $MM3_2$, $MM4_2$, $MM5_2$, BC_{21} , and BC_{23} hold.

On attribute 3, we have $(p, q) \succ_3^* [(p, p) \sim_3^* (q, q)] \succ_3^* (q, p)$. This shows that $RC1_3$ and $RC2_3$ hold. We have $p \succ_3^+ q$ and $p \succ_3^- q$. Hence, $AC1_3$, $AC2_3$ and $AC3_3$ hold. It is easy to check that UC_3 and LC_3 hold so that $MM3_3$, $MM4_3$, $MM5_3$, BC_{31} , and BC_{32} hold.

On attribute 1, all ordered pairs, except (x, y) , (z, y) and (z, w) , are in the same equivalence class of \sim_1^* and above these three pairs. The ordered pairs (x, y) and (z, w) are clearly incomparable w.r.t. \lesssim_1^* and are both above (z, y) .

This shows that $RC1_1$ fails but that $RC2_1$ holds. It is not difficult to check that we have $[y \sim_1^+ w] \succ_1^+ x \succ_1^+ z$ and $y \succ_1^- w \succ_1^- [x \sim_1^+ z]$. Hence, $AC1_1$, $AC2_1$ and $AC3_1$ hold. It is easy to see that UC_1 holds. Hence, $MM4_1$, $MM5_1$, BC_{12} , and BC_{13} hold.

It remains to check that $MM3_1$ holds. Let us check that $M3_1$ holds.

Suppose that $(x_1, a_{-1}) \mathcal{P} (y_1, b_{-1})$, $(y_1, c_{-1}) \mathcal{P} (x_1, d_{-1})$ and $(z_1, e_{-1}) \mathcal{P} (w_1, f_{-1})$. We want to show that we have one of the following three relations: $(y_1, a_{-1}) \mathcal{P} (x_1, b_{-1})$, $(z_1, a_{-1}) \mathcal{P} (w_1, b_{-1})$, $(z_1, c_{-1}) \mathcal{P} (w_1, d_{-1})$.

If $(y_1, x_1) \in \Gamma$ then the first conclusion always holds. If $(y_1, x_1) = (z, y)$, the second premise never holds, so that the condition is trivially satisfied. It remains to deal with the following two cases: $(y_1, x_1) = (x, y)$ and $(y_1, x_1) = (z, w)$. We deal with the first case, the treatment of the second being entirely similar.

Suppose that $(y_1, x_1) = (x, y)$. If $(z_1, w_1) \neq (z, y)$ and $(z_1, w_1) \neq (z, w)$, we have $(z_1, w_1) \lesssim_1^* (x, y)$, so that the third conclusion always holds. If $(z_1, w_1) =$

(z, w) , the third premise never holds, so that the condition is trivially satisfied. If $(z_1, w_1) = (z, y)$, it is easy to check that there are no $a_{-1}, b_{-1} \in X_{-1}$ such that $(y_1, a_{-1}) \mathcal{P} (x_1, b_{-1})$, $\text{Not}[(x_1, a_{-1}) \mathcal{P} (y_1, b_{-1})]$ and $\text{Not}[(z_1, a_{-1}) \mathcal{P} (w_1, b_{-1})]$, so that no violation of $M3_1$ is possible in this case. \diamond

5 Discussion

We have analyzed and characterized binary relations \mathcal{P} on product sets that can be obtained as the asymmetric part of an outranking relation \mathcal{S} . We have seen that these relations are more complex than outranking relations. Indeed, they can be seen as outranking relations only in degenerate cases.

Consider two alternatives (x_i, a_{-i}) and (y_i, b_{-i}) that are such that there is no relation V_j on the attributes $j \neq i$. This complexity stems from the fact that $y_i V_i x_i$ can have two different effects. If $S[(x_i, a_{-i}), (y_i, b_{-i})] \supset S[(y_i, b_{-i}), (x_i, a_{-i})]$, then $y_i V_i x_i$ acts as a normal veto. It forbids to have $(x_i, a_{-i}) \mathcal{P} (y_i, b_{-i})$ because the negative preference difference between y_i and x_i is “too large”. If $S[(x_i, a_{-i}), (y_i, b_{-i})] \triangleq S[(y_i, b_{-i}), (x_i, a_{-i})]$ then $y_i V_i x_i$ acts as a bonus. It implies that $(y_i, b_{-i}) \mathcal{P} (x_i, a_{-i})$, because it forbids to have $(x_i, a_{-i}) \mathcal{S} (y_i, b_{-i})$. This makes an AP-R-CDR-AT an object that is far more complex than an outranking relation.

This is reflected in our analysis by the fact that conditions *MM4*, *MM5* and *BC* are complex conditions, even though it remains possible to interpret them in terms of the relation \succsim_i^* . This complexity also explains the difference between the pseudo-disjunctive (also known as “optimistic”) and pseudo-conjunctive (also known as “pessimistic”) versions of ELECTRE TRI-B. While the latter uses a reflexive outranking relation (i.e., an R-CDR-AT), the former uses an AP-R-CDR-AT. Because these two objects are different, we should not be surprised by the fact that the theoretical analysis of the pseudo-conjunctive version given in Bouyssou and Marchant (2007a,b) does not carry over to the pseudo-disjunctive version. Similarly, we should not be surprised by the fact that most of elicitation techniques developed for the parameters of ELECTRE TRI-B only deal with the pseudo-conjunctive version (see Cailloux, Meyer, and Mousseau, 2012; Damart, Dias, and Mousseau, 2007; Dias and Clímaco, 2000; Dias and Mousseau, 2003, 2006; Dias, Mousseau, Figueira, and Clímaco, 2002; Leroy, Mousseau, and Pirlot, 2011; Mousseau and Dias, 2004; Mousseau and Słowiński, 1998; Mousseau, Słowiński, and Zieliński, 2000; Mousseau, Figueira, and Naux, 2001; Mousseau, Figueira, Dias, da Silva, and Clímaco, 2003; Ngo The and Mousseau, 2002). There has been a number of recent works proposing elicitation techniques for the pseudo-disjunctive version (see Zheng, 2012; Zheng, Takougang, Mousseau, and Pirlot, 2012). These exceptions do not invalidate the above remark, since these elicitation techniques are far

more complex than the ones developed for the pseudo-conjunctive version (they use MILP instead of LP). For more details on this point, we refer to Bouyssou and Marchant (2013).

A clear limitation of the present work as well as our previous works on outranking relations is that it only deals with *crisp* relations. Since there are outranking methods such as ELECTRE III (Roy, 1978) or PROMETHEE (Brans and Vincke, 1985; Brans, Vincke, and Mareschal, 1986) that use valued relations, an important direction for future research would be to analyze such relations. Although this will surely imply the development of a framework different from the ones used here, this does not seem to be out of reach.

References

- J. Almeida-Dias, J. R. Figueira, and B. Roy. ELECTRE TRI-C: A multiple criteria sorting method based on characteristic reference actions. *European Journal of Operational Research*, 204(3):565–580, 2010.
- D. Bouyssou and T. Marchant. An axiomatic approach to noncompensatory sorting methods in MCDM, I: The case of two categories. *European Journal of Operational Research*, 178(1):217–245, 2007a.
- D. Bouyssou and T. Marchant. An axiomatic approach to noncompensatory sorting methods in MCDM, II: More than two categories. *European Journal of Operational Research*, 178(1):246–276, 2007b.
- D. Bouyssou and T. Marchant. Some remarks on ELECTRE TRI. Cahier du LAMSADE 347, LAMSADE, Université Paris Dauphine, 2013. Available from <http://lamsade.dauphine.fr/~bouyssou>.
- D. Bouyssou and M. Pirlot. Conjoint measurement without additivity and transitivity. In N. Meskens and M. Roubens, editors, *Advances in Decision Analysis*, pages 13–29. Kluwer, Dordrecht, 1999.
- D. Bouyssou and M. Pirlot. Nontransitive decomposable conjoint measurement. *Journal of Mathematical Psychology*, 46:677–703, 2002a.
- D. Bouyssou and M. Pirlot. A characterization of strict concordance relations. In D. Bouyssou, É. Jacquet-Lagrèze, P. Perny, R. Słowiński, D. Vanderpooten, and Ph. Vincke, editors, *Aiding Decisions with Multiple Criteria: Essays in Honour of Bernard Roy*, pages 121–145. Kluwer, Dordrecht, 2002b.
- D. Bouyssou and M. Pirlot. Preferences for multiattributed alternatives: Traces, dominance, and numerical representations. *Journal of Mathematical Psychology*, 48:167–185, 2004a.
- D. Bouyssou and M. Pirlot. ‘Additive difference’ models without additivity and subtractivity. *Journal of Mathematical Psychology*, 48:263–291, 2004b.
- D. Bouyssou and M. Pirlot. A characterization of concordance relations. *European Journal of Operational Research*, 167(2):427–443, 2005a.

- D. Bouyssou and M. Pirlot. Following the traces: An introduction to conjoint measurement without transitivity and additivity. *European Journal of Operational Research*, 163(2):287–337, 2005b. Invited Review.
- D. Bouyssou and M. Pirlot. Further results on concordance relations. *European Journal of Operational Research*, 181:505–514, 2007.
- D. Bouyssou and M. Pirlot. An axiomatic analysis of concordance-discordance relations. *European Journal of Operational Research*, 199:468–477, 2009.
- D. Bouyssou and M. Pirlot. An axiomatic approach to TACTIC. *Studia Informatica Universalis*, 10(2):45–71, 2012.
- D. Bouyssou and M. Pirlot. On the relationship between strict and non-strict outranking relations. Cahier du LAMSADE 346, LAMSADE, Université Paris Dauphine, 2013. Available from <http://lamsade.dauphine.fr/~bouyssou>.
- D. Bouyssou, M. Pirlot, and Ph. Vincke. A general model of preference aggregation. In M. H. Karwan, J. Spronk, and J. Wallenius, editors, *Essays in Decision Making*, pages 120–134. Springer Verlag, Berlin, 1997.
- J.-P. Brans and Ph. Vincke. A preference ranking organisation method. (The PROMETHEE method for multiple criteria decision-making). *Management Science*, 31(6):647–656, 1985.
- J.-P. Brans, Ph. Vincke, and B. Mareschal. How to select and how to rank projects: the PROMETHEE method. *European Journal of Operational Research*, 24:228–238, 1986.
- O. Cailloux, P. Meyer, and V. Mousseau. Eliciting ELECTRE TRI category limits for a group of decision makers. *European Journal of Operational Research*, 223(1):133–140, 2012.
- S. Damart, L. C. Dias, and V. Mousseau. Supporting groups in sorting decisions: Methodology and use of a multi-criteria aggregation / disaggregation DSS. *Decision Support Systems*, 43(4):1464–1475, 2007.
- L. C. Dias and J. Clímaco. ELECTRE TRI for groups with imprecise information on parameter values. *Group Decision and Negotiation*, 9(5):355–377, 2000.
- L. C. Dias and V. Mousseau. IRIS: A DSS for multiple criteria sorting problems. *Journal of Multi-Criteria Decision Analysis*, 12:285–298, 2003.
- L. C. Dias and V. Mousseau. Inferring ELECTRE’s veto-related parameters from outranking examples. *European Journal of Operational Research*, 170(1):172–191, 2006.
- L. C. Dias, V. Mousseau, J. Figueira, and J. Clímaco. An aggregation / disaggregation approach to obtain robust conclusions with ELECTRE TRI. *European Journal of Operational Research*, 138(2):332–48, 2002.
- D. Dubois, H. Fargier, P. Perny, and H. Prade. A characterization of generalized concordance rules in multicriteria decision-making. *International Journal of Intelligent Systems*, 18(7):751–774, 2003.
- S. Greco, B. Matarazzo, and R. Słowiński. Axiomatic basis of noncompensatory preferences. Communication to *FUR X*, 30 May–2 June, Torino, Italy, 2001.
- A. Leroy, V. Mousseau, and M. Pirlot. Learning the parameters of a multiple criteria

- sorting method. In R. I. Brafman, F. S. Roberts, and A. Tsoukiàs, editors, *ADT 2011*, number 6992 in LNAI, pages 219–233. Springer, 2011.
- R. D. Luce. Semiorders and a theory of utility discrimination. *Econometrica*, 24:178–191, 1956.
- V. Mousseau and L. C. Dias. Valued outranking relations in ELECTRE providing manageable disaggregation procedures. *European Journal of Operational Research*, 156(2):467–482, 2004.
- V. Mousseau and R. Słowiński. Inferring an ELECTRE TRI model from assignment examples. *Journal of Global Optimization*, 12(2):157–174, 1998.
- V. Mousseau, R. Słowiński, and P. Zielniewicz. A user-oriented implementation of the ELECTRE TRI method integrating preference elicitation support. *Computers & Operations Research*, 27(7–8):757–777, 2000.
- V. Mousseau, J. Figueira, and J.-Ph. Naux. Using assignment examples to infer weights for ELECTRE TRI method: Some experimental results. *European Journal of Operational Research*, 130(2):263–275, 2001.
- V. Mousseau, J. R. Figueira, L. C. Dias, C. Gomes da Silva, and J. Clímaco. Resolving inconsistencies among constraints on the parameters of an MCDA model. *European Journal of Operational Research*, 147(1):72–93, 2003.
- A. Ngo The and V. Mousseau. Using assignment examples to infer category limits for the ELECTRE TRI method. *Journal of Multi-Criteria Decision Analysis*, 11(1):29–43, 2002.
- B. Roy. Classement et choix en présence de points de vue multiples (la méthode ELECTRE). *RIRO*, 2:57–75, 1968.
- B. Roy. ELECTRE III : un algorithme de classement fondé sur une représentation floue des préférences en présence de critères multiples. *Cahiers du CERO*, 20:3–24, 1978.
- B. Roy. Présentation et interprétation de la méthode ELECTRE TRI pour affecter des zones dans des catégories de risque. Document du LAMSADE 124, LAMSADE, Université Paris Dauphine, Paris, France, March 2002. 25 pages.
- B. Roy and P. Bertier. La méthode ELECTRE II : une application au media-planning. In M. Ross, editor, *OR'72*, pages 291–302. North Holland, Amsterdam, 1973.
- B. Roy and D. Bouyssou. *Aide multicritère à la décision : Méthodes et cas*. Economica, Paris, 1993.
- J.-C. Vansnick. On the problems of weights in MCDM (the noncompensatory approach). *European Journal of Operational Research*, 24:288–294, 1986.
- J. Zheng. *Preference Elicitation for reference based aggregation models: Algorithms and Procedures*. PhD thesis, École Centrale, Paris, 2012.
- J. Zheng, S. A. Metchebon Takougang, V. Mousseau, and M. Pirlot. Learning an optimistic ELECTRE TRI sorting rule. Research Report CER 12–06, École Centrale, Paris, Laboratoire de Génie Industriel, 2012.