

Cooperative Games

Lecture 4: Game with Coalition structures, Core and Bargaining Set

Stéphane Airiau

ILLC - University of Amsterdam



Today

- If agents desire the kind of stability offered by the core, they will be unable to reach an agreement.
- they have no choice but to **relax** their stability requirements.
- Need a solution that allows agents to reach an agreement, but maintain some stability.
- First we will consider the problem of stability of coalition structure, i.e., a partition of the set of agents is formed first, and then, members of a coalition negotiate their payoff.
- Then, we will consider the bargaining set, which relaxes the requirements of the core.

Coalition Structure

Definition (Coalition Structure)

A **coalition structure (CS)** is a partition of the grand coalition into coalitions.
 $\mathcal{S} = \{C_1, \dots, C_k\}$ where $\cup_{i \in \{1, \dots, k\}} C_i = N$ and $i \neq j \Rightarrow C_i \cap C_j = \emptyset$.
 We note \mathcal{S}_N the set of all coalition structures over the set N .

ex: $\{\{1, 3, 4\}, \{2, 7\}, \{5\}, \{6, 8\}\}$ is a coalition structure for $n = 8$ agents.

We will study three solution concepts: the **bargaining set**, the **nucleolus** and the **kernel**. They form the "**bargaining set family**" and we will see later why. In addition, the definition of each of these solution concepts uses a CS.

We start by defining a game with coalition structure, and see how we can define the core of such a game. Then, we'll start studying the bargaining set family.

Game with Coalition Structure

Definition (TU game)

A TU game is a pair (N, v) where N is a set of agents and where v is a valuation function.

Definition (Game with Coalition Structures)

A **TU-game with coalition structure (N, v, \mathcal{S})** consists of a TU game (N, v) and a CS $\mathcal{S} \in \mathcal{S}_N$.

- We assume that the players agreed upon the formation of \mathcal{S} and only the payoff distribution choice is left open.
- The CS may model affinities among agents, may be due to external causes (e.g. affinities based on locations).
- The agents may refer to the value of coalitions with agents outside their coalition (i.e. opportunities they would have outside of their coalition).
- (N, v) and $(N, v, \{N\})$ represent the same game.

Definition (core of a game (N, v))

The core of a TU game (N, v) is defined as
 $Core(N, v) = \{x \in \mathbb{R}^n \mid x(N) \leq v(N) \wedge x(C) \geq v(C) \forall C \subseteq N\}$

The set of **feasible** payoff vectors for (N, v, \mathcal{S}) is
 $X_{(N, v, \mathcal{S})} = \{x \in \mathbb{R}^n \mid \text{for every } C \in \mathcal{S} \ x(C) \leq v(C)\}$.

Definition (Core of a game with CS)

The **core $Core(N, v, \mathcal{S})$** of (N, v, \mathcal{S}) is defined by
 $\{x \in \mathbb{R}^n \mid (\forall C \in \mathcal{S}, x(C) \leq v(C)) \text{ and } (\forall C \subseteq N, x(C) \geq v(C))\}$

We have $Core(N, v, \{N\}) = Core(N, v)$.

The next theorems are due to Aumann and Drèze.

R.J. Aumann and J.H. Drèze. **Cooperative games with coalition structures**, *International Journal of Game Theory*, 1974

Definition (Superadditive cover)

The **superadditive cover** of (N, v) is the game (N, \hat{v}) defined by

$$\begin{cases} \hat{v}(C) = \max_{\mathcal{P} \in \mathcal{S}_N} \left\{ \sum_{T \in \mathcal{P}} v(T) \right\} & \forall C \subseteq N \setminus \emptyset \\ \hat{v}(\emptyset) = 0 \end{cases}$$

- We have $\hat{v}(N) = \max_{\mathcal{P} \in \mathcal{S}_N} \left\{ \sum_{T \in \mathcal{P}} v(T) \right\}$, i.e., $\hat{v}(N)$ is the maximum value that can be produced by N . We call it the **value of the optimal coalition structure**.
- The superadditive cover is a superadditive game (**why?**).

Theorem

Let (N, v, \mathcal{S}) be a game with coalition structure. Then

- $Core(N, v, \mathcal{S}) \neq \emptyset$ iff $Core(N, \hat{v}) \neq \emptyset \wedge \hat{v}(N) = \sum_{C \in \mathcal{S}} v(C)$
- if $Core(N, v, \mathcal{S}) \neq \emptyset$, then $Core(N, v, \mathcal{S}) = Core(N, \hat{v})$

Proof of part a)

\Rightarrow Let $x \in Core(N, v, \mathcal{S})$. We show that $x \in Core(N, \hat{v})$ as well. Let $C \subseteq N \setminus \emptyset$ and $\mathcal{P}_C \in \mathcal{S}_C$ be a partition of C .

By definition of the core, for every $S \subseteq N$ $x(S) \geq v(S)$.
 $x(C) = \sum_{i \in C} x_i = \sum_{S \in \mathcal{P}_C} x(S) \geq \sum_{S \in \mathcal{P}_C} v(S)$, which is valid for all

partitions of C . Hence, $x(C) \geq \max_{\mathcal{P}_C \in \mathcal{S}_C} \sum_{S \in \mathcal{P}_C} v(S) = \hat{v}(C)$.

We have just proved $\forall C \subseteq N \setminus \emptyset, x(C) \geq \hat{v}(C)$, and so x is **group rational**. ✓

We now need to prove that $\hat{v}(N) = \sum_{C \in \mathcal{S}} v(C)$.

$x(N) = \sum_{C \in \mathcal{S}} v(C)$ since x is in the core of (N, v, \mathcal{S}) (efficient).

Applying the inequality above, we have

$$x(N) = \sum_{C \in \mathcal{S}} v(C) \geq \hat{v}(N).$$

Applying the definition of the valuation function \hat{v} , we have

$$\hat{v}(N) \geq \sum_{C \in \mathcal{S}} v(C). \text{ Consequently, } \hat{v}(N) = \sum_{C \in \mathcal{S}} v(C) \text{ and it follows that } x \text{ is } \textbf{efficient}$$

Hence $x \in Core(N, \hat{v})$.

Proof of part a)

\Leftarrow Let's assume $x \in Core(N, \hat{v})$ and $\hat{v}(N) = \sum_{C \in \mathcal{S}} v(C)$.

We need to prove that $x \in Core(N, v, \mathcal{S})$.

For every $C \subseteq N, x(C) \geq \hat{v}(C)$ since x is in the core of $Core(N, \hat{v})$. Then $x(C) \geq \max_{\mathcal{P}_C \in \mathcal{S}_C} \sum_{S \in \mathcal{P}_C} v(S) \geq v(C)$ using $\{C\}$

as a partition of C .

This proves x is group rational. ✓

$x(N) = \hat{v}(N) = \sum_{C \in \mathcal{S}} v(C)$ since x is efficient.

It follows that $\forall C \in \mathcal{S}, x(C) \geq v(C)$, which proves x is feasible for the CS \mathcal{S} , and that x is efficient. ✓

- proof of part b): we have just proved that $x \in Core(N, \hat{v})$ implies that $x \in Core(N, v, \mathcal{S})$ and $x \in Core(N, v, \mathcal{S})$ implies that $x \in Core(N, \hat{v})$. This proves that if $Core(N, v, \mathcal{S}) \neq \emptyset, Core(N, \hat{v}) = Core(N, v, \mathcal{S})$.

Definition (Substitutes)

Let (N, v) be a game and $(i, j) \in N^2$. Agents i and j are **substitutes** iff $\forall \mathcal{C} \subseteq N \setminus \{i, j\}, v(\mathcal{C} \cup \{i\}) = v(\mathcal{C} \cup \{j\})$.

A nice property of the core related to fairness:

Theorem

Let (N, v, \mathcal{S}) be a game with coalition structure, let i and j be substitutes, and let $x \in \text{Core}(N, v, \mathcal{S})$. If i and j belong to different members of \mathcal{S} , then $x_i = x_j$.

Proof

Let $(i, j) \in N^2$ be substitutes, $\mathcal{C} \in \mathcal{S}$ such that $i \in \mathcal{C}$ and $j \notin \mathcal{C}$. Let $x \in \text{Core}(N, v, \mathcal{S})$. Since i and j are substitutes, we have $v((\mathcal{C} \setminus \{i\}) \cup \{j\}) = v(\mathcal{C})$. Since $x \in \text{Core}(N, v, \mathcal{S})$, we have $\forall \mathcal{C} \subseteq N, x(\mathcal{C}) \geq v(\mathcal{C})$, we apply this to $(\mathcal{C} \setminus \{i\}) \cup \{j\}$: $0 \geq v((\mathcal{C} \setminus \{i\}) \cup \{j\}) - x((\mathcal{C} \setminus \{i\}) \cup \{j\}) = v(\mathcal{C}) - x(\mathcal{C}) + x_i - x_j$, since $\mathcal{C} \in \mathcal{S}$ and $x \in \text{Core}(N, v, \mathcal{S})$, $x(\mathcal{C}) = v(\mathcal{C})$ we have $x_j \geq x_i$. Since i and j play symmetric roles, we have $x_i = x_j$. ✓ □

A second solution concept:

The bargaining set.

R.J. Aumann and M. Maschler. **The bargaining set for cooperative games**, in *Advances in game theory (Annals of mathematics study)*, 1964.

M. Davis and M. Maschler. **Existence of stable payoff configurations for cooperative games**, *Bulletin of the American mathematical society*, 1963.

Let (N, v, \mathcal{S}) be a game with coalition structure and x an imputation. The bargaining set models stability in the following sense:

Any **argument** from an agent i against a payoff distribution x is of the following form:

I get too little in the imputation x , and agent j gets too much! I can form a coalition that excludes j in which some members benefit and all members are at least as well off as in x .

The argument is **ineffective** for the bargaining set if agent j can answer the following:

I can form a coalition that excludes agent i in which all agents are at least as well off as in x , and as well off as in the payoff proposed by i for those who were offered to join i in the argument.

Definition (Objection)

Let (N, v, \mathcal{S}) be a game with coalition structure, $x \in X_{(N, v, \mathcal{S})}$ (the set of all feasible payoff vectors for (N, v, \mathcal{S})), $\mathcal{C} \in \mathcal{S}$ be a coalition, and i and j two distinct members of \mathcal{C} ($(i, j) \in \mathcal{C}^2, i \neq j$).

An **objection of i against j** is a pair (P, y) where

- $P \subseteq N$ is a coalition such that $i \in P$ and $j \notin P$.
- $y \in \mathbb{R}^P$ where p is the size of P
- $y(P) \leq v(P)$ (y is a feasible payoff distribution for the agents in P)
- $\forall k \in P, y_k \geq x_k$ and $y_i > x_i$ (agent i strictly benefits from y , and the other members of P do not do worse in y than in x .)

An objection (P, y) of i against j is a **potential threat** by coalition P , which contains i but not j , to deviate from x . The goal is not to change \mathcal{S} , but to obtain a side payment from j to i , i.e., to modify x within $X_{(N, v, \mathcal{S})}$.

Definition (Counter-objection)

An **counter-objection to (P, y)** is a pair (Q, z) where

- $Q \subseteq N$ is a coalition such that $j \in Q$ and $i \notin Q$.
- $z \in \mathbb{R}^Q$ where q is the size of Q
- $z(Q) \leq v(Q)$ (z is a feasible payoff distribution for the agents in Q)
- $\forall k \in Q, z_k \geq x_k$ (the members of Q get at least the value in x)
- $\forall k \in Q \cap P, z_k \geq y_k$ (the members of Q which are also members of P get at least the value promised in the objection)

In a counter-objection, agent j must show that it can protect its payoff x_j in spite of the existing objection of i .

Definition (Stability)

Let (N, v, \mathcal{S}) a game with coalition structure. A vector $x \in X_{(N, v, \mathcal{S})}$ is **stable** iff for each objection at x there is a counter-objection.

Definition (Pre-bargaining set)

The **pre-bargaining set (preBS)** is the set of all stable members of $X_{(N, v, \mathcal{S})}$.

Lemma

Let (N, v, \mathcal{S}) a game with coalition structure, we have $\text{Core}(N, v, \mathcal{S}) \subseteq \text{preBS}(N, v, \mathcal{S})$.

This is true since, if $x \in \text{Core}(N, v, \mathcal{S})$, no agent i has any objection against any other agent j .

Example

Let (N, v) be a 7-player simple majority game, i.e.

$$v(\mathcal{C}) = \begin{cases} 1 & \text{if } |\mathcal{C}| \geq 4 \\ 0 & \text{otherwise} \end{cases}$$

Let us consider $x = (\frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \dots, \frac{1}{5})$. It is clear that $x(N) = 1$.

Let us prove that x is in the **pre-bargaining set** of the game $(N, v, \{N\})$.

Objections within members of $\{2, 3, 4, 5, 6, 7\}$ will have a counter-objection by symmetry. ✓

Let us consider the objections (P, y) of 1 against another member of $\{2, 3, 4, 5, 6, 7\}$. Since the players $\{2, \dots, 7\}$ play symmetric roles, we consider an objection of 1 against 7 using successively $P = \{1, 2, 3, 4, 5, 6\}, P = \{1, 2, 3, 4, 5\}, P = \{1, 2, 3, 4\}, P = \{1, 2, 3\}, P = \{1, 2\}$ and $P = \{1\}$. We will look for a counter-objection of player 7 using (Q, z) .

- We consider that $P = \{1, 2, 3, 4, 5, 6\}$. We need to find the payoff vector $y \in \mathbb{R}^6$ so that (P, y) is an objection.

$$y = (\alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \dots, \frac{1}{5} + \alpha_6),$$

The conditions for (P, y) to be an objection are the following:

- each agent is as well off as in x : $\alpha > -\frac{1}{5}, \alpha_i \geq 0$
- y is feasible for coalition P : $\sum_{i=2}^6 (\alpha_i + \frac{1}{5}) + \alpha \leq 1$.

w.l.o.g $0 \leq \alpha_2 \leq \alpha_3 \leq \alpha_4 \leq \alpha_5 \leq \alpha_6$.

$$\text{Then } \sum_{i=2}^6 (\frac{1}{5} + \alpha_i) + \alpha = \frac{5}{5} + \sum_{i=2}^6 \alpha_i + \alpha = 1 + \sum_{i=2}^6 \alpha_i + \alpha \leq 1.$$

$$\text{Then } \sum_{i=2}^6 \alpha_i \leq -\alpha < \frac{1}{5}.$$

- We need to find a counter-objection for (P, y) .

claim: we can choose $Q = \{2, 3, 4, 7\}$ and $z = (\frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5} + \alpha_4, \frac{1}{5} + \alpha_5)$

$$z(Q) = \frac{1}{5} + \alpha_2 + \frac{1}{5} + \alpha_3 + \frac{1}{5} + \alpha_4 + \frac{1}{5} + \alpha_5 = \frac{4}{5} + \sum_{i=2}^5 \alpha_i \leq 1 \text{ since } \sum_{i=2}^5 \alpha_i \leq \sum_{i=2}^6 \alpha_i < \frac{1}{5} \text{ so } z \text{ is feasible.}$$

It is clear that $\forall i \in Q, z_i \geq x_i$ and that $\forall i \in Q \cap P, z_i \geq y_i$. ✓ Hence, (Q, z) is a counter-objection. ✓

- Now, let us consider that $P = \{1, 2, 3, 4, 5\}$. The vector $y = (\alpha, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5} + \alpha_4, \frac{1}{5} + \alpha_5)$ is an objection when

$$\alpha > -\frac{1}{5}, \alpha_i \geq 0, \sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha \leq 1$$

This time, we have $\sum_{i=2}^5 (\frac{1}{5} + \alpha_i) + \alpha = \frac{4}{5} + \sum_{i=2}^5 \alpha_i + \alpha \leq 1$

then $\sum_{i=2}^5 \alpha_i \leq 1 - \frac{4}{5} - \alpha = \frac{1}{5} - \alpha$ and finally $\sum_{i=2}^5 \alpha_i \leq \frac{1}{5} - \alpha < \frac{2}{5}$.

- We need to find a counter-objection to (P, y)

claim: we can choose $Q = \{2, 3, 6, 7\}$, $z = (\frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3, \frac{1}{5}, \frac{1}{5})$

It is clear that $\forall i \in Q, z_i \geq x_i$ and $\forall i \in P \cap Q, z_i \geq y_i$ (for agent 2 and 3).

$z(Q) = \frac{1}{5} + \alpha_2 + \frac{1}{5} + \alpha_3 + \frac{1}{5} + \frac{1}{5} = \frac{4}{5} + \alpha_2 + \alpha_3$. We have $\alpha_2 + \alpha_3 < \frac{1}{5}$, otherwise, we would have $\alpha_2 + \alpha_3 \geq \frac{1}{5}$ and since the α_i are

ordered, we would then have $\sum_{i=2}^5 \alpha_i \geq \frac{2}{5}$, which is not possible.

Hence $z(Q) \leq 1$ which proves z is feasible ✓

Using similar arguments, we find a counter-objection for each other objections (you might want to fill in the details at home).

- $P = \{1, 2, 3, 4\}$, $y = (\alpha, \frac{1}{5} + \alpha_1, \frac{1}{5} + \alpha_2, \frac{1}{5} + \alpha_3)$, $\alpha > -\frac{1}{5}$, $\alpha_i \geq 0$, $\sum_{i=2}^4 \alpha_i + \alpha \leq \frac{2}{5} \Rightarrow \sum_{i=2}^4 \alpha_i \leq \frac{2}{5} - \alpha < \frac{3}{5}$.
- $Q = \{2, 5, 6, 7\}$, $z = (\frac{1}{5} + \alpha_2, \frac{1}{5}, \frac{1}{5}, \frac{1}{5})$ since $\alpha_2 \leq \frac{1}{5}$
- $|P| \leq 3$ $P = \{1, 2, 3\}$, $v(P) = 0$, $y = (\alpha, \alpha_1, \alpha_2)$, $\alpha > -\frac{1}{5}$, $\alpha_i \geq 0$, $\alpha_1 + \alpha_2 \leq -\alpha < \frac{1}{5}$
- $Q = \{4, 5, 6, 7\}$, $z = (\frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5})$ will be a counter argument (1 cannot provide more than $\frac{1}{5}$ to any other agent).
- For each possible objection of 1, we found a counter-objection. Using similar arguments, we can find a counter-objection to any objection of player 7 against player 1.
- $x \in \text{preBS}(N, v, S)$ ✓

Bargaining set

In the example, agent 1 gets $-\frac{1}{5}$ when $v(\mathcal{C}) \geq 0$ for all coalition $\mathcal{C} \subseteq N!$ This shows that the pre-bargaining set may **not** be individually rational.

Let $I(N, v, S) = \{x \in X_{(N, v, S)} \mid x_i \geq v(\{i\}) \forall i \in N\}$ be the **set of individually rational payoff vector** in $X_{(N, v, S)}$.

Lemma

If a game is weakly superadditive, $I(N, v, S) \neq \emptyset$.

Definition (Bargaining set)

Let (N, v, S) a game in coalition structure.

The **bargaining set (BS)** is defined by $BS(N, v, S) = I(N, v, S) \cap \text{preBS}(N, v, S)$.

Lemma

We have $\text{Core}(N, v, S) \subseteq BS(N, v, S)$.

Theorem

Let (N, v, S) a game with coalition structure. Assume that $I(N, v, S) \neq \emptyset$. Then the bargaining set $BS(N, v, S) \neq \emptyset$.

Proof

It is possible to give a direct proof of this theorem (a bit long, (Section 4.2 in **Introduction to the Theory of Cooperative Games**)).

We will show this result in a different way in the lecture about the nucleolus next week. □

B. Peleg and P. Sudhölter **Introduction to the Theory of Cooperative Games**, Springer, 2007.

Definition (weighted voting games)

A game $(N, w_{i \in N}, q, v)$ is a **weighted voting game** when v satisfies unanimity, monotonicity and the valuation function is defined as

$$v(S) = \begin{cases} 1 & \text{when } \sum_{i \in S} w_i \geq q \\ 0 & \text{otherwise} \end{cases}$$

We note such a game by $(q; w_1, \dots, w_n)$

Let (N, v) be the game associated with the 6-player weighted majority game $(3; 1, 1, 1, 1, 1, 0)$. Agent 6 is a null player since its weight is 0. Nevertheless $x = (\frac{1}{7}, \dots, \frac{1}{7}, \frac{2}{7}) \in BS(N, v)$.

Show it at home (a solution will be posted online). We need to consider all objections (P, y) from the an agent against the null agent, and find a counter objection (Q, z) .

Agent 6 is a dummy, however, it receives a payoff of $\frac{2}{7}$, which is larger than agents who are not dummy!

Remember: $mc_i^{\max} = \max_{\mathcal{C} \subseteq N \setminus \{i\}} v(\mathcal{C} \cup \{i\}) - v(\mathcal{C})$

x is **reasonable from above** if $\forall i \in N, x^i < mc_i^{\max}$

⇔ mc_i^{\max} is the strongest **threat** that an agent can use against a coalition.

The bargaining set is not **Reasonable from above**: the dummy agent gets more than $\max_{\mathcal{C} \subseteq N \setminus \{6\}} (v(\mathcal{C} \cup \{6\}) - v(\mathcal{C})) = 0$. ✗

Summary

- We introduced the notion of games with coalition structures.
- We looked at the definition of the core, i.e., stability of the coalition structure. Games with coalition structure may have an empty core (e.g., $(N, v, [N])$ and v in exercise 4 of homework 1).
- We introduced the bargaining set, and looked at some examples.
 - pros:** it is guaranteed to be non-empty, when the core is non-empty, it is contained in the bargaining set.
 - cons:** it may not be reasonable from above.

Coming next

- We will consider the Nucleolus. It can also be defined in terms of objections and counter objections, but the nature of the objection is different from the bargaining set.