

# Cooperative Games

## Lecture 7: The Kernel (end) and The Shapley Value

Stéphane Airiau

ILLC - University of Amsterdam



Today

- We prove one property of the kernel, and we consider an algorithm to compute an element in the kernel
- We introduce a solution concept called the Shapley value.

Last week

### Theorem

Let  $(N, v, \mathcal{S})$  a game with coalition structure, and let  $\mathcal{I}mp \neq \emptyset$ . Then we have:

- (i)  $Nu(N, v, \mathcal{S}) \subseteq K(N, v, \mathcal{S})$   
Proof ✓
- (ii)  $K(N, v, \mathcal{S}) \subseteq BS(N, v, \mathcal{S})$   
Proof ✗

Proof of (ii)

Let  $(N, v, \mathcal{S})$  a TU game with coalition structure. Let  $x \in K(N, v, \mathcal{S})$ . We want to prove that  $x \in BS(N, v, \mathcal{S})$ . To do so, we need to show that for any objection  $(P, y)$  from any player  $i$  against any player  $j$  at  $x$ , there is a counter objection  $(Q, z)$  to  $(P, y)$ . For the bargaining set, An **objection of  $i$  against  $j$**  is a pair  $(P, y)$  where

- $P \subseteq N$  is a coalition such that  $i \in P$  and  $j \notin P$ .
- $y \in \mathbb{R}^P$  where  $p$  is the size of  $P$
- $y(P) \leq v(P)$  ( $y$  is a feasible payoff for members of  $P$ )
- $\forall k \in P, y_k \geq x_k$  and  $y_i > x_i$

An **counter-objection to  $(P, y)$**  is a pair  $(Q, z)$  where

- $Q \subseteq N$  is a coalition such that  $j \in Q$  and  $i \notin Q$ .
- $z \in \mathbb{R}^Q$  where  $q$  is the size of  $Q$
- $z(Q) \leq v(Q)$  ( $z$  is a feasible payoff for members of  $Q$ )
- $\forall k \in Q, z_k \geq x_k$
- $\forall k \in Q \cap P, z_k \geq y_k$

Proof of (ii)

Let  $(P, y)$  be an objection of player  $i$  against player  $j$  to  $x$ .  $i \in P, j \notin P, y(P) \leq v(P)$  and  $y(P) > x(P)$ . We choose  $y(P) = v(P)$ .

- $x_j = v(\{j\})$ : Then  $(\{j\}, v(\{j\}))$  is a counter objection to  $(P, y)$ . ✓
- $x_j > v(\{j\})$ : Since  $x \in K(N, v, \mathcal{S})$  we have  $s_{ji}(x) \geq s_{ij}(x) \geq v(P) - x(P) \geq y(P) - x(P)$  since  $i \in P, j \notin P$ . Let  $Q \subseteq N$  such that  $j \in Q, i \notin Q$  and  $s_{ji}(x) = v(Q) - x(Q)$ . We have  $v(Q) - x(Q) \geq y(P) - x(P)$ . Then, we have

$$\begin{aligned} v(Q) &\geq y(P) + x(Q) - x(P) \\ &\geq y(P \cap Q) + y(P \setminus Q) + x(Q \setminus P) - x(P \setminus Q) \\ &> y(P \cap Q) + x(Q \setminus P) \text{ since } i \in P \setminus Q, y(P \setminus Q) > x(P \setminus Q) \end{aligned}$$

Let us define  $z$  as follows  $\begin{cases} x_k & \text{if } k \in Q \setminus P \\ y_k & \text{if } k \in Q \cap P \end{cases}$   
 $(Q, z)$  is a counter-objection to  $(P, y)$ . ✓

Finally  $x \in BS(N, v, \mathcal{S})$ .

Computing a kernel-stable payoff distribution

- There is a transfer scheme converging to an element in the kernel.
- It may require an infinite number of small steps.
- We can consider the  $\epsilon$ -kernel where the inequality are defined up to an arbitrary small constant  $\epsilon$ .

R. E. Stearns. **Convergent transfer schemes for n-person games.** *Transactions of the American Mathematical Society*, 1968.

Computing a kernel-stable payoff distribution

### Algorithm 1: Transfer scheme converging to a $\epsilon$ -Kernel-stable payoff distribution for the CS $\mathcal{S}$

```

compute- $\epsilon$ -Kernel-Stable( $N, v, \mathcal{S}, \epsilon$ )
repeat
  for each coalition  $C \in \mathcal{S}$  do
    for each member  $(i, j) \in C, i \neq j$  do // compute the maximum surplus
      // for two members of a coalition in  $\mathcal{S}$ 
       $s_{ij} \leftarrow \max_{R \subseteq N \setminus \{i, j\}} v(R) - x(R)$ 
     $\delta \leftarrow \max_{(i, j) \in C, i \neq j} s_{ij} - s_{ji}$ 
     $(i^*, j^*) \leftarrow \text{argmax}_{(i, j) \in C} (s_{ij} - s_{ji})$ 
    if  $(x_{i^*} - v(\{j^*\})) < \frac{\delta}{2}$  then // payment should be individually rational
       $d \leftarrow x_{i^*} - v(\{j^*\})$ 
    else
       $d \leftarrow \frac{\delta}{2}$ 
     $x_{i^*} \leftarrow x_{i^*} + d$ 
     $x_{j^*} \leftarrow x_{j^*} - d$ 
until  $\frac{\delta}{v(\mathcal{S})} \leq \epsilon$ 

```

- The complexity for one side-payment is  $O(n \cdot 2^n)$ .
- Upper bound for the number of iterations for converging to an element of the  $\epsilon$ -kernel:  $n \cdot \log_2(\frac{\delta_0}{\epsilon \cdot v(\mathcal{S})})$ , where  $\delta_0$  is the maximum surplus difference in the initial payoff distribution.
- To derive a polynomial algorithm, the number of coalitions must be bounded. For example, only consider coalitions which size is bounded in  $[K_1, K_2]$ . The complexity of the truncated algorithm is  $O(n^2 \cdot n_{\text{coalitions}})$  where  $n_{\text{coalitions}}$  is the number of coalitions with size in  $[K_1, K_2]$ , which is a polynomial of order  $K_2$ .

• M. Klusch and O. Shehory. **A polynomial kernel-oriented coalition algorithm for rational information agents.** In *Proceedings of the Second International Conference on Multi-Agent Systems*, 1996.

• O. Shehory and S. Kraus. **Feasible formation of coalitions among autonomous agents in non-superadditive environments.** *Computational Intelligence*, 1999.

## Summary

- We saw another way to use the excess to make objections and counter-objections.
- We defined the kernel.
- We proved that both the kernel and the bargaining set are non-empty if the set of imputations is non-empty.
- pros:**
  - If the set of imputations is non-empty, the nucleolus, kernel, bargaining set are non-empty.
  - There is an algorithm to compute a payoff in the kernel.
- cons:** The algorithm is not polynomial

## The Shapley value

Lloyd S. Shapley. **A Value for  $n$ -person Games.** In *Contributions to the Theory of Games, volume II (Annals of Mathematical Studies)*, 1953.

### Definition (marginal contribution)

The **marginal contribution** of agent  $i$  for a coalition  $\mathcal{C} \subseteq N \setminus \{i\}$  is  $mc_i(\mathcal{C}) = v(\mathcal{C} \cup \{i\}) - v(\mathcal{C})$ .

$(mc_1(\emptyset), mc_2(\{1\}), mc_3(\{1,2\}))$  is an efficient payoff distribution for any game  $((1,2,3), v)$ . This payoff distribution may model a dynamic process in which 1 starts a coalition, is joined by 2, and finally 3 joins the coalition  $\{1,2\}$ , and where the incoming agent gets its marginal contribution.

An agent's payoff depends on which agents are already in the coalition. This payoff may not be **fair**. To increase fairness, one could take the average marginal contribution over all possible joining orders.

Let  $\sigma$  represent a joining order of the grand coalition  $N$ , i.e.,  $\sigma$  is a permutation of  $(1, \dots, n)$ .

We write  $mc(\sigma) \in \mathbb{R}^n$  the payoff vector where agent  $i$  obtains  $mc_i(\{\sigma(j) \mid j < i\})$ . The vector  $mc$  is called a **marginal vector**.

### Shapley value: version based on marginal contributions

Let  $(N, v)$  be a TU game. Let  $\Pi(N)$  denote the set of all permutations of the sequence  $(1, \dots, n)$ .

$$Sh(N, v) = \frac{\sum_{\sigma \in \Pi(N)} mc(\sigma)}{n!}$$

the Shapley value is a **fair** payoff distribution based on marginal contributions of agents averaged over joining orders of the coalition.

### An example

$N = \{1, 2, 3\}$ ,  $v(\{1\}) = 0$ ,  $v(\{2\}) = 0$ ,  $v(\{3\}) = 0$ ,  
 $v(\{1, 2\}) = 90$ ,  $v(\{1, 3\}) = 80$ ,  $v(\{2, 3\}) = 70$ ,  
 $v(\{1, 2, 3\}) = 120$ .

	1	2	3	Let $y = \langle 50, 40, 30 \rangle$
$1 \leftarrow 2 \leftarrow 3$	0	90	30	$\mathcal{C}$
$1 \leftarrow 3 \leftarrow 2$	0	40	80	$e(\mathcal{C}, x)$
$2 \leftarrow 1 \leftarrow 3$	90	0	30	$e(\mathcal{C}, y)$
$2 \leftarrow 3 \leftarrow 1$	50	0	70	$\{1\}$
$3 \leftarrow 1 \leftarrow 2$	80	40	0	$\{2\}$
$3 \leftarrow 2 \leftarrow 1$	50	70	0	$\{3\}$
total	270	240	210	$\{1, 2\}$
Shapley value	45	40	35	$\{1, 3\}$
				$\{2, 3\}$
				$\{1, 2, 3\}$

This example shows that the Shapley value may not be in the core, and may not be the nucleolus.

- There are  $|\mathcal{C}|!$  permutations in which all members of  $\mathcal{C}$  precede  $i$ .
- There are  $|N \setminus (\mathcal{C} \cup \{i\})|!$  permutations in which the remaining members succeed  $i$ , i.e.  $(|N| - |\mathcal{C}| - 1)!$ .

The Shapley value  $Sh_i(N, v)$  of the TU game  $(N, v)$  for player  $i$  can also be written

$$Sh_i(N, v) = \sum_{\mathcal{C} \subseteq N \setminus \{i\}} \frac{|\mathcal{C}|!(|N| - |\mathcal{C}| - 1)!}{|N|!} (v(\mathcal{C} \cup \{i\}) - v(\mathcal{C})).$$

Using definition, the sum is over  $2^{|N|-1}$  instead of  $|N|!$ .

### Notion of value

#### Definition (value function)

Let  $\mathcal{G}_N$  the set of all TU games  $(N, v)$ . A **value function**  $\phi$  is a function that assigns to each TU game  $(N, v)$  an efficient allocation, i.e.  $\phi : \mathcal{G}_N \rightarrow \mathbb{R}^{|N|}$  such that  $\phi(N, v)(N) = v(N)$ .

- The Shapley value is a value function.
- None of the concepts presented thus far were a value function (the nucleolus is guaranteed to be non-empty only for games with a non-empty set of imputations).

### Some interesting properties

Let  $(N, v)$  and  $(N, u)$  be TU games and  $\phi$  be a value function.

- **Symmetry or substitution (SYM):** If  $\forall (i, j) \in N$ ,  $\forall \mathcal{C} \subseteq N \setminus \{i, j\}$ ,  $v(\mathcal{C} \cup \{i\}) = v(\mathcal{C} \cup \{j\})$  then  $\phi_i(N, v) = \phi_j(N, v)$
- **Dummy (DUM):** If  $\forall \mathcal{C} \subseteq N \setminus \{i\}$   $v(\mathcal{C}) = v(\mathcal{C} \cup \{i\})$ , then  $\phi_i(N, v) = 0$ .
- **Additivity (ADD):** Let  $(N, u + v)$  be a TU game defined by  $\forall \mathcal{C} \subseteq N$ ,  $(u + v)(N) = u(N) + v(N)$ .  $\phi(u + v) = \phi(u) + \phi(v)$ .

#### Theorem

The Shapley value is the unique value function  $\phi$  that satisfies (SYM), (DUM) and (ADD).

## Unanimity game

Let  $N$  be a set of agents and  $T \subseteq N \setminus \emptyset$ .

The **unanimity game**  $(N, v_T)$  is defined as follows:

$$\forall \mathcal{C} \subseteq N, v_T(\mathcal{C}) = \begin{cases} 1, & \text{if } T \subseteq \mathcal{C}, \\ 0 & \text{otherwise.} \end{cases}$$

We note that

- if  $i \in N \setminus T$ ,  $i$  is a null player.
- if  $(i, j) \in T^2$ ,  $i$  and  $j$  are substitutes.

### Lemma

The set  $\{v_T \mid T \subseteq N \setminus \emptyset\}$  is a linear basis of  $\mathcal{G}_N$ .

This means that a TU game  $(N, v)$  can be represented by a unique set of values  $(\alpha_T)_{T \subseteq N \setminus \emptyset}$  such that

$$\forall \mathcal{C} \subseteq N, v(\mathcal{C}) = \left( \sum_{T \subseteq N \setminus \emptyset} \alpha_T v_T \right) (\mathcal{C}).$$

## Proof of the lemma

There are  $2^n - 1$  unanimity games and the dimension of  $\mathcal{G}_N$  is also  $2^n - 1$ .

We only need to prove that the unanimity games are linearly independent.

Towards a contradiction, let us assume that  $\sum_{T \subseteq N \setminus \emptyset} \alpha_T v_T = 0$  where  $(\alpha_T)_{T \subseteq N \setminus \emptyset} \neq 0_{\mathbb{R}^{2^n - 1}}$ .

Let  $T_0$  be a minimal set in  $\{T \subseteq N \mid \alpha_T \neq 0\}$ .

Then,  $\left( \sum_{T \subseteq N \setminus \emptyset} \alpha_T v_T \right) (T_0) = \alpha_{T_0} \neq 0$ , which is a contradiction.

## Proof of the theorem: Uniqueness (1/2)

Let  $\phi$  a feasible solution on  $\mathcal{G}_N$  that is non-empty and satisfies the axioms SYM, DUM and ADD. Let us prove that  $\phi$  is a value function.

Let  $(N, v) \in \mathcal{G}_N$ .

- if  $v = 0_{\mathcal{G}_N}$ , all players are dummy. Since the solution is non-empty,  $0^{\mathbb{R}^{|N|}}$  is the unique member of  $\phi(N, v)$ .
- otherwise,  $(N, -v) \in \mathcal{G}_N$ .  
Let  $x \in \phi(N, v)$  and  $y \in \phi(N, -v)$ .  
By ADD,  $x + y \in \phi(v - v)$ , and then,  $x = -y$  is unique. Moreover,  $x(N) \leq v(N)$  as  $\phi$  is a feasible solution. Also  $y(N) \leq -v(N)$ .  
Since  $x = -y$ , we have  $v(N) \leq x(N) \leq v(N)$ , i.e.  $x$  is efficient.

Hence,  $\phi$  is a value function.

## Proof of the theorem: Uniqueness (2/2)

Let  $T \subseteq N \setminus \emptyset$  and  $\alpha \in \mathbb{R}$ . Let us prove that  $\phi(N, \alpha \cdot v_T)$  is uniquely defined.

- Let  $i \notin T$ . We have trivially  $T \subseteq \mathcal{C}$  iff  $T \subseteq \mathcal{C} \cup \{i\}$ . Then  $\forall \mathcal{C} \subseteq N \setminus \{i\}$ ,  $\alpha v_T(\mathcal{C}) = \alpha v_T(\mathcal{C} \cup \{i\})$ . Hence, all agent  $i \notin T$  are dummies. By DUM,  $\forall i \notin T$ ,  $\phi_i(N, \alpha \cdot v_T) = 0$ .
- Let  $(i, j) \in T^2$ . Then for all  $\mathcal{C} \subseteq N \setminus \{i, j\}$ ,  $v(\mathcal{C} \cup \{i\}) = v(\mathcal{C} \cup \{j\})$ . By SYM,  $\phi_i(N, \alpha \cdot v_T) = \phi_j(N, \alpha \cdot v_T)$ .
- Since  $\phi$  is a value function, it is efficient. Then,  $\sum_{i \in N} \phi_i(N, \alpha \cdot v_T) = \alpha v_T(N) = \alpha$ . Hence,  $\forall i \in T$ ,  $\phi_i(N, \alpha \cdot v_T) = \frac{\alpha}{|T|}$ .

This proves that  $\phi(N, \alpha \cdot v_T)$  is uniquely defined. Since any TU game  $(N, v)$  can be written as  $\sum_{T \subseteq N \setminus \emptyset} \alpha_T v_T$  and because of ADD, there is a unique value function that satisfies the three axioms.

## Proof of the theorem: Existence

We need to show that the Shapley value satisfies the three axioms. Let  $(N, v)$  a TU game.

- Let us assume that  $\forall \mathcal{C} \subseteq N \setminus \{i, j\}$ , we have  $v(\mathcal{C} \cup \{i\}) = v(\mathcal{C} \cup \{j\})$ . Then  $\forall \mathcal{C} \subseteq N \setminus \{i, j\}$ , we have
  - $mc_i(\mathcal{C}) = mc_j(\mathcal{C})$
  - $v(\mathcal{C} \cup \{i, j\}) - v(\mathcal{C} \cup \{i\}) = v(\mathcal{C} \cup \{i, j\}) - v(\mathcal{C} \cup \{j\})$ , hence, we have  $mc_j(\mathcal{C} \cup \{j\}) = mc_i(\mathcal{C} \cup \{i\})$ .
- $Sh_i(N, v) = Sh_j(N, v)$ ,  $Sh$  satisfies SYM.
- Let us assume there is an agent  $i$  such that for all  $\mathcal{C} \subseteq N \setminus \{i\}$  we have  $v(\mathcal{C}) = v(\mathcal{C} \cup \{i\})$ . Then, each marginal contribution of player  $i$  is zero, and it follows that  $Sh_i(N, v) = 0$ .  $Sh$  satisfies DUM.
- $Sh$  is clearly additive.

✓

## Coming next

- Voting games and power indices.