

Fine-grained complexity of coloring geometric intersection graphs

Édouard Bonnet joint work with Csaba Biró, Dániel Marx,
Tillmann Miltzow, and Paweł Rzażewski and Stéphan
Thomassé

Middlesex University, London

ACiD seminar, Durham, February 20th 2017

NP-hardness vs ETH-hardness

NP-hardness:

- ▶ your problem is not solvable in polynomial, unless 3-SAT is
- ▶ very widely believed but do not give evidence against algorithms running in say, $2^{n^{1/100}}$.

NP-hardness vs ETH-hardness


NP-hardness:

- ▶ your problem is not solvable in polynomial, unless 3-SAT is
- ▶ very widely believed but do not give evidence against algorithms running in say, $2^{n^{1/100}}$.

ETH-hardness:


- ▶ stronger assumption than $P \neq NP$ is ETH asserting that no $2^{o(n)}$ algorithm exists for 3-SAT
- ▶ Allows to prove stronger conditional lower bounds
- ▶ linear reduction from 3-SAT: no $2^{o(n)}$ algorithm for your problem, quadratic reduction: no $2^{o(\sqrt{n})}$ algorithm, etc.

Square root phenomenon on planar graphs




Many problems are solvable in $2^{O(\sqrt{n})}$ in **planar graphs**, and unlikely solvable in $2^{o(n)}$ in general graphs.

Square root phenomenon on planar graphs




Many problems are solvable in $2^{O(\sqrt{n})}$ in **planar graphs**, and unlikely solvable in $2^{o(n)}$ in general graphs.

Square root phenomenon on planar graphs




MAX INDEPENDENT SET, 3-COLORING, HAMILTONIAN PATH...

Square root phenomenon on planar graphs




MAX INDEPENDENT SET, 3-COLORING, HAMILTONIAN PATH...

Square root phenomenon on planar graphs




MAX INDEPENDENT SET, 3-COLORING, HAMILTONIAN PATH...

Square root phenomenon on planar graphs




MAX INDEPENDENT SET, 3-COLORING, HAMILTONIAN PATH...
Dynamic programming would spare a $\log n$ in the exponent.

Square root phenomenon on planar graphs




MAX INDEPENDENT SET, 3-COLORING, HAMILTONIAN PATH...

Square root phenomenon on planar graphs




MAX INDEPENDENT SET, 3-COLORING, HAMILTONIAN PATH...

Square root phenomenon on planar graphs




MAX INDEPENDENT SET, 3-COLORING, HAMILTONIAN PATH...

Coloring Unit Disks




It might also be that only the intersection graph is given and not a geometric representation.

Coloring Unit Disks




It might also be that only the intersection graph is given and not a geometric representation.

Coloring Unit Disks




It might also be that only the intersection graph is given and not a geometric representation.

Coloring Unit Disks




It might also be that only the intersection graph is given and not a geometric representation.

Coloring Unit Disks



It might also be that only the intersection graph is given and not a geometric representation.

Coloring Unit Disks



It might also be that only the intersection graph is given and not a geometric representation.

Balanced separators

Theorem (Smith, Wormald '98)

For every $d \geq 1$ and $B \geq 0$, there exists a constant $c = c(d, B)$, such that for every B -fat collection S of n d -dimensional convex sets with ply at most ℓ , there exists a d -dimensional sphere Q , such that:

- ▶ *at most $\frac{d+1}{d+2}n$ elements of S are entirely inside Q ,*
- ▶ *at most $\frac{d+1}{d+2}n$ elements of S are entirely outside Q ,*
- ▶ *at most $cn^{1-1/d}\ell^{1/d}$ elements of S intersect Q .*

Balanced separators

Theorem (Smith, Wormald '98)

For every $d \geq 1$ and $B \geq 0$, there exists a constant $c = c(d, B)$, such that for every B -fat collection S of n d -dimensional convex sets with ply at most ℓ , there exists a d -dimensional sphere Q , such that:

- ▶ at most $\frac{d+1}{d+2}n$ elements of S are entirely inside Q ,
- ▶ at most $\frac{d+1}{d+2}n$ elements of S are entirely outside Q ,
- ▶ at most $cn^{1-1/d}\ell^{1/d}$ elements of S intersect Q .

ply: maximum number of objects covering a point.

B -fat objects: aspect ratios diameter/width are bounded by B .



Balanced separators for unit disks

Theorem (Smith, Wormald '98, special case)

Given a collection S of n disks with ply at most ℓ , there exists a circle Q , such that:

- ▶ *at most $3n/4$ disks of S are entirely inside Q ,*
- ▶ *at most $3n/4$ disks of S are entirely outside Q ,*
- ▶ *at most $O(\sqrt{n\ell})$ disks of S intersect Q .*

Standard algorithm for ℓ -coloring (for unit disks)

If the ply is greater than ℓ , then more than ℓ colors are needed.

Otherwise, there is a balanced separator of size $O(\sqrt{n\ell})$ which can be exhaustively found in time $O(2^{\sqrt{n\ell} \log n})$.

Trying all the ℓ -colorings on S takes time $O(2^{\sqrt{n\ell} \log \ell})$.

Standard algorithm for ℓ -coloring (for unit disks)

If the ply is greater than ℓ , then more than ℓ colors are needed.

Otherwise, there is a balanced separator of size $O(\sqrt{n\ell})$ which can be exhaustively found in time $O(2^{\sqrt{n\ell} \log n})$.

Trying all the ℓ -colorings on S takes time $O(2^{\sqrt{n\ell} \log \ell})$.

Overall running time: $O(2^{\sqrt{n\ell} \log n})$.

We will see that this running time is optimal up to logarithmic factors in the exponent.

We will see that this running time is optimal up to logarithmic factors in the exponent.

Theorem

For any $\alpha \in [0, 1]$, coloring n unit disks with $\ell = \Theta(n^\alpha)$ colors cannot be solved in time $2^{o(n^{\frac{1+\alpha}{2}})} = 2^{o(\sqrt{n\ell})}$, under the ETH.

We will see that this running time is optimal up to logarithmic factors in the exponent.

Theorem

For any $\alpha \in [0, 1]$, coloring n unit disks with $\ell = \Theta(n^\alpha)$ colors cannot be solved in time $2^{o(n^{\frac{1+\alpha}{2}})} = 2^{o(\sqrt{n\ell})}$, under the ETH.

Constant number of colors \rightsquigarrow square root phenomenon.

Linear number of colors \rightsquigarrow no subexponential-time algorithm.

We will see that this running time is optimal up to logarithmic factors in the exponent.

Theorem

For any $\alpha \in [0, 1]$, coloring n unit disks with $\ell = \Theta(n^\alpha)$ colors cannot be solved in time $2^{o(n^{\frac{1+\alpha}{2}})} = 2^{o(\sqrt{n\ell})}$, under the ETH.

Constant number of colors \rightsquigarrow square root phenomenon.

Linear number of colors \rightsquigarrow no subexponential-time algorithm.


And everything in between (hard part).

For instance, \sqrt{n} -coloring cannot be done in $2^{o(n^{3/4})}$.

Roadmap

3-SAT \rightarrow 2-grid 3-SAT \rightarrow Partial 2-grid Coloring \rightarrow coloring unit disks

Partial 2-grid Coloring \rightarrow coloring unit disks



Partial 2-Grid Coloring

Input: An induced subgraph G of the $g \times g$ -grid, a positive integer ℓ . Each cell of this grid is mapped to a set of ℓ points (in a smaller grid $[\ell]^2$).

Question: Is there an ℓ -coloring of all the points such that:

- ▶ two points in the same cell get different colors;
- ▶ if v and w are adjacent in G , say, $w = v + (1, 0)$, p , resp. q , are points in the smaller grid of v resp. w , receiving the same color, then q has at a second coordinate which is at least the second coordinate of p ?


2-Grid 3-SAT

Input: A $g \times g$ grid, a positive integer k , each vertex (or cell) of the grid is associated to k variables, and a set \mathcal{C} of constraints of two kinds:

- ▶ **clause constraints:** for each cell of the grid, a set of pairwise variable-disjoint 3-clauses on its variables;
- ▶ **equality constraints:** for two adjacent cells of the grid, a set of pairwise variable-disjoint equality constraints.

Question: Is there an assignment of the variables such that all constraints are satisfied?

3-SAT \rightarrow 2-Grid 3-SAT




3-SAT on N variables with bounded number of occurrences (Sparsification Lemma) \rightsquigarrow split the clauses into $\approx g$ blocks \rightsquigarrow split again the clauses on one block into a constant number of sub-blocks (clauses vertex-disjoint)


The size of the created instance is $n = g^2 k$.

$$N = \Theta(gk) = \Theta(\sqrt{nk})$$



2-Grid 3-SAT \rightarrow Partial 2-Grid Coloring





2-Grid 3-SAT \rightarrow Partial 2-Grid Coloring




Encoding information and reference coloring




Wires




Permutation




Forget




Independence




Clauses




reference coloring



Consistency gadget (also crossing)





Higher dimension

Theorem

For $\alpha \in [0, 1]$ and dimension $d \geq 2$, coloring n unit d -balls with $\ell = \Theta(n^\alpha)$ colors cannot be solved in time $2^{n^{\frac{d-1+\alpha}{d}-\epsilon}}$ for any $\epsilon > 0$, under the ETH.



The first step in the chain is trickier: the higher dimensional grid should embed the SAT instance in a more compact way.

The second and third steps work similarly.

(Longer and longer) Segments

Theorem

6-coloring 2-Dir is not solvable in $2^{o(n)}$, under the ETH.




Reduction from 3-coloring on degree-4 graphs to list 6-coloring of segment intersection graphs.

The x_i 's lists are $[1, 2, 3]$, the y_j 's lists are $[4, 5, 6]$.


Circles are equality gadgets ($1 \equiv 4, 2 \equiv 5, 3 \equiv 6$), squares are inequality gadgets.

Equality




vertex	list
x_i	1,2,3
y_i	4,5,6
a_1	1,4
b_1	4,5
c_1	4,6
a_2	2,5
b_2	4,5
c_2	5,6
a_3	3,6
b_3	4,6
c_3	5,6

Inequality



vertex	list
x_j	1,2,3
y_j	4,5,6
x'	4,5,6
p_1	1,5
p_2	1,6
q_1	2,4
q_2	2,6
r_1	3,4
r_2	3,5

Inequality



vertex	list
x_j	1,2,3
y_j	4,5,6
x'	4,5,6
p_1	1,5
p_2	1,6
q_1	2,4
q_2	2,6
r_1	3,4
r_2	3,5

Some extra gadgets permit to remove the lists.

Same lower bound for 4 colors.

What happens with 3-colors? (whiteboard)

Thanks for your attention!