

ELECTRE TRI-nB, pseudo-disjunctive: axiomatic and combinatorial results

Denis Bouyssou¹ · Thierry Marchant² · Marc Pirlot³

Accepted: 13 March 2025 © The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2025

Abstract

ELECTRE TRI-nB is a method designed to sort alternatives evaluated on several attributes into ordered categories. It is an extension of ELECTRE TRI-B, using several limiting profiles, instead of just one, to delimit each category. ELECTRE TRI-nB comes in two flavours: pseudo-conjunctive and pseudo-disjunctive. In a previous paper, we have characterized the ordered partitions that can be obtained with ELECTRE TRI-nB, pseudo-conjunctive, using a simple axiom called linearity. The present paper is dedicated to the axiomatic analysis of ELECTRE TRI-nB, pseudo-disjunctive. It also provides some combinatorial results.

Keywords Multiple criteria analysis · Sorting models · ELECTRE TRI-nB

1 Introduction

Given a set of alternatives, multiple criteria decision analysis (MCDA, aka multiple criteria decision making) is a sub-field of operations research simultaneously dealing with several criteria or objectives over this set (for an overview of this field, see e.g. Belton & Stewart 2001, Greco et al., 2016). The goal is usually to choose a single most desirable alternative or to rank them by decreasing order of preference.

Authors are listed alphabetically. They have contributed equally.

Thierry Marchant thierry.marchant@UGent.be

> Denis Bouyssou dbouyssou@gmail.com

Marc Pirlot marc.pirlot@umons.ac.be

- ¹ CNRS, Paris, France
- ² Department of Data Analysis, Ghent University, H. Dunantlaan, 1, 9000 Ghent, Belgium

³ Université de Mons, rue de Houdain 9, 7000 Mons, Belgium

A problem with such an approach is that the best alternative may be a poor one, if all other alternatives are even poorer. That is why some researchers have proposed methods aiming at sorting the alternatives into a few ordered categories (e.g. good, passable and unacceptable) that are defined using some kind of norms or references. In sorting methods, the norms defining a category originally are alternatives that stand at the border of two consecutive categories¹. They are called *limiting profiles* For a recent survey of sorting methods, see Belahcène et al. (2023a, 2023b). See also the Special issue on multiple criteria sorting methods (Marchant & Pirlot, 2021). Notice that sorting models are studied not only in MCDA, but also in decision under risk (Nakamura, 2004) and decision under uncertainty (Bouyssou & Marchant, 2011).

An important difficulty when applying multiple criteria decision analysis is the choice of an adequate method (there are hundreds of them in the literature) and the choice—or elicitation—of parameters (often tens of them) that are reasonably compatible with the preferences of the decision-maker (for a survey about elicitation, see Dias et al., 2018). Since the 1960s, many papers and books (Bouyssou et al., 2006; Keeney & Raiffa, 1976) have shown that the axiomatic analysis of MCDA methods (for choosing or ranking) within the framework of Multi-attribute Utility Theory (Krantz et al., 1971) can provide valuable results for guiding the choice of the method and of the parameters. More recently, a thread of papers have done a similar job for MCDA methods aimed at sorting (e.g. Bouyssou & Marchant, 2007a; Słowiński et al., 2002).

ELECTRE TRI (or ETRI for short) is a family of sorting methods. The first method in this family was ETRI-B (Roy & Bouyssou, 1993; Yu, 1992). Then came several variants, that we do not detail.² The principle of all ELECTRE methods is to build a binary relation on the set of alternatives, which represents the preferences of the decision maker. Basically, an alternative is said to be preferred to another (or to "outrank" the other) if it is at least as good as the other on a sufficient coalition of criteria (concordance) and there is no criterion on which it is unacceptably worse than the other (non discordance). Such a relation, which is not necessarily transitive, is known as an outranking relation. In the ETRI-B sorting method, this relation is used to assign alternatives to categories by comparing them to limiting profiles. There are two versions of ETRI-B. The simpler one, called pseudo-conjunctive (aka pessimistic) assigns an alternative to a category if this alternative is preferred to (i.e., outranks) the lower limiting profile of the category and is not preferred to its upper limiting profile. The other variant, called pseudo-disjunctive (or optimistic), assigns an alternative to a category if the upper limiting profile of the category is strictly preferred to the alternative while the lower limiting profile of the category is not strictly preferred to it.

¹ Sorting methods have been recently proposed in which the norms defining a category stand central in that category

 $^{^{2}}$ For an overview of ELECTRE methods, we refer to Roy and Bouyssou (1993, Ch. 5 & 6), Figueira et al. (2013, 2016).

Recently, Fernández et al. (2017) proposed a new variant (named ELECTRE TRI-nB or ETRI-nB for short) that uses several limiting profiles instead of merely one as in the original ETRI-B. Like ETRI-B, the new ETRI-nB has two versions: pseudo-conjunctive (pc) and pseudo-disjunctive (pd).

A simplified version of ETRI-B-pc received a detailed axiomatic analysis in Bouyssou and Marchant (2007a, 2007b), Słowiński et al. (2002), Greco et al. (2001). Later, Bouyssou and Marchant (2015) have shown that ETRI-B-pd is much more difficult to analyze than ETRI-B-pc, although their definitions may seem dual to each other at first sight.

Bouyssou et al. (2023)—hereafter referred to as BMP23—have characterized the pseudo-conjunctive version of ETRI-nB making auxiliary use of a simplified version thereof. This characterization uses a single axiom—Linearity—that was first proposed by Goldstein (1991). It turns out that ETRI-nB-pc is a very general sorting model, if the number of limiting profiles is unbounded: any monotone assignment to ordered categories is representable in this model, using an appropriate number of limiting profiles. Bouyssou et al. (2022) have characterized the particular case of ETRI-nB using at most 2 limiting profiles.

The present paper intends to axiomatically analyze ETRI-nB-pd or a simplified version thereof. Is this model able to represent any monotone assignment rule as ETRI-nB-pc does, using an appropriate number of limiting profiles? Otherwise, what are the monotone rules that can be represented in this model? Our main findings are twofold. The first one is similar to that in Bouyssou and Marchant (2015): ETRI-nB-pd is much more difficult to analyze than ETRI-nB-pc, although their definitions may seem dual to each other. Only a subset of the monotone assignment rules can be represented in the pseudo-disjunctive model, but we are not able to characterize which ones do. Our second main finding is a characterization of a special case of ETRI-nB-pd, involving Linearity and a new condition, raising some interesting combinatorial questions about maximal antichains in direct products of chains.

2 Framework and notation

As in BMP23, we will restrict our attention to the case of two categories. This allows us to use a simple framework while not concealing any important difficulty.³ For the same reasons, we suppose throughout that the set of objects to be sorted is finite.

The finite set of alternatives is $X = X_1 \times ... \times X_n$, with $n \ge 2$. The set of attributes is $N = \{1, ..., n\}$. For $x, y \in X, i \in N$ and $J \subseteq N$, we use $X_J, X_{-J}, X_i, X_{-i}, (x_J, y_{-J})$ and (x_i, y_{-i}) as usual. Our primitives consist of a twofold partition $\langle \mathcal{A}, \mathcal{U} \rangle$ of the set X, where \mathcal{A} (resp. \mathcal{U}) contains the s<u>A</u>tisfactory (resp. <u>U</u>nsatisfactory) alternatives.

An attribute *i* is influential for $\langle \mathcal{A}, \mathcal{U} \rangle$ if there exist $x_i, y_i \in X_i$ and $a_{-i} \in X_{-i}$ such that $(x_i, a_{-i}) \in \mathcal{A}$ and $(y_i, a_{-i}) \in \mathcal{U}$. If an attribute is not influential, it does not play

³ Bouyssou and Marchant (2007b) have shown how to extend the axiomatic analysis to the case of more than two categories, in the case of ETRI-B. Their technique applies mutatis mutandis to ETRI-nB.

any role and can be suppressed. We therefore suppose without loss of generality that all attributes are influential.

3 Axiomatic analysis of ETRI-nB-pc: a digest

In this section, we recall some definitions and results presented in BMP23. All ETRI methods start with a preference modelling step during which a preference relation is built for each attribute. This valued preference relation depends on a number of parameters that we do not detail here. In a second step, these n valued preference relations are aggregated into a single valued preference relation that is afterwards cut to define a crisp outranking relation S. The assignment of alternatives to categories occurs in a third step. In order to save space, we do not present the exact definition of ETRI-nB-pc, but an idealization thereof: Model E. It mostly simplifies steps 1 and 2 and we will later see that this does not entail any loss of generality. See Fernández et al. (2017) for a complete description of ETRI-nB-pc and BMP23 for the relationship between Model E and ETRI-nB-pc.

Definition 1 (*Models* E, E^c, E^u) We say that a partition $\langle \mathcal{A}, \mathcal{U} \rangle$ has a representation in Model E if:

- for all *i* ∈ *N*, there is a semiorder *S_i* on *X_i* (with asymmetric part *P_i* and symmetric part *I_i*),
- for all *i* ∈ *N*, there is a strict semiorder *V_i* on *X_i* that is included in *P_i* and is the asymmetric part of a semiorder *U_i*,
- (S_i, U_i) is a homogeneous nested chain of semiorders and W_i is a weak order that is compatible with both S_i and U_i ,⁴
- there is a set of subsets of attributes $\mathcal{F} \subseteq 2^N$ such that, for all $I, J \in 2^N$, $[I \in \mathcal{F}]$ and $I \subseteq J] \Rightarrow J \in \mathcal{F}$,
- there is a binary relation S on X (with symmetric part I and asymmetric part P) defined by

$$x S y \iff [S(x, y) \in \mathcal{F} \text{ and } V(y, x) = \emptyset],$$

where $S(x, y) = \{i \in N : x_i S_i y_i\}$ and $V(x, y) = \{i \in N : x_i V_i y_i\}$,

• there is a set $\mathcal{P} = \{p^1, \dots, p^k\} \subseteq X$ of k limiting profiles, such that for all $p, q \in \mathcal{P}, Not[p P q],$

and

$$x \in \mathcal{A} \iff \begin{cases} x \, S \, p & \text{for some } p \in \mathcal{P} \\ Not[q \, P \, x] & \text{for all } q \in \mathcal{P}. \end{cases}$$
(1)

⁴ W_i is the intersection of the weak orders S_i^{wo} and U_i^{wo} , respectively induced by S_i and U_i . See Appendix A of the supplementary material of BMP23.

We then say that $\langle (S_i, V_i)_{i \in N}, \mathcal{F}, \mathcal{P} \rangle$ is a representation of $\langle \mathcal{A}, \mathcal{U} \rangle$ in Model *E*. Model *E*^{*c*} is the particular case of Model *E*, in which there is a representation that shows no discordance effects, i.e. in which all relations V_i are empty. Model E^u is the particular case of Model E^c , in which there is a representation that requires unanimity, i.e. such that $\mathcal{F} = \{N\}$.

In this definition, S_i is the idealization of the preference relation on attribute *i*, V_i represents all pairs of levels on attribute *i* for which a discordance could occur (step 1).⁵ S is the idealization of the outranking relation (step 2). The third step (the assignment of alternatives to categories) is described by (1).

Goldstein (1991) has proposed a simple condition that may be satisfied by some partitions:

Definition 2 (*Linearity*) The partition $\langle \mathcal{A}, \mathcal{U} \rangle$ is linear on attribute *i* if, for all $x_i, y_i \in X_i$ and all $a_{-i}, b_{-i} \in X_{-i}$,

$$\begin{array}{c} (x_i, a_{-i}) \in \mathcal{A} \\ \text{and} \\ (y_i, b_{-i}) \in \mathcal{A} \end{array} \right\} \Rightarrow \begin{cases} (y_i, a_{-i}) \in \mathcal{A} \\ \text{or} \\ (x_i, b_{-i}) \in \mathcal{A}. \end{cases}$$
(2)

The partition $\langle \mathcal{A}, \mathcal{U} \rangle$ is linear if it is linear on all attributes. If all partitions that can be represented in some Model *M* are linear, we say that Model *M* satisfies Linearity.

Replacing \mathcal{A} by \mathcal{U} in (2) yields an equivalent definition of Linearity. On each attribute X_i , we define the relation \succeq_i letting, for all $x_i, y_i \in X_i$,

$$x_i \gtrsim_i y_i$$
 if [for all $a_{-i} \in X_{-i}, (y_i, a_{-i}) \in \mathcal{A} \Rightarrow (x_i, a_{-i}) \in \mathcal{A}$].

By construction, \succeq_i is transitive and reflexive; it is complete if and only if the partition is linear on attribute *i*. The symmetric part of \succeq_i is denoted by \sim_i . It is not useful to keep in X_i elements that are equivalent w.r.t. the equivalence relation \sim_i . Indeed, if $x_i \sim y_i$ then $(x_i, a_{-i}) \in \mathcal{A}$ iff $(y_i, a_{-i}) \in \mathcal{A}$. In order to simplify notation, we suppose throughout the paper that we are dealing with partitions on $\prod_{i=1}^n X_i$ for which all relations \sim_i are trivial⁶. This non-restrictive convention implies that each relation \succeq_i is antisymmetric.

Let \succeq be the relation on *X* defined by $x \succeq y$ iff $x_i \succeq_i y_i$ for all $i \in N$. This relation is a partial order (reflexive, transitive and antisymmetric). Let $\mathcal{A}_* = \min(\mathcal{A}, \succeq)$ be the set of minimal elements in \mathcal{A} for \succeq . By construction, for any $x \in \mathcal{A}_*$ and $y_i \prec_i x_i$, we have $(y_i, x_{-i}) \in \mathcal{U}$.

⁵ In Definition 1, S_i and V_i are supposed to be semiorders. The reason of this assumption is that the notion of semiorder is related to the existence of thresholds, as they appear in the modelling of preference and veto in the classical ELECTRE methods.

⁶ If \sim_i is not trivial, we can work without loss of generality with the quotient X_i / \sim_i .

We say Model M is nested in—or is a special case of—Model M' (denoted $M \subseteq M'$) if all partitions that can be represented in M can also be represented in M'. Models M and M' are equivalent (denoted $M \equiv M'$) if $M \subseteq M'$ and $M' \subseteq M$. We note $M \subsetneq M'$ if $M \subseteq M'$ and M is not equivalent to M'. By construction, we have $E^u \subseteq E^c \subseteq E$. The main results in BMP23 can now be summarized in the following theorem.

Theorem 1

- 1. ETRI-nB-pc $\equiv E \equiv E^c \equiv E^u$.
- 2. A partition $\langle \mathcal{A}, \mathcal{U} \rangle$ has a representation in any of these models iff it is linear.
- 3. This representation can always be taken to be $\langle (\succeq_i, V_i = \emptyset)_{i \in \mathbb{N}}, \mathcal{F} = \{N\}, \mathcal{P} = \mathcal{A}_* \rangle$, that is a representation in Model E^u .

We like to stress point 1: although model E and the nested models E^c and E^u seem to be simplifications of ETRI-nB-pc, they are not: all four models are fully equivalent.

4 ETRI-nB-pd: definition and difficulties

The pseudo-disjunctive version of ETRI-nB consists of three steps. The first and the second one are identical to steps 1 and 2 in ETRI-nB-pc. The only difference is the third step: the assignment of alternatives to categories. With ETRI-nB-pc, an alternative x is assigned to \mathcal{A} iff it is weakly preferred (in terms of S) to a limiting profile and no limiting profile is strictly preferred to x (in terms of P), as in (1). With ETRI-nB-pd, an alternative x is assigned to \mathcal{U} iff (i) there is a limiting profile strictly preferred (in terms of P) to x and (ii) x is not strictly preferred to any limiting profile. As in Sect. 3, in order to save space, we do not present the exact definition of ETRI-nB-pd, but an idealization thereof: Model F. We define Model F that is to ETRI-nB-pd what Model E is to ETRI-nB-pc.

Definition 3 Model F is defined exactly as Model E, except that we now replace (1) by:

$$x \in \mathcal{U} \iff \begin{cases} p P x & \text{for some } p \in \mathcal{P} \text{ and} \\ Not[x P q] & \text{for all } q \in \mathcal{P}. \end{cases}$$
(3)

The definition of Models F^c and F^u parallels that of E^c and E^u .

All pseudo-disjunctive models mentioned so far satisfy linearity.

Lemma 1 If $\langle \mathcal{A}, \mathcal{U} \rangle$ has a representation in Model *F*, then it is linear. The same holds for ETRI-nB-pd, F^{u} and F^{c} .

Proof Consider first Model *F*. Suppose that we have $(x_i, a_{-i}) \in \mathcal{U}$ and $(y_i, b_{-i}) \in \mathcal{U}$. We have either $x_i W_i y_i$ or $y_i W_i x_i$ since W_i is a weak order. Suppose wlog that $y_i W_i x_i$. Because $(y_i, b_{-i}) \in \mathcal{U}$, we know that $p P (y_i, b_{-i})$, for some $p \in \mathcal{P}$, and $Not[(y_i, b_{-i}) P q]$ for all $q \in \mathcal{P}$. Using Lemma 3 in BMP23, we obtain $p P (x_i, b_{-i})$ and $Not[(x_i, b_{-i}) P q]$ for all $q \in \mathcal{P}$.⁷ Hence, $(x_i, b_{-i}) \in \mathcal{U}$ and linearity holds for Model *F*. By construction, $F^u \subseteq F^c \subseteq F$ and linearity thus also holds for these models.

Since we did not formally define ETRI-nB-pd, we cannot provide the proof that linearity holds for partitions generated by ETRI-nB-pd. For the interested reader, this proof closely follows that of Corollary 1 in BMP23.

Hence, combining Lemma 1 with Theorem 1, we obtain the next proposition.

Proposition 1 $F^u \subseteq F^c \subseteq F \subseteq E$ and ETRI-nB-pd $\subseteq E \equiv$ ETRI-nB-pc.

At this stage, given the apparent duality between the definitions of the pseudoconjunctive and pseudo-disjunctive models, we can suspect that $F^u \equiv F^c \equiv F \equiv$ ETRI-nB-pd $\equiv E$, but the next result shows that it does not hold.

Proposition 2 $F^u \subsetneq F^c$ and $F \subsetneq E$.

Proof Part 1 $F^u \subsetneq F^c$: Let $N = \{1, 2, 3\}$ and $X_i = \{0, 1\}$ for all $i \in N$, so that X has $2^3 = 8$ elements. Consider the partition $\langle \mathcal{A}, \mathcal{U} \rangle$ such that $\mathcal{A} = \{111, 101, 011\}$ and $\mathcal{U} = \{110, 100, 010, 001, 000\}$, abusing notation in an obvious way. It is simple to check that all attributes are influential for $\langle \mathcal{A}, \mathcal{U} \rangle$ and that, for all $i \in N$, we have $1_i \succ_i 0_i$. Notice that we have $\mathcal{A}_* = \text{Min}(\succeq, \mathcal{A}) = \{101, 011\}$ and $\mathcal{U}^* = \text{Max}(\succeq, \mathcal{U}) = \{110, 010, 001\}$.

Let us show that this partition *cannot* be obtained with Model F^u . Observe first that, here, since all attributes are influential and can only take two values, we must have that $S_i = \geq_i$, for all $i \in N$.

Since $110 \in \mathcal{U}$, there must be $p \in \mathcal{P}$ such that p P 110. Since we are looking for a representation in Model F^u and we know that $S_i = \succeq_i$, for all $i \in N$, we must find a profile $p \in \mathcal{P}$ such that p > 110. The only candidate is 111. But taking $\mathcal{P} = \{111\}$ together with $\mathcal{F} = \{N\}$ does not lead to the desired partition. Indeed, we have 111 > 101, so that 101 should be in \mathcal{U} .

This partition can be obtained with Model F^c , taking $S_i = \succeq_i$, for all $i \in N$, $\mathcal{P} = \{111\}$ and $\mathcal{F} = \{\{1,3\}, \{2,3\}\}.$

Part 2 $F \subsetneq E$: Let n = 4 and $X_1 = X_2 = X_3 = \{2, 1, 0\}$ and $X_4 = \{0, 1\}$, so that X has 54 elements. Consider the partition $\langle \mathcal{A}, \mathcal{U} \rangle$ such that $\mathcal{A} = \{2221, 2211, 2121, 1221, 2111, 1211, 1121, 1111, 2220\}$. Notice that $\mathcal{A}_* = \{1111, 2220\}$. It is easy to check that all attributes are influential for $\langle \mathcal{A}, \mathcal{U} \rangle$ and that, for all $i \in \{1, 2, 3\}$, we

⁷ Lemma 3 in BMP23 is established under the hypothesis that the partition $[\mathcal{A}, \mathcal{U}]$ is representable in Model E. Since the proof only uses the properties of relation *S*, which are common to Models E and F, the result also holds for partitions representable in Model F.

have $2_i >_i 1_i >_i 0_i$, while $1_4 >_4 0_4$. Hence, the partition is linear and, by Theorem 1, it can be represented in Model *E*.

In order to show that this partition cannot be obtained in Model F, we have to examine, all cases of indifference thresholds (associated with the strict semiorders S_i), combined with all cases of veto thresholds (associated with the strict semiorders V_i), and combined with all choices for \mathcal{F} .

Notice that if an attribute in $i \in \{1, 2, 3\}$ has thresholds (i.e. S_i is not a weak order), this means that $2_i I_i 1_i$ and $1_i I_i 0_i$. But veto effects can only occur among the elements that are strictly preferred. Hence, in this case, the only possibility is to take $2_i V_i 0_i$.

If $\{1, 2, 4\} \in \mathcal{F}$, then, without veto, $2201 \in \mathcal{U}$ outranks all elements in \mathcal{A} , a contradiction. This will remain true unless, there is a veto effect on attribute 3.

If $2_3 V_3 0_3$, the only elements in \mathcal{A} that are not strictly beaten by another element in \mathcal{A} are 2220, 1111, and 1121. It is easy to check that taking all of them or any subset of them as the set of profiles does not lead to the desired partition (consider $2201 \in \mathcal{U}$). If, furthermore, $2_3 V_3 1_3$, the only elements in \mathcal{A} that are not strictly beaten by another element in \mathcal{A} are 2220 and 1111. It is easy to check that taking all of them or any subset of them as the set of profiles does not lead to the desired partition (consider $2201 \in \mathcal{U}$).

The analysis of the cases $\{1, 3, 4\} \in \mathcal{F}$ and $\{2, 3, 4\} \in \mathcal{F}$ is entirely similar.

Suppose now that $\mathcal{F} = \{\{1, 2, 3\}, N\}$. Suppose that only attribute 1 has thresholds. Without veto, it is easy to check that $1220 \in \mathcal{U}$ outranks all elements in \mathcal{A} . This remains true, whatever the choice of veto thresholds on attributes 2 and 3. This also remains true if $1_4 V_4 0_4$. But veto effects on attribute 1 are immaterial since 1_1 is indifferent to both 2_1 and 0_1 .

The situation is entirely similar if 2 (resp. 3) is the only attribute to have thresholds.

Suppose that only attributes 1 and 2 have thresholds. Without veto, it is easy to check that $1120 \in \mathcal{U}$ outranks all elements in \mathcal{A} . This remains true, whatever the choice of veto thresholds on attributes 1 and 2 since 1_1 (resp. 1_2) is indifferent to 2_1 and 0_1 (resp. 2_2 and 0_2). This also remains true if $1_4 V_4 0_4$. Clearly, the veto threshold on attribute 3 is immaterial.

The analysis of the cases in which attributes (1 and 3) or (2 and 3) have thresholds is entirely similar.

It remains to tackle the case $\mathcal{F} = \{N\}$.

Suppose that only attribute 1 has thresholds. Without veto, there are only 3 elements in A that are not strictly beaten by another element in A: 2220, 1111 and 2111. It is easy to check that taking all of them or any subset of them as the set of profiles does not lead to the desired partition. It is simple to check that whatever the choice of veto we make on attributes 2, 3 and 4, the situation remains the same.

There is only one possibility to put a veto on attribute 1, i.e. $2_1 V_1 0_1$. In this case there are only 2 elements in A that are not strictly beaten by another element in A: 2220 and 1111. In any case, it is impossible to recover the desired partition.

The situation is entirely symmetric in the case only attribute 2 or only attribute 3 has thresholds.

Suppose that both attributes 1 and 2 have thresholds. Without veto, there are only 5 elements in \mathcal{A} that are not strictly beaten by another element in \mathcal{A} : 2220, 1111, 2211, 2111, and 1211. It is easy to check that taking all of them or any subset of them as the set of profiles does not lead to the desired partition. It is simple to check that whatever the choice of veto we make on attributes 3 and 4, the situation remains the same. There is only one possibility to put a veto on attribute 1 (resp. 2), i.e. $2_1 V_1 0_1$ (resp. $2_2 V_2 0_2$). It is simple to check that any of the three possible choices for the veto on these attributes does not alter the situation.

The situation is entirely symmetric in the case only attributes 1 and 3 or only attributes 2 and 3 have thresholds.

Finally, if all attributes have thresholds, there is only one element in \mathcal{A} that is not strictly beaten by another element in \mathcal{A} : 2220. It is easy to check that taking this element to be the unique profile, does not lead to the desired partition. Now, the choice of veto thresholds (they must be of the type $2_i V_i 0_i$) on attributes 1, 2, and 3 is immaterial. But it is also simple to check that adding a veto on attribute 4 does not change the situation.

Given Propositions 1 and 2, it would be highly desirable to know whether ETRInB-pd $\equiv E$ or ETRI-nB-pd $\equiv F$. Unfortunately, we are presently unable to prove or disprove these equivalences. This shows that the relations between the pseudodisjunctive models are more complex than between the corresponding pseudoconjunctive models.

5 Two characterizations

In view of the above-metioned difficulties, we devote this section to two simpler problems: (1) the characterization of Models F^c , F, E and ETRI-nB-pc when all attributes are binary and (2) the characterization of a special case of Model F^u .

5.1 The case of binary attributes

Suppose the partition $\langle \mathcal{A}, \mathcal{U} \rangle$ is linear on attribute *i*. We say attribute *i* is binary if the weak order \gtrsim_i has exactly two equivalence classes. Such attributes are common in many applications. The case in which all attributes are binary corresponds to the well-developed theory of monotone Boolean functions (see Crama and Hammer, 2011).

Proposition 3 $F^c \equiv F \equiv E \equiv \text{ETRI-nB-pc}$ whenever all attributes are binary.

Proof When all attributes are binary, each X_i contains only two elements that we can denote by 1_i and 0_i with $1_i \succ_i 0_i$. Each element in *X* corresponds to a unique coalition $C(x) = \{i \in N : x_i = 1_i\} \subseteq 2^N$. Hence, all linear partitions have a representation in F^c with $S_i = \succeq_i$, for all $i \in N$, $\mathcal{P} = \{111\}$ and $\mathcal{F} = \{C(x) : x \in \mathcal{A}\}$.

Since the proof uses a set \mathcal{P} containing only one limiting profile, the reader may have the impression that Proposition 3 only applies to ETRI-B and not to ETRI-nB. What the result actually says is that any partition generated by a model F^c , F, E or ETRI-nB-pc (irrespective of the number of limiting profiles) can be represented in the other three models. The proof further shows that the representation can be chosen so that \mathcal{P} is a singleton.

5.2 A special case of Model F^u

In order to reduce the complexity of the models, let us assume that the data are of good quality—in the sense of Roy (1996, Section 8.2)—meaning that there is no imprecision, uncertainty, or inaccurate determination. In that case, there is no need to use preference or indifference thresholds and the relation S_i is a weak order. Since \gtrsim_i is also a weak order and we cannot have $x_i >_i y_i$ while $y_i S_i x_i$, it must be the case that S_i is a refinement of \gtrsim_i (i.e. $S_i \subseteq \gtrsim_i$). But since we have assumed that the relation \sim_i is trivial, the equality $S_i = \gtrsim_i$ must hold.

So, in this section, we restrict our attention to partitions having a representation in Model F^u such that $S_i = \succeq_i$ is a weak order for all $i \in N$. Model F^u together with this additional constraint will be denoted by $F^{\underline{u}}$. In such a model, $S = \succeq$, $P = \succ$ and Condition 3 reduces to $x \in \mathcal{U}$ iff $p \succ x$, for some $p \in \mathcal{P}$. Indeed, we may not have $p \succ x \succ q$ for $q \in \mathcal{P}$, otherwise $p \succ q$, a contradiction. Notice that $F^{\underline{u}} \subseteq F^u \subsetneq F^c \subseteq F \subsetneq E$.

By construction, the set $\mathcal{A}_* = \operatorname{Min}(\succeq, \mathcal{A})$ is an antichain in the poset (X, \succeq) , remembering our convention that each relation \sim_i is trivial. Observe that in the first part of the proof of Proposition 2, the antichain $\mathcal{A}_* = \{101, 011\}$ is not a maximal antichain, i.e. it is strictly included in the antichain $\{110, 101, 011\}$. As shown below, a characteristic feature of partitions that can be represented in Model $F^{\underline{\mu}}$ is that \mathcal{A}_* is a maximal antichain in the poset (X, \succeq) .

Theorem 2 Let $X = \prod_{i=1}^{n} X_i$ be a finite set and $\langle \mathcal{A}, \mathcal{U} \rangle$ be a twofold linear partition of *X*. The partition $\langle \mathcal{A}, \mathcal{U} \rangle$ has a representation in Model $F^{\underline{u}}$ iff the antichain \mathcal{A}_* , in the poset (X, \geq) , is maximal.

Proof Necessity. Suppose that \mathcal{A}_* is not a maximal antichain. Hence there is $x \in X$ such that x is incomparable, using \succeq , w.r.t. all elements in \mathcal{A}_* . In view of the definition of \mathcal{A}_* , it is impossible that $x \in \mathcal{A}$ (since this would imply that $x \succeq z$, for some $z \in \mathcal{A}_*$). Hence, we must have $x \in \mathcal{U}$, so that there must be a profile $p \in \mathcal{P}$ such that p > x. This profile must be in \mathcal{A} . But, by hypothesis, this profile cannot belong to \mathcal{A}_* . Hence, by construction, we know that p > y, for some $y \in \mathcal{A}_* \subseteq \mathcal{A}$, which implies $y \in \mathcal{U}$, a contradiction.

Sufficiency. Since $\langle \mathcal{A}, \mathcal{U} \rangle$ is linear, we know that it has a representation in Model E^u using the representation $\langle (\succeq_i, V_i = \emptyset)_{i \in N}, \mathcal{F} = \{N\}, \mathcal{P} = \mathcal{A}_* \rangle$. Since \mathcal{A}_* is a maximal antichain, it is easy to see that this representation is also a representation in Model $F^{\underline{u}}$. Indeed, by construction, it is impossible that $x \in \mathcal{U}$ is incomparable, using \succeq , to all $p \in \mathcal{P} = \mathcal{A}_*$. Let $q \in \mathcal{P}$ be such that x and q are comparable using

 \gtrsim . It is impossible that *x* ≥ *q* since this would imply that *x* ∈ A, in view of the definition of $A_* = P$. Hence, we must have that *q* > *x*.

The next result shows that $F^{\underline{u}} \subsetneq F^{u}$, thereby showing that the hypothesis that the representation is such that $S_i = \succeq_i$, for all $i \in N$, is *not* innocuous.

Proposition 4 $F^{\underline{u}} \subsetneq F^{\underline{u}}$.

Proof Let $N = \{1, 2, 3, 4\}$ and $X_i = \{0, 1, 2\}$ for all $i \in N$, so that X has $3^4 = 81$ elements. Consider the partition $\langle \mathcal{A}, \mathcal{U} \rangle$ such that $\mathcal{A} = \{2222, 2221, 2220, 2212, 2211, 2210, 2202, 2201, 2200, 2122, 2121, 2120, 2112, 2111, 2110, 2102, 2101, 2022, 2021, 2020, 2012, 2011, 2010, 2002, 2001, 1222, 1221, 1220, 1212, 1211, 1210, 1202, 1201, 1200, 1122, 1121, 1120, 1112, 1111, 1110, 1102, 1101, 1022, 0222, 0221, 0220, 0212, 0211, 0210, 0202, 0201, 0122, 0121, 0120, 0112, 0111, 0110, 0102, 0101, 0022\}$. The set \mathcal{A} has 60 elements. It is easy to check that we have $\mathcal{A}_* = \{2010, 2001, 1200, 0110, 0101, 0022\}$.

We have $2012 \in \mathcal{A}$, $1012 \in \mathcal{U}$, $1200 \in \mathcal{A}$, $0200 \in \mathcal{U}$, so that $2_1 >_1 1_1 >_1 0_1$. Similarly, we have: $2200 \in \mathcal{A}$, $2100 \in \mathcal{U}$, $1101 \in \mathcal{A}$, $1001 \in \mathcal{U}$, so that $2_2 >_2 1_2 >_2 0_2$. We also have: $0022 \in \mathcal{A}$, $0012 \in \mathcal{U}$, $2110 \in \mathcal{A}$, $2100 \in \mathcal{U}$, so that $2_3 >_3 1_3 >_3 0_3$. Finally, we have: $0022 \in \mathcal{A}$, $0021 \in \mathcal{U}$, $2101 \in \mathcal{A}$, $2100 \in \mathcal{U}$, so that $2_4 >_4 1_4 >_4 0_4$ (notice that the role of attributes 3 and 4 is entirely symmetric, in this example).

Hence, using Theorem 2, this partition *cannot* be represented in Model $F^{\underline{u}}$. Indeed, the antichain \mathcal{A}_* is not maximal: the element 2100 is incomparable, using \geq , to all elements in \mathcal{A}_* .

Yet it is cumbersome but easy to check that this partition can be obtained in Model F^{u} , taking $\mathcal{P} = \{2200, 0022\}, \mathcal{F} = \{N\}, S_{i} = \succeq_{i}$, for i = 2, 3, 4, and $2_{1} P_{1} 0_{1}$, $2_{1} I_{1} 1_{1}$, and $1_{1} I_{1} 0_{1}$.

Let us define $E^{\underline{u}}$ in the same way as $F^{\underline{u}}$. By Theorem 1, $E^{\underline{u}}$ is equivalent to E^{u} . Summarizing Proposition 4 and previous results, we have that $F^{\underline{u}} \subsetneq F^{u} \subsetneq F^{c} \subseteq F \subsetneq E \equiv E^{c} \equiv E^{u} \equiv E^{\underline{u}}$. This long chain of inclusions and equivalences illustrates the strong asymmetry between the families of pseudoconjunctive and pseudo-disjunctive models. In order to explore the gap between both families, we devote the rest of the paper to comparing the numbers of partitions that can be represented in models $F^{\underline{u}}$ and $E^{\underline{u}}$ (or any of the pseudo-conjunctive models discussed in this paper). This will help us quantify how restrictive $F^{\underline{u}}$ is compared to $E^{\underline{u}}$.

6 Counting maximal antichains

The number of partitions that can be represented in model $F^{\underline{u}}$ (resp. model $E^{\underline{u}}$) is the number of maximal antichains (resp. antichains) in the poset (X, \geq) . This poset can be seen as a direct product of *n* chains, where *n* is the number of attributes and the

*i*th chain $(i \in \{1, ..., n\})$ is the set $[m_i] = \{1, ..., m_i\}$ ordered by \geq (the natural order on the integer interval $[m_i]$), with m_i being the number of equivalence classes of the weak order \gtrsim_i . Notice that antichains in the direct product of *n* chains also plays an important role in the analysis of multichoice cooperative games, as shown by Grabisch (2016a). More generally, the importance of studying discrete mathematics structures in decision theory was powerfully stressed in Grabisch (2016b).

The number of antichains (maximal antichains) in $[m_1] \times \ldots \times [m_n]$ will be denoted by $d_E(m_1, \ldots, m_n)$ (resp. $d_F(m_1, \ldots, m_n)$). When $m_1 = \ldots = m_n = m$, the numbers $d_E(m_1, \ldots, m_n)$ and $d_F(m_1, \ldots, m_n)$ are respectively denoted by $D_E(m, n)$ and $D_F(m, n)$. We first tackle two special cases (n = 2 and m = 2) and then the general case, for which we have few results.

6.1 The case *n* = 2

Let \mathbb{N} denote the set of positive integers. The next result, due to Covington (2004), presents a recurrence relation for $d_F(m_1, m_2)$.

Theorem 3 For all $m_1, m_2 \in \mathbb{N}$, $d_F(m_1, m_2)$ is equal to

$$d_F(m_1 - 1, m_2 - 1) + \sum_{i=0}^{m_1 - 2} d_F(i, m_2 - 1) + \sum_{i=0}^{m_2 - 2} d_F(m_1 - 1, i).$$
(4)

A detailed proof of this result can be found in Bouyssou et al. (2024). For $d_E(m_1, m_2)$, the following result easily follows from Berman and Köhler (1976).

Corollary 1 For all $m_1, m_2 \in \mathbb{N}$, we have

$$d_E(m_1, m_2) = \begin{pmatrix} m_1 + m_2 \\ m_1 \end{pmatrix}.$$

Proof According to Berman and Köhler (1976), the number of antichains in $[m_1] \times [m_2] \times [m_3]$ is equal to

$$\prod_{i=0}^{m_3-1} \frac{\binom{m_1+m_2+i}{m_1}}{\binom{m_1+i}{m_1}}.$$
(5)

Setting $m_3 = 1$ in this expression yields the desired result.

For illustration purpose, we computed some numerical results under the constraint that $m_1 = m_2$ (to save space). Some terms of the sequences $D_E(m, 2)$ and $D_F(m, 2)$ can be found in Table 1, with the corresponding ratios $D_F(m, 2)/D_E(m, 2)$. For small values of *m*, the difference of expressivity between models $F^{\underline{u}}$ and $E^{\underline{u}}$ is not very large, but it grows for large values of *m*, since the ratio seems to converge to 0.

Table 1 Number $D_F(m, 2)$ of maximal antichains, number $D_E(m, 2)$ of antichains and ratio of these numbers in $[m]^2$ for $m \in [15]$ and $m = 100$. Values of $D_F(m, 2)$ are computed by means of (4)	m	$D_F(m,2)$	$D_E(m,2)$	$D_F(m,2)/D_E(m,2)$
	1	1	2	0.5
	2	3	6	0.5
	3	9	20	0.45
	4	27	70	0.385714286
	5	83	252	0.329365079
	6	259	924	0.28030303
	7	817	3432	0.238053613
	8	2599	12870	0.201942502
	9	8323	48620	0.171184698
	10	26797	184756	0.145039945
	11	86659	705432	0.122845292
	12	281287	2704156	0.104020256
	13	915907	10400600	0.088062900
	14	2990383	40116600	0.074542284
	15	9786369	155117520	0.06309003
	100	3.76527E+51	9.05485E+58	4.15829E-08

Table 2 Number $D_F(2, n)$ of
maximal antichains, number
$D_E(2, n)$ of antichains and ratio
of these numbers in $[2]^n$ for
$n \in [7]$

n	$D_F(2,n)$	$D_E(2,n)$	$D_F(2,n)/D_E(2,n)$
1	2	3	0.6666667
2	3	6	0.5
3	7	20	0.35
4	29	168	0.172619
5	376	7581	0.04959768
6	31746	7828354	0.004055259
7	123805914	2414682040998	0.00005127214

 $D_F(m, 2)$ and $D_F(m, 2)$ are respectively sequences A171155 and A000984 in the On-line Encyclopedia of Integer Sequences OEIS (2023). A recurrence relation is mentioned by Alois P. Heinz (without proof) for $D_F(m, 2)$ in OEIS (2023): $D_F(m, 2)$ is equal to

$$\frac{(4m-3)D_F(m-1,2) - (2m-5)D_F(m-2,2) + D_F(m-3,2) - (m-3)D_F(m-4,2)}{m}$$

Some other results (old and new) about the case n = 2 are presented in Bouyssou et al. (2024). Therein, in addition to enumeration results, correspondences (bijections) between (maximal) antichains in products of chains and other mathematical structures are established.

6.2 The case *m* = 2

 $D_F(2, n)$ is sequence A326358 in OEIS (2023). No expression seems to be known for this sequence and the highest known value corresponds to n = 7. Some terms can be found in Table 2.

 $D_E(2, n)$ corresponds to the Dedekind numbers (sequence A000372 in OEIS (2023)), for which no expression is known. The highest known value corresponds to n = 9. Some terms can be found in Table 2 with the corresponding ratios $D_F(2, n)/D_E(2, n)$. Here again, for small values of n, the difference of expressivity between models $F^{\underline{u}}$ and $E^{\underline{u}}$ is not very large, but for large values of n, the ratio seems to converge to 0.

6.3 The general case

In the general case, analytic expressions for $D_F(m, n)$ and $D_E(m, n)$ are difficult to obtain and we therefore only provide a lower bound for $D_F(m, n)$ and some numerical results.

6.3.1 A lower bound for $D_F(m, n)$

Proposition 5 The number of maximal antichains in $[m]^n$ is at least the number of antichains of $[m]^{n-1}$, that is $D_F(m, n) \ge D_F(m, n-1)$.

Proof The set $\{x \in [m]^n : x_i = m\}$ is the set of elements $x \in X$ having their *i*th coordinate equal to *m*. We shall prove that any antichain, not necessarily maximal, in $\{x \in [m]^n : x_i = m\}$ can be extended into a maximal antichain of *X*, which has no other element with its *i*th coordinate x_i equal to *m*. This will establish Proposition 5 since any antichain of X_{-i} is in one-to-one correspondence with an antichain of $\{x \in [m]^n : x_i = m\}$.

We take wlog i = 1. If the antichain in $\{x \in [m]^n : x_1 = m\}$ is maximal, the result is obvious. Otherwise, let A be any non-maximal antichain in $\{x \in [m]^n : x_1 = m\}$. Since A is not maximal in $\{x \in [m]^n : x_1 = m\}$, there is at least one element $x = (m, x_2, ..., x_n)$ that is incomparable to all elements in A. Let $x' = (m - 1, x_2, ..., x_n)$. We have that x > x' and x' is incomparable to any element in A. Indeed, for no $y \in A$, we have $x' \gtrsim y$ (otherwise $x \gtrsim y$ would hold too) and, for no $y \in A$, we have $y \gtrsim x'$ (otherwise $y \gtrsim x$ would also hold). Consider the set of all elements in $\{x \in [m]^n : x_1 = m\}$ that are incomparable to all elements in A. Select the minimal elements from this set. Change the first coordinate of each minimal element x into $x_1 = m - 1$, yielding an element x'. Let A' be the set obtained by adding all such elements x' to the antichain A. These elements are incomparable to all elements in A and incomparable to one another. Therefore, A' is an antichain. It is easy to see that it is maximal in X. Furthermore, the intersection of A' with the set $\{x \in [m]^n : x_1 = m\}$ is exactly A. Since $D_F(m, n)$ is nondecreasing with *m* and *n*, we may conclude in particular that the number of maximal antichains in *X* is at least the number of antichains in [2]^{*n*-1}, which is Dedekind number $D_E(2, n-1)$. Table 2 suggests that this bound is very weak. It also suggests that $D_F(m, n)$ grows extremely fast with *n* even for m = 2.

6.3.2 Some numerical results

Table 3 presents some values of $D_F(m, n)$ for small values of m and n, computed with the help of the software system Macaulay2 (Grayson & Stillman, 2021). For [3]³, we used the function maximalAntichains provided by the package Posets in the software system Macaulay2 (Grayson & Stillman, 2021) and manually checked the result. For [4]³ and [3]⁴, we also used the function maximalAntichains, but without manual check. For larger values (except when m = 2 or n = 2), the calculations are prohibitively long (indicated by question marks in Table 3).

For [3]³, using (5), we find $D_E(3,3) = 980$ so that the ratio $D_F(3,3)/D_E(3,3)$ is equal to 0.14693878. Similarly, for [4]³, we obtain $D_E(4,3) = 232848$ so that the ratio $D_F(4,3)/D_E(4,3)$ is equal to 0.04565639, which implies a huge difference of expressivity between $F^{\underline{u}}$ and $E^{\underline{u}}$.

7 Conclusion

We provided characterizations for two special cases of ETRI-nB-pd. Theses cases are not extremely restrictive and definitely correspond to applications. For these two special cases, our results can be used to develop elicitation techniques for the parameters of the method.

In addition, although our results about ETRI-nB-pd and its special cases are partial, we have axiomatic and combinatorial results showing that

- 1. the analysis of the pseudo-disjunctive models is far more complex than that of the pseudo-conjunctive models;
- 2. there is a whole variety of pseudo-disjunctive models that are not all equivalent, contrary to what we observed for pseudo-conjunctive models;
- most pseudo-disjunctive models are strict special cases of the corresponding pseudo-conjunctive models;

Table 3 Number of maximal	$D_{-}(m, n)$	n-1	2	3	4
antichains $(D_F(m, n))$ for small values of <i>m</i> and <i>n</i> . Boldface entries are new	$ \frac{D_F(m,n)}{m=1} $	n = 1 1 2 3 4 5	2 1 3 9 27	1 7 144 10631	4 1 29 116547 ?
	5	5	83	?	?

4. the pseudo-disjunctive model $F^{\underline{u}}$ is much more restrictive than the corresponding pseudo-conjunctive model.

The strong asymmetry between the pseudo-conjunctive and pseudo-disjunctive models can be ascribed to the central role played by the relation P in the definition of ETRI-nB-pd while S is central in ETRI-nB-pc. Indeed, Bouyssou and Pirlot (2015a, 2015b) have shown that the nature of the relation P is rather different from that of the relation S in the ELECTRE methods.

Hence, paralleling Bouyssou and Marchant (2015), we suggest to define the *dual* of ETRI-nB-pc not by means of (3), but rather by

$$x \in \mathcal{U} \iff \begin{cases} p \ S \ x & \text{for some } p \in \mathcal{P} \\ Not[x \ P \ q] & \text{for all } q \in \mathcal{P}. \end{cases}$$
(6)

It is easy to see that ETRI-nB-pc and its dual now correspond via the transposition operation consisting in inverting the direction of preference on all criteria and permuting A and U (see Almeida-Dias, Figueira, and Roy, 2010, Bouyssou and Marchant, 2015, Roy, 2002).

Mimicking Bouyssou et al. (2023, Th. 15), it is clear this dual model is characterized by Linearity. Instead of taking \mathcal{A}_* as the set of profiles to delimit \mathcal{A} , we now take $\mathcal{U}^* = \text{Max}(\succeq, \mathcal{U})$ to delimit the category \mathcal{U} , still using $S_i = \succeq_i$ and $\mathcal{F} = \{N\}$.

If we replace (3) by (6) in the definition of Models F, F^c and F^u , it is also simple to see that they are all equivalent to the dual of ETRI-nB-pc.

Declarations

Conflict of interest The authors have no relevant financial or non-financial interests to disclose.

References

- Almeida-Dias, J., Figueira, J. R., & Roy, B. (2010). ELECTRE TRI-C: A multiple criteria sorting method based on characteristic reference actions. *European Journal of Operational Research*, 204(3), 565–580.
- Belahcène, K., Mousseau, V., Ouerdane, W., Pirlot, M., & Sobrie, O. (2023a). Multiple criteria sorting models and methods—part I: Survey of the literature. 40R, 21(1), 1–46.
- Belahcène, K., Mousseau, V., Ouerdane, W., Pirlot, M., & Sobrie, O. (2023b). Multiple criteria sorting models and methods. Part II: Theoretical results and general issues. 4OR, 21(2), 181–204.
- Belton, V., & Stewart, T. (2001). Multiple criteria decision analysis: An integrated approach. Kluwer.
- Berman, J., & Köhler, P. (1976). Cardinalities of finite distributive lattices. Mitteilungen aus dem Mathematischen Seminar Giessen, 121, 103–124.
- Bouyssou, D., & Marchant, T. (2007a). An axiomatic approach to noncompensatory sorting methods in MCDM, I: The case of two categories. *European Journal of Operational Research*, 178(1), 217–245.
- Bouyssou, D., & Marchant, T. (2007b). An axiomatic approach to noncompensatory sorting methods in MCDM, II: More than two categories. *European Journal of Operational Research*, 178(1), 246–276.

- Bouyssou, D., & Marchant, T. (2011). Subjective expected utility without preferences. Journal of Mathematical Psychology, 55(6), 457–468. https://doi.org/10.1016/j.jmp.2011.08.006
- Bouyssou, D., & Marchant, T. (2015). On the relation between ELECTRE TRI-B and ELECTRE TRI-C and on a new variant of ELECTRE TRI-B. *European Journal of Operational Research*, 242, 201–211.
- Bouyssou, D., Marchant, T., Pirlot, M., Tsoukiàs, A., & Vincke, P. (2006). Evaluation and decision models with multiple criteria: Stepping stones for the analyst. Springer.
- Bouyssou, D., Marchant, T., & Pirlot, M. (2022). A note on ELECTRE TRI-nB with few limiting profiles. 40R, 20, 443–463. https://doi.org/10.1007/s10288-021-00485-y. Published online on June 17, (2021).
- Bouyssou, D., Marchant, T., & Pirlot, M. (2023). A theoretical look at ELECTRE TRI-nB and related sorting models. 4OR, A Quarterly Journal of Operations Research, 21, 73–103.
- Bouyssou, D., Marchant, T., & Pirlot, M. (2024). About maximal antichains in a product of two chains: A catch-all note, https://arxiv.org/abs/2410.16243.
- Bouyssou, D., & Pirlot, M. (2015a). A consolidated approach to the axiomatization of outranking relations: A survey and new results. Annals of Operations Research, 229(1), 159–212. https://doi.org/ 10.1007/s10479-015-1803-y
- Bouyssou, D., & Pirlot, M. (2015b). A note on the asymmetric part of an outranking relation. *Interna*tional Transactions in Operational Research, 22(5), 883–912. https://doi.org/10.1111/itor.12135
- Covington, M. A. (2004). The number of distinct alignments of two strings. *Journal of Quantitative Linguistics*, 11(3), 173–182.
- Crama, Y., & Hammer, P. L. (2011). Boolean functions: Theory, algorithms, and applications. Encyclopedia of mathematics and its applications. Cambridge University Press.
- Dias, L. C., Morton, A., Quigley, J. (2018). *Elicitation : The Science and Art of Structuring Judgement*. International Series in Operations Research & Management Science, 261. Springer International Publishing, Cham, 1st ed. 2018. edition, ISBN 3-319-65052-1.
- Fernández, E., Figueira, J. R., Navarro, J., & Roy, B. (2017). ELECTRE TRI-nB: A new multiple criteria ordinal classification method. *European Journal of Operational Research*, 263(1), 214–224. https:// doi.org/10.1016/j.ejor.2017.04.048
- Figueira, J. R., Greco, S., Roy, B., & Słowiński, R. (2013). An overview of ELECTRE methods and their recent extensions. *Journal of Multi-Criteria Decision Analysis*, 20(1–2), 61–85.
- Figueira, J. R., Mousseau, V., & Roy, B. (2005). ELECTRE methods. In J. R. Figueira, S. Greco, and M. Ehrgott, editors, *Multiple Criteria Decision Analysis: State of the Art Surveys*, number 233 in International Series in Operations Research & Management Science, pages 155–185. Springer, New York, NY, 2016. ISBN 978-1-4939-3094-4. https://doi.org/10.1007/978-1-4939-3094-4_5. First Edition:
- Goldstein, W. M. (1991). Decomposable threshold models. *Journal of Mathematical Psychology*, 35(1), 64–79.
- Grabisch, M. (2016). Remarkable polyhedra related to set functions, games and capacities. TOP, 24, 301– 326. https://doi.org/10.1007/s11750-016-0421-4
- Grabisch, Michel. (2016). Set functions, games and capacities in decision making (Vol. 46). Springer. https://doi.org/10.1007/978-3-319-30690-2_2 978-3-319-30690-2.
- Grayson, D. R., & Stillman, M. E. (2021). Macaulay2, a software system for research in algebraic geometry. https://macaulay2.com/. accessed on Apr 8, 2025
- Greco, S., Ehrgott, M., Figueira, J.R. (2005). Multiple Criteria Decision Analysis: State of the Art Surveys. Number 233 in International Series in Operations Research & Management Science. Springer, New York, NY, 2016. ISBN 978-1-4939-3094-4. First Edition:
- Greco, S., Matarazzo, B., & Słowiński, R. (2001). Conjoint measurement and rough set approach for multicriteria sorting problems in presence of ordinal criteria. Aide Multicritère à la Décision / Multiple Criteria Decision AidIn A. Colorni, M. Paruccini, & B. Roy (Eds.), A-MCD-A (pp. 117–144). European Commission, Joint Research Centre.
- Keeney, R. L., & Raiffa, H. (1976). Decisions with multiple objectives: Preferences and value tradeoffs. Wiley.
- Krantz, D. H., Luce, R. D., Suppes, P., & Tversky, A. (1971). Foundations of measurement, volume 1: Additive and polynomial representations. Academic Press.
- Marchant, T., & Pirlot, M. (2021). eds. Journal of Multi-Criteria Decision Analysis, Special issue on multiple criteria sorting methods, volume 28.

- Nakamura, Y. (2004). Trichotomic preferences for gambles. Journal of Mathematical Psychology, 48(6), 385–398.
- OEIS. (2023). The On-line Encyclopaedia of Integer Sequences, Sloane, N. J. A. (Ed.). https://oeis.org. Accessed on Apr 8, 2025
- Roy, B. (2002). Présentation et interprétation de la méthode ELECTRE TRI pour affecter des zones dans des catégories de risque. Document du LAMSADE 124, LAMSADE, Université Paris Dauphine, Paris, France, 25 pages.
- Roy, B., & Bouyssou, D. (1993). Aide multicritère à la décision:méthodes et cas. Paris:Economica., 2-7178-2473-1
- Roy, B., & Multicriteria methodology for decision aiding. Kluwer, Dordrecht, (1996). Original version in French: "Méthodologie multicritère d'aide à la décision" (p. 1985). Economica.
- Słowiński, R., Greco, S., & Matarazzo, B. (2002). Axiomatization of utility, outranking and decision-rule preference models for multiple-criteria classification problems under partial inconsistency with the dominance principle. *Control and Cybernetics*, 31(4), 1005–1035.
- Yu, W. (1992). Aide multicritère à la décision dans le cadre de la problématique du tri:concepts, méthodes et applications. Thèse de doctorat: Université Paris Dauphine, Paris, France. (in French).

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.